

Quality of Life of Cognitively Impaired Older Adults Living in Community

Shailendra Mohan Tripathi¹, Rakesh Kumar Tripathi²

¹Assistant Professor, Department of Geriatric Mental Health, King George's Medical University UP, Lucknow, India.

²Assistant Professor cum Clinical Psychologist, Department of Geriatric Mental Health, King George's Medical University UP, Lucknow, India

Abstract: *Quality of life (QOL) conveys an overall sense of well-being, including aspects of happiness and satisfaction with life as a whole. With graying scenario worldwide, knowing about the QOL is important to developing future strategies for enhancing the same. The study aimed at finding QOL among community dwelling urban older adults from Lucknow. Two randomly selected urban localities (Musahibganj & Jankipuram) of Lucknow were visited for data collection in a consecutive series. Out of 7351 homes visited and 1888 elderlies were identified. Of them the first 300 consecutively recruited persons fulfilling inclusion & exclusion criteria constituted the study sample. They were screened by HCST (Hindi Cognitive Screening Test) in such a way that 75 persons from either of the sexes with HCST positive (Score less than or equal to 23) and Negative scores (Score more than 23) scores were considered for analysis. World Health Organization Quality of Life-BREF (WHOQOL-BREF) was administered to assess QOL. The results revealed that there was significant difference between QOL of normal and cognitively impaired older adults. QOL in physical, psychological and social domains were found to be significantly poor amongst cognitively impaired individuals.*

Keywords: *Cognitive impairment, HCST, Quality of life, WHOQOL-BREF, older adults*

1. Introduction

QOL is extraordinarily broad and conceptually complex multidimensional construct, with some legitimate controversies regarding its definition. Quality of life (QOL) is a popular term that conveys an overall sense of well-being, including aspects of happiness and satisfaction with life as a whole. It is broad and subjective rather than specific and objective. Apart from health the other domains of QOL are finance, housing, security education, and the neighborhood (Tripathi, 2012). It is also affected by culture, values, and spirituality which add to the complexity of its measurement. Nevertheless, researchers in the fields of psychology and sociology have developed useful techniques that have helped to conceptualize and measure these multiple domains and how they relate to each other. (Center for Disease Control and Prevention, 2000). The World Health Organization (WHO) has attempted defining the concept and to developing appropriate assessment tools (WHOQOL Group, 1998, 1999). Developers of these tools have attempted to incorporate both subjective and objective elements of QOL by identifying and testing various facets of quality of life and examining it in cross-cultural studies.

H.S. Asthana (2009) from India in his work proposed that, the term quality of life (QOL) is an amorphous concept; it is normative and value laden. Its contours are difficult to be defined and its empirical inquiry is still more challenging. QOL overflows the boundaries of social sciences including economics and psychology. One must take into account what humanistic disciplines have to say in this regard (QOL) and examine the issue in the context of the world view one holds.

In a study by Tiwari, Tripathi and Kumar (2011) statistically significant difference was found between normally aging subjects and subjects with mental health problems in QOL

of urban elderlies. Most of the normally aging subjects (67.7%) were significantly highly satisfied with their QOL when compared to cognitively impaired (12.6%) and subjects with psychiatric disorders (10.6%). Significantly higher number of subjects (46.7%) suffering with psychiatric disorders reported dissatisfaction from their QOL when compared to subjects having cognitive disorder (13.3%).

Plethora of studies on QOL of elderly people living in the community and old age homes (Hall *et al.* 2011; Akbar *et al.* 2013; 2014); suffering from different physical (Phillips-Bute *et al.* 2006) and mental illnesses (Selwood, Thorgrimsen & Orrell, 2005; Bartels and Pratt, 2009) have been conducted. Some reported issues regarding health related QOL and others reported issues regarding assessment (Hall *et al.* 2011) and administration of tools specially in elderly patient with severe cognitive impairment or with dementia (Logsdon *et al.* 2002; De Rooij *et al.* 2011; Schifczczyk *et al.* 2010). Bartels and Pratt (2009) found that poor functional outcomes and lower quality of life among older people with severe mental illnesses are strongly associated with social isolation, depression, cognitive impairment, and chronic medical illness.

Scattered studies on QOL in psychiatric illnesses, such as in schizophrenia (Nanda, Das & Bhalla, 2002), patients with severe mental illness (Jha *et al.* 2003), in adolescent bipolar affective disorder (Kumari, 2006), QOL on residents of old age home (Mehra *et al.* 2005; Akbar *et al.* 2014) have been conducted in India. There is hardly any study about QOL amongst community dwelling elderlies in India. Relationship between cognitive functioning and quality of life has also not thoroughly been researched in India though, there is some study about cognitive functioning of community dwelling older adults (Tripathi & Tiwari, 2011; 2013; Tiwari *et al.* 2012). Studying this relation may be important since it may provide a window of opportunity for

therapeutic intervention in the near future. Present study is an effort in this direction.

Keeping this in mind the study was planned to delineate the association of QOL with cognitive functioning and the QOL of elderly which was part of doctoral thesis entitled, "A Clinical Psychological study of cognitive functioning as a determinant of quality of life amongst urban elderly". (Tripathi & Tiwari, 2012). The thesis work was conducted along with ICMR funded project titled: 'Lucknow urban elderly epidemiological project' (Tiwari *et al.* 2009). The present study is adopted from the thesis to compare the QOL of normal and cognitively impaired community dwelling older adults.

2. Methodology

Institutional ethical clearance was taken for the main study on which data this paper is based. It was a community based survey under aegis of ICMR sponsored Lucknow urban elderly project (Tiwari *et al.* 2009, 2014) in which a team of trained social workers and clinical psychologists with established inter rater reliability visited two randomly selected urban localities (Musahibganj & Jankipuram) of Lucknow. Out of 7351 homes visited 1888 elderly were identified who consented participating in the present study.

2.1 Sample

Of the total sample of persons screened the first 300 consecutively recruited persons fulfilling inclusion & exclusion criteria constituted the study sample. They were screened by Hindi Cognitive Screening Test (Tiwari & Tripathi, 2011) in such a way that 75 persons from either of the sexes with HCST positive (Score less than or equal to 23) and Negative scores (Score more than 23) scores were considered for analysis. (Table 1)

Table 1: Sex and cognitive status (HCST positive and negative) wise distribution of the sample:

Classification of Elderlies by Sex	HCST Positive (score at or below 23)	HCST Negative (score above 23)	Total
Males (60 years and above)	75	75	150
Females (60years and above)	75	75	150
Total	150	150	300

Inclusion Criteria

1. Elderly (Males and Females) aged 60 years and above; Confirmation of the age was done using one or more of the following:
 - i. Government document; Retirement year (if retired)
 - ii. Year of marriage + gap period of his/her eldest child birth + age of eldest son/daughter = Age of the subject
 - iii. Age at independence year (1947) of India respect to freedom of India + duration between year 1947 to date of screening/interview = Age of the subject
2. Cooperative persons
3. Persons giving written informed consent (by the elderly / their family members)

Exclusion Criteria:

1. Uncooperative persons;
2. Having been diagnosed as suffering from any psychiatric disorder (as per ICD-10 criteria) other than Mild Cognitive Impairment and Dementia. The findings of the diagnosis from the ICMR funded research project entitled, "An epidemiological study of prevalence of neuropsychiatric disorders with special reference to cognitive disorders amongst urban elderly" (Tiwari *et al.* 2009b) were adopted for making the diagnoses. Having any indicators of significant organic pathology like head injury, seizure, mental retardation, substance abuse etc. or having significant physical ailments.
3. Having problems with speech, hearing and vision, which can impede the interview.

Assessment Tools:

Following tools were administered for the assessment:

Semi-structured socio-demographic data sheet: Semi structured socio-demographic data sheet was used to collect the information about social, personal and demographic details of the subjects.

Hindi Cognitive Screening Test (HCST): Tiwari and Tripathi (2011) developed a cognitive screening instrument in which items suited to both literate and illiterate subjects and could be interchanged depending upon the literacy, MMSE and HMSE. HCST has a high level of sensitivity (0.93), specificity (0.96), and high positive (0.96), and negative (0.94) predictive value when compared to BCRS. A significant ($p < 0.01$) negative correlation ($r = -0.87$) with BCRS total scores and different axes of BCRS was found for concentration ($r = -0.79$), recent memory ($r = -0.83$), past memory ($r = -.79$), Orientation ($r = -.73$), and functioning / self care ($r = -.77$).

World Health Organization Quality of Life-BREF (WHOQOL-BREF) Hindi version (Saxena *et al.* 1998): The WHOQOL-BREF, an abbreviated 26 item version of the WHOQOL-100, was developed using data from the field-trial version of the WHOQOL-100. The items are distributed into 4 domains (physical; psychological; social and environmental health) and 25 facets. WHOQOL-BREF has been shown to have good discriminant validity, content validity and test retest validity. Question number 1 and 2 of the WHOQOL-BREF is related to overall Quality of life and general health respectively as perceived by the individual.

WHOQOL-BREF is a five point rating scale questionnaire which consists of scoring as follows 1: very poor; 2: poor; 3: neither poor nor good; 4: good and 5: very good. It is subjective rating of QOL of the participants on different domains.

The following cut off scores for the Total QOL scores was used in the study to delineate the level of quality of life.

Level of Quality of life	QOL-BREF total Score
Very poor	1-26
Poor	27-52
Average	53-78
Good	79-104
Very good	105-130

3. Procedure of the Study

Information about the subjects was obtained on semi-structured Socio-demographic proforma. HCST (Tiwari and Tripathi, 2011) was administered on included subjects to identify subjects with Cognitive Impairment. A cut off score of 24 was applied to differentiate cognitively impaired subjects from normal (Table 1). WHOQOL-BREF was administered on all 300 subjects. As and when required, the family members were cross checked about the information given by the elderly subjects at the time of administration of tests. The information which was difficult to elicit from the subjects with cognitive impairment were obtained from their family members who were living with the elderly subjects for at least last one year.

Statistical Analysis: Data was analyzed using statistical software Statistical Packages for Social Sciences (SPSS) version 12.0. and GraphPad InStat demo version 3.05 Inc year 2000. Mean, SD and Chi square test was used for analysis. Fisher's exact test was used for variables with two categories only. A minimum cut off value of $p < 0.05$ was considered as statistically significant.

4. Results

Table 2: Cognitive status wise distribution of QOL of urban older adults

QOL	Cognitive status on HCST	Very Poor N (%)	Poor N (%)	Average N (%)	Good N (%)	Very good	Total	Pearson Chi square
Physical QOL	Cognitive Impairment	8 (5.3)	35(23.3)	63 (42.0)	41 (27.3)	3 (2.0)	150	111.825 P<.0001 Significant
	Normal	1(0.7)	1 (0.7)	28 (18.7)	52 (34.7)	68 (45.3)	150	
	Total	9 (3.0)	36 (12.0)	91 (30.3)	93 (31.0)	71 (23.7)	300	
Psychological QOL	Cognitive Impairment	4 (2.7)	23 (15.3)	96 (64.0)	27 (18.0)	0 (0.0)	150	184.084 P<.0001 Significant
	Normal	0 (0.0)	0 (0.0)	10 (6.7)	76 (50.7)	64 (42.7)	150	
	Total	4 (1.3)	23 (7.7)	106 (35.3)	103 (34.3)	64 (21.3)	300	
Social QOL	Cognitive Impairment	3 (0.5)	21 (14.0)	105 (70.0)	21 (14.0)	0 (0.0)	150	104.955 P<.0001 Significant
	Normal	0 (0.0)	1 (0.7)	44 (29.3)	84 (56.0)	21 (14.0)	150	
	Total	3 (1.0)	22 (7.3)	149 (49.7)	105 (35.0)	21 (7.0)	300	
Environmental QOL	Cognitive Impairment	2 (1.3)	25 (16.7)	85 (56.7)	38 (25.3)	0 (0.0)	150	17.249 P<.140 not Significant
	Normal	0 (0.0)	0 (0.0)	26 (17.3)	102 (68.0)	22 (14.7)	150	
	Total	2 (0.7)	25 (8.3)	111 (37.0)	140 (46.7)	22 (7.3)	300	

5. Discussion

The study was conducted with the objective to study the QOL of cognitively impaired older adults aged 60 years and above residing in urban locality of Lucknow, Uttar Pradesh, India. Statistically significant ($p < 0.0001$) difference on QOL was found between normal (HCST negative) and cognitively impaired (HCST positive) older adults in physical, psychological and social domains of WHOQOL-BREF. QOL on all these domains were found to be poor amongst cognitively impaired older adults. A similar finding that is poor QOL amongst cognitively impaired older adults were

The mean age of the older adults was 67.98 ± 8.7 years and there was significant difference ($p, 0.001$) between mean age of normal (65.2 ± 6.6 years) and cognitively impaired (70.7 ± 9.6) group. Most of the older adults were illiterate (normal- 74.7% and cognitively impaired- 65.7%), married (normal- 66% and cognitively impaired- 58%) followed by widowed (34% & 40.7%) in both the groups.

Table 2 shows that significantly higher number of older adults from HCST negative (normal) group reported very good (45.3%) QOL on 'physical' domain, when compared to cognitively impaired (2.0%) on HCST. However, significantly higher percentage of cognitively impaired older adults reported average (42.0%), followed by poor (32.3%) and very poor (2.7%) QOL in 'physical' domain when compared to HCST negative older adults.

On 'psychological' domain significantly higher percentage of normal older adults on HCST reported good (50.7%) and very good (42.7%) QOL when compared to cognitively impaired older adults (18.0% and 0% respectively). However this pattern was found to be reverse amongst normal and cognitively impaired older adults.

Similar results were found in 'social' domain where significantly higher number of older adults from HCST negative (normal) group reported good (56.0%) and very good (14%) QOL as compared to HCST positive (cognitively impaired) group where significantly higher number of cognitively impaired older adults reported average (70.0%), poor (14.0%) and very poor (0.5%) 'Social' relationships when compared to normal older adults.

reported by Bartels and Pratt (2009) Tiwari *et al.* (2011) and B arrrios *et al.* (2013).

There was insignificant difference in environmental domain between normal and cognitively impaired subjects in our study. It was in accordance with the findings of Abrahamson *et al.* (2012) where more severe cognitively impaired subjects reported higher QOL in the domains of comfort and environment and lower QOL in activities, individuality, privacy and meaningful relationships, and the mood scale. It may be explained on the basis of thought process prevalent in the Indian society that older adults should given shelter inside the home and their daily basic needs to be addressed

irrespective of whether they are normal or having some illness. The other domains are affected in older adults with cognitive impairment as they require individual's effort and somehow it lacks due to symptomatology of the disease.

Our study validates the findings of Logsdon et al. (2002) where they reported poor QOL of cognitively impaired subjects but their emphasis was more on to study whether assessment of QOL in severely cognitively impaired individuals is possible from the subject itself or it would be better to rate on the basis of information provided by caregiver. They found that cognitively impaired elderly can rate their QOL with the progression of disease. WHOQOL-BREF was administered on all the subjects in our study and there were only 2 subjects where information provided was required confirmation of family members.

In our study mean age of the cognitively impaired group was found to be significantly higher than normal's. Poor QOL of cognitively impaired is thus associated with increasing age. This might be due to the fact that with advancing age older adults weaken biologically, physically, psychologically and economically and are not able to cope with their physical disabilities. Similar findings were reported by Gureje *et al.* (2008) for QOL. Joshi *et al.* (2003) also found that health related QOL was associated with advancing age. It is obvious that with advancing age there is progression in cognitive impairment and thus it affects the QOL of older adults. But if, psycho-social and physical wellbeing of older adults can be improved then maximum utilization of remaining cognitive function will be done thus, QOL of the older adults.

6. Conclusion

The results demonstrate that QOL of cognitively impaired older adults was found to be significantly poor when compared to normal older adults. It was more pronounced in the areas of physical, psychological, and social domain of QOL.

7. Acknowledgement

Authors are grateful to Indian Council of Medical Research, New Delhi for funding support, Dr S.C. Tiwari., Professor & Head, Department of Geriatric Mental Health and Dr Aditya Kumar for guidance, Mrs Reema Sinha, Reetu Shukla, Mrs. Sandhya Rani and Dr. Samridhi Tandon for data collection and compilation. The paper was presented in 39th National Annual Conference of Indian Association of Clinical Psychologists (NACIACP), AIIMS, New Delhi, 18th to 20th Feb. 2013 by 2nd author.

References:

- [1] Abrahamson, K., Clark, D., Perkins, A., Arling, G. (2012). Does Cognitive Impairment Influence Quality of Life Among Nursing Home Residents? *The Gerontologist*, 1-9. doi: 10.1093/geront/gnr137
- [2] Akbar, S., Tiwari, S.C., Garg, R.K., Pandey, N.M., Tripathi, R.K., Kumar A. (2013). Psychiatric morbidity

- and quality of life of inmates in old age homes. *Indian Journal of Geriatric Mental Health*, 9 (suppl 1), 18.
- [3] Akbar, S., Tiwari, S.C., Tripathi, R.K., Pandey, N.M., Kumar, A. (2014). Psychiatric morbidity and quality of life of residents in old age homes in Bareilly district of Uttar Pradesh. *International Journal of Psychosocial Research*, 3 (1):126-130.
- [4] Asthana, H.S. (2009). Quality of life: a psychological analysis. *Psychological Studies*, 54, 109-113.
- [5] Bárrios, H., Narciso, S., Guerreiro, M., Maroco, J., Logsdon, R., de Mendonça, A. (2013). Quality of life in patients with mild cognitive impairment. *Aging Ment Health*; 17(3):287-92. doi: 10.1080/13607863.2012.747083.
- [6] Bartels, S.J. and Pratt, S.I. (2009). Psychosocial Rehabilitation and Quality of Life for Older Adults With Serious Mental Illness: Recent Findings and Future Research Directions. *Curr Opin Psychiatry*, 22(4), 396-400.
- [7] Centers for Disease Control and Prevention. (2000). *Measuring Healthy Days*. US Department of Health and Human Services. Atlanta, Georgia: CDC, November 2000.
- [8] De Rooij, A.H.P.M., Luijckx, K. G., Declercq, A. G., & Schols, J.M.G.A. (2011). Quality of life of residents with dementia in long term care settings in the Netherlands and Belgium: design of a longitudinal comparative study in traditional nursing homes and small scale living facilities. *BMC Geriatrics*, 11, 20. Retrieved from <http://www.biomedcentral.com/1471-2318/11/20>
- [9] Gureje, O., Kola, L., Afolabi, E., & Olley, B.O. (2008). Determinants of quality of life of elderly Nigerians: results from the Ibadan study of ageing. *Afr J Med Med Sci.*, 37 (3), 329. UKPMC Funders Group Author Manuscript; available in PMC 2010 February 12.
- [10] Hall, S., Opio, D., Dodd, R.H., and Higginson, I.J. (2011). Assessing quality-of-life in older people in care homes. *Age and Ageing*, 40 (4), 507-512.
- [11] Jha, M., Singh, A.R., & Verma, A.N. (2003). Quality of life for patients with severe mental illness. *Industrial Psychiatry Journal*, 12(1), 7-9.
- [12] Joshi, K., Avasthi, A., & Kumar, R. (2003b). Health related quality of life among the elderly in northern India. *Health and population –perspectives and Issues*, 26 (4), 141-153.
- [13] Kumari, S. (2006). Quality of life and social support in adolescent bipolar affective disorder (Laxmi Shanti Award). *Industrial Psychiatry Journal*, 15(1),14 -18.
- [14] Logsdon, R.G., Gibbons, L.E., McCurry, S.M., & Teri, L. (2002). Assessing quality of life in older adults with cognitive impairment. *Psychosomatic Medicine*, 64(3),510-519.
- [15] Mehra, H.D., Kasar, N., Waykar, V., Dhadphale, M. (2005). Evaluation of Quality of life (QOL) functional and socio-demographic status of residents of old age home. *Industrial Psychiatry Journal*, 14 (1-2), 32-34.
- [16] Nanda, P.K., Das P.K., & Bhalla, K.C. (2002): Quality of life and job stress in schizophrenics working in a steel industry. *Industrial Journal of Psychiatry*, 11(1), 39-41.
- [17] Orley, J., Saxena, S. & Herrman, H. (1998). *Quality of life and mental illness. Reflections from the perspective*

- of the WHOQOL. *British Journal of Psychiatry*, 172, 291-293.
- [18] Phillips-Bute, B., Mathew, J.P., Blumenthal, J.A., Grocott, H.P., Laskowitz, D.T., Jones, R.H., Mark, D.B., and Newman, M.F. (2006) Association of Neurocognitive Function and Quality of Life 1 Year After Coronary Artery Bypass Graft (CABG) Surgery. *Psychosomatic Medicine*, 68,369-375.
- [19] Saxena, S., Chandiramani, K., and Bhargava, R.(1998). WHOQOL-Hindi : A questionnaire for assessing quality of life in health care settings in India. *The National Medical Journal of India*,11(4), 160-65.
- [20] Schifczyk, C., Romero, B., Jonas, C., Lahmeyer, C., Müller, F. and Riepe, M.W. (2010). Generic quality of life assessment in dementia patients: a prospective cohort study. *BMC Neurology*, 10, 48.
- [21] Selwood, A., Thørgrensen, L. and Orrell, M. (2005). Quality of life in dementia—a one-year follow-up study. *International Journal of Geriatric Psychiatry*, 20, 232-237. doi: 10.1002/gps.1271
- [22] Sivasilam, S., and Gopalakrishnan, G. (2006). A comparative study on the quality of life (QOL) and activities in daily (ADL) among non-cognitively impaired (MMSE>24) and cognitively impaired (MMSE ≤ 24) elderly persons from an urban based institutional care setting for elderly. *Indian Journal of Geriatric Mental Health*, 2 (1), 37-46.
- [23] Tiwari S C, Kar A M, Singh R, Kohli V K, Agarwal GG (2009). An epidemiological study of prevalence of neuro-psychiatric disorders with special reference to cognitive disorders amongst urban elderly. Report submitted to Indian Council of Medical Research, New Delhi, India.
- [24] Tiwari, S., & Tripathi, R. (2009). Neuropsychiatric morbidity amongst urban and rural older adults in Northern India: Report from community based epidemiological study. *International Psychogeriatrics*, 21, supplement 2, S136.
- [25] Tiwari, S.C. & Tripathi, R.K. (2011). “Development of an Education and Culture Fair Hindi Cognitive Screening Test (HCST) for the Elderly Population of India”. IAGMH / INTAS award-2011 at 3rd place. *Indian Journal of Geriatric Mental Health*, 7(2), 83-96.
- [26] Tiwari, S.C., Tripathi, R.K., Kumar A (2011). Predictors of healthy and unhealthy aging in north Indian Older adults with reference to mental health. IAGMH / INTAS award-2010 at 3rd place in 6th Annual National Conference of IAGMH held on 4th-5th Sep. 2010, Mumbai. *Indian Journal of Geriatric Mental Health*, 7 (1); 24-35.
- [27] Tiwari, S.C., Tripathi, R.K., Kumar, A., Kar A.M., Singh, R., Kohli V.K., Agarwal, G.G. (2014) Prevalence of Psychiatric Morbidity amongst Urban Older adults: Lucknow Elderly Study. *Indian Journal of Psychiatry*, 56 (2), 154-160.
- [28] Tiwari, S.C., Tripathi, R.K., Farooqi, S.A., Kumar, R., Srivastava, G., Kumar, A (2012). Diabetes Mellitus: A risk factor of cognitive impairment. *Industrial Psychiatry Journal*, 21 (1), 44-48. DOI: 10.4103/0972-6748.110950
- [29] Tripathi R K & Tiwari S. C. (2012). PhD Thesis submitted to CSMMU titled “A Clinical Psychological study of cognitive functioning as a determinant of quality of life amongst urban elderlies.” PhD Thesis submitted to CSM Medical University, Lucknow, UP, India.
- [30] Tripathi RK, & Tiwari SC (2013). Cognitive functioning of community dwelling urban older adults with reference to socio-demographic variables. S. C. Gupta Young Scientist Award. *Indian Journal of Clinical Psychology*, 40(2), 92-102.
- [31] Tripathi, R.K. (2012). Quality of Life: An Important Issue in Geriatric Research. *Journal of Gerontology and Geriatric Research*, 1(4).1000e114. <http://dx.doi.org/10.4172/2167-7182.1000e114>
- [32] Tripathi, R.K. and Tiwari, S.C. Cognitive dysfunction in normally aging urban older adults: a community based study. *Indian Journal of Psychological Medicine*, 33 (2), 177-181, 2011.
- [33] WHOQOL Group. (1998). The World Health Organization Quality of Life Assessment (WHOQOL): Development and general psychometric properties. *Social Science & Medicine*, 46(12), 1569-1585.
- [34] WHOQOL Group. (1999). The World Health Organization WHOQOL: Tests of the universality of quality of life in 15 different cultural groups worldwide. *Health Psychology*, 18(5), 495-505.