

Conditions of Boundedness for Compact operators on Banach spaces

Mohammed Ahamed Osman Tom^{1, 2}

¹Tabuk University, College of Science, Department of Mathematics, ²University of Bahri & College of applied Science and Industrial

Abstract: In this paper we study Compact operator between Banach spaces which are bounded . We establish two conditions that must be satisfies for boundedness of Compact operator.

Keywords: Closure of image ; Continuous at every points ; Compact operator ; Bounded

1. Introduction

A Banach space is a normed linear space that a complete metric space with respect to the metric derived from its norm . See [7] .This space has studied by several authors in different context such a in [3] , [1] and [6] . We give a brief discussion of conditions for boundedness of Compact operator. First Condition is the existence of the constant $M \geq 0$ as we done in Remark 3 that the operator $T: X \rightarrow Y$ between two normed operator bounded by letting T is compact which associate with the closures of the images of $B(0,2)$ and $B(0,n)$ producing $\|T\| = \sup_{\|x\|=1} \|Tx\| \leq \sup_{x \in B(0,2)} \|Tx\|$. Since $\sup_{x \in B(0,2)} \|Tx\| \in \mathbb{R}_{\geq 0}$ we use a constant M instead of $\sup_{x \in B(0,2)} \|Tx\|$ for some $M \in \mathbb{R}_{\geq 0}$, hence $\|Tx\| \leq M$ is bounded as long as $TB(0,2) \subset \overline{TB(0,2)}$ and $TB(0,2)$ is compact and hence bounded .In Lemma 4 we put $S = \{X \in x: \|x\| = 1\}$. Since T(S) is compact and $\|x\| = 1$ this result $\|T\| \leq M$ which is bounded by constant M .Second condition is the operator T must be continuous at every point . See Theorem 6 we show that the operator $T: X \rightarrow Y$ is bounded by letting T is continuous for all $x, y \in X$ we have $\|Tx - Ty\| = \|T(x - y)\| \leq M\|x - y\|$ where M is a constant, then by letting T is continuous at 0. Since T is linear we have $T(0) = 0$ there exists $\delta > 0$ with $\|x\| \leq \delta$ and $x \neq 0$ we define \tilde{x} by $\tilde{x} = \frac{\delta x}{\|x\|}$. Then $\|\tilde{x}\| \leq \delta$. From linearity of T that $\|Tx\| = \frac{\|x\|}{\delta} \|T\tilde{x}\| \leq M\|x\|$ where $M = \frac{1}{\delta}$. Thus T is bounded and continuous at zero , then at every point . (See Remark 3, Lemma 4, Definition 5 and Theorem 6) .

2. Compact operators on Banach space

A continuous linear operator $T: X \rightarrow Y$ on Banach spaces is compact when T maps bounded sets in X pre-compact sets in Y , that is , sets with compact closure . Since bounded sets lies in some ball in X , and since T is linear , its suffices to verify that T maps the unit ball in X to a pre-compact set in Y . Finite -rank operators are clearly compact . Both right and left compositions ToS and SoT of compact T with continuous S produce compact operators . Recall a criterion for pre-compactness : a set E in a complete metric space is pre-compact if and only if its totally bounded , sense that , given $\varepsilon > 0$, E is covered by finitely-many open balls of radius ε . We claim that operator-norm limits $T = \lim_i T_i$ of compact operators T_i are compact : given $\varepsilon > 0$, choose T_i

so that $\|T_i - T\|_{op} < \varepsilon$, cover the image of the Unit ball B_1 under T_i by finitely-many open ball U_k of radius ε .

Since $\|T_i x - Tx\| < \varepsilon$ for all $x \in X$, enlarging the balls U_k to radius 2ε covers TB_1 .

(If desired , rewrite the proof replacing ε by $\frac{\varepsilon}{2}$) . So there exist Banach spaces with compact operators which are not norm-limits of finite -rank operators . Recall that every compact operators $T: X \rightarrow Y$ on Hilbert space is an operator norm limit of finite rank operators . Given $\varepsilon > 0$, let y_1, \dots, y_n be the centers of open ε - balls in Y covering TB_1 , where B_1 is the unit ball in X . Let $T_\varepsilon = PoT$ where P is the projection of Y to the span of the y_i , then similarly , the sum of two compact operator is compact .

Definition 1

A compact operator is a linear operator T from Banach space X to another Banach space Y , such that the image under T of any bounded subset of X is necessarily a bounded operator , and so continuous . Any bounded operator Y that has finite rank is a compact operator , the class of compact operators is a natural generalization of the class of finite -rank operators in an infinite-dimensional setting . When Y is a Hilbert space , it is true that any compact is a limit of finite-rank operators can be defined alternatively as the closure in the operator norm of the finite -rank operators .

Lemma 2 : Let K be a compact operator on \mathcal{H} and suppose (T_n) is a bounded sequence in $\mathcal{B}(\mathcal{H})$ such that , for each $x \in \mathcal{H}$ the sequence $(T_n x)$ converges to Tx , where $T \in \mathcal{B}(\mathcal{H})$. Then $(T_n K)$ converges to TK in norm. Briefly, the above can be rephrased as:

If $K \in \mathcal{K}(\mathcal{H})$ and $\|T_n x - Tx\| \rightarrow 0$ for all $x \in \mathcal{H}$ then $\|T_n K - TK\| \rightarrow 0$.

In words: Multiplying by a compact operator on the right converts a pointwise convergent sequence of operators into a norm convergent one .

Proof

Since (T_n) is bounded sequence , $\|T_n\| \leq M$ for some constant M . Then for all $x \in \mathcal{H}$, $\|Tx\| = \lim_n \|T_n x\| \leq M\|x\|$ and so $\|T\| \leq M$.

Let K be compact and suppose that $\|TK - T_n K\| \rightarrow 0$. Then there exists some $\delta > 0$.

And a subsequence $(T_{n_i}K)$ such that $\|TK - T_{n_i}K\| > \delta$. Choose unit vectors (x_{n_i}) of \mathcal{H} such that $\|(TK - T_{n_i}K)x_{n_i}\| > \delta$. [That this can be done follows directly from the definition of the norm of an operator .]. Using the fact that K is compact , we can find a subsequence (x_{n_j}) of (x_{n_i}) such that (Kx_{n_j}) is convergent . Let the limit of this sequence be y . Then for all j

$$\delta < \|(TK - T_{n_j}K)x_{n_j}\| \leq \|(T - T_{n_j})(Kx_{n_j} - y)\| + \|(T - T_{n_j})y\|$$

Now , using the convergence of (Kx_{n_j}) to y , there exists n so that , for $n_j > n$,

$$\|Kx_{n_j} - y\| < \frac{\delta}{8M}$$

Also , using the convergence of (T_{n_j}) to T , there exists m so that , for $n_j > m$, $\|(T - T_{n_j})y\| < \frac{\delta}{4}$. Then , for $j > \max[n, m]$ that right hand side of the displayed inequality is less than $\frac{\delta}{2}$, and this contradiction shows that the supposition that $\|TK - T_nK\| \rightarrow 0$ is false .

3. First Condition of Boundedness of Compact operator

Remark 3: A compact operator is bounded .

Proof

First we show that a compact linear operator $T : X \rightarrow Y$ between normed spaces is bounded , if T is compact , then the closure of the image of $B(0,1)$ and $B(0, n)$ is compact , then

$$\|T\| = \sup_{\|x\|=1} \|Tx\| \leq \sup_{x \in B(0,2)} \|Tx\| \leq M$$

For some $M \in \mathbb{R}_{\geq 0}$ since $TB(0,2) \subset \overline{TB(0,2)}$ and $\overline{TB(0,2)}$ is compact hence bounded .

Lemma 4 : Every compact operator is bounded .

Proof

Put $S = \{x \in X : \|x\| = 1\}$, then $T(S)$ is relatively compact , hence bounded (by M) therefore , $\|T\| \leq M$.

Definition 5

Let X and Y be two normed linear spaces . We denote that both X and Y norms by $\|\cdot\|$. A linear operator $T : X \rightarrow Y$ is bounded if there is constant $M \geq 0$ such that

$$\|Tx\| \leq M\|x\| \text{ for all } x \in X \quad (1)$$

If no such constant exists , the we say that T is unbounded .

If $T : X \rightarrow Y$ is a bounded linear compact operator we define operator norm $\|T\|$ of T by

$$\|T\| = \sup\{M \mid \|Tx\| \leq M\|x\| \text{ for all } x \in X\} \quad (2)$$

We denote the set of all linear maps $T : X \rightarrow Y$ by $\mathcal{L}(X, Y)$, and the set of all bounded linear maps by $\mathcal{B}(X, Y)$. Then the domain and range spaces are the same , we write

$\mathcal{L}(X, X) = \mathcal{L}(X)$ and $\mathcal{B}(X, X) = \mathcal{B}(X)$ equivalent expressions for $\|Tx\|$ are :

$$\|T\| = \sup_{x \neq 0} \frac{\|Tx\|}{\|x\|} ; \|T\| = \sup_{\|x\| < 1} \|Tx\| ;$$

$$\|T\| = \sup_{\|x\|=1} \|Tx\| \quad (3)$$

4. Second Condition of Boundedness of Compact operator

Theorem 6

Linear compact operator is bounded if it is continuous .

Proof

First , suppose that $T : X \rightarrow Y$ is bounded .

Then , for all $x, y \in X$, we have

$$\|Tx - Ty\| = \|T(x - y)\| \leq M\|x - y\|$$

Where M is a constant for which (1) holds, therefore we can take $\epsilon = \epsilon/M$.

In the definition of continuity, T is continuous.

Second , suppose that T is continuous at 0 . Since T is linear, we have $T(0) = 0$.

Choosing $\epsilon = 1$ in the definition of continuity , we conclude that there is a $\delta > 0$.

Such that $\|Tx\| \leq 1$ whenever $\|x\| \leq \delta$ for any $x \in X$, with $x \neq 0$, we define

$$\tilde{x} \text{ by } \tilde{x} = \delta \frac{x}{\|x\|}$$

Then $\|\tilde{x}\| \leq \delta$, so $\|T\tilde{x}\| \leq 1$.

It follow from linearity of T that $\|Tx\| = \frac{\|x\|}{\delta} \|T\tilde{x}\| \leq M\|x\|$.

Where $M = \frac{1}{\delta}$. Thus T is bounded . The proof shows that if any compact linear map is continuous at zero , then its continuous at every point .

References

- [1] Bounded operators .Jan Dereziński ,Department of Mathematical Methods in Physics, Warsaw University , Hoza74 , 00-682,Warszawa , Poland ,Lecture Notes ,version of Jan ,2007, January 31,2007 .
- [2] Compact operator and Classical Function Theory , Joel . Shapiro , Portland State University CMFT 2013 .
- [3] Enflo ,A counterexample to the approximation problem in Banach spaces , Acta.130 (1973) , 309-317.
- [4] Factorization of weakly Compact operators between Banach spaces and Frechet or (LB)- spaces New Mexico State University , U.S.A , Received :July 2009 , Accepted : August 2009 .
- [5] Kerss , R.,Linear Integral Equations . Applied Mathematical Science 82 . New York : Springer 1999 .
- [6] T. Kato , Perturbation theory for Linear operators ,Springer ,1966 ,reprinted 1955.
- [7] Michael Reed and Barry Simon , Methods of Modern Mathematics Physics I Functional Analysis , Academic press,1980.