

Pulmonary Radiographic Findings in Leptospirosis Patients

Dr. Ajinkya Maheshwaril, Dr. Purvi Desai², Dr. Bhagwati Ukani³

^{2,3}Associate Professor

Abstract: We studied chest radiographs of 45 patients who presented during June 2015 to October 2016 at New civil hospital, Surat who had a high serum immunofluorescent assay titer for leptospirosis. Thirty five of 45 (77%) patients had abnormal chest films. Nearly all of these patients had respiratory symptoms and needed oxygen therapy. 40 of the 45 (65%) had respiratory symptoms, but only 35/40 (87%) had abnormal radiographs. 23 of the 35 (65%) and 20/35 (57%) who had abnormal chest radiographs had coexisting impaired renal function or jaundice, respectively. There were 6/35 (17%) patients who had abnormal chest radiographs without renal or liver impairment. **Methodology:** Patients with high serum immunofluorescent titre for leptospirosis underwent chest xray with SIEMENS HELIPHOS-D 500 mA xray machine. Abnormality on chest xray were compared with disease severity and organ involvement. **Result:** The most common finding on the abnormal chest radiograph was bilateral diffuse air space disease, which resolved within 7 days. No permanent lung damage was seen. The patients who had an abnormal chest radiograph needed longer hospitalization, than those without an abnormal chest radiograph (average 12 days and 5 days, respectively).

Keywords: Leptospirosis, Chest Xray, Diffuse airway disease, Multi organ failure

1. Introduction

Leptospirosis is a systemic infectious disease caused by spirochetes of the genus *Leptospira*. Human infection typically results from exposure to infected animal urine, by either direct contact or indirect exposure through water or soil. Clinical features vary, and multiorgan involvement (mainly involving skeletal muscles, kidneys, adrenal glands, liver, stomach, spleen and lung) is common. Respiratory manifestations have been reported in 20-70% of patients, but such features of the disease are often overshadowed by other more serious expressions. Chest radiographic abnormalities have been observed in 11-64% of these patients. Even though it has a world-wide distribution, it is more common in tropical countries. Reports of chest radiographic findings in leptospirosis patients in Asian countries are few. We review a large number of leptospirosis patients in Surat, India who were infected and had chest radiographs between June 2016 to October 2016.

2. Methodology

The names and hospital numbers of leptospirosis patients admitted to our hospital from June 2016 to October 2016, were collected from the serology unit. There were serum samples in 100 patients that had an immunofluorescent antibody (IFA) titer higher than 400. 45 patients were admitted to our hospital. The medical records of these patients were studied, noting four particular features:

- 1) Respiratory system involvement, especially symptoms (dyspnea, cough and hemoptysis) and oxygen therapy, if any.
- 2) The renal function, especially the serum creatinine level and need for dialysis.
- 3) The hepatic function, especially the serum bilirubin and transaminase enzyme levels.
- 4) The final patient outcome.

Chest radiographic abnormalities were analyzed with regard to any patterns of lung infiltration.

The distribution of such infiltration and the extrapulmonary findings (adenopathy and pleural effusion) were also evaluated. The progression and resolution of the abnormalities were also studied.

3. Discussion

The definitive diagnosis of leptospirosis depends mainly on serologic tests. The microscopic agglutination (MA) method is the reference standard serodiagnostic test. However, it is time-consuming and increases the risk of laboratory-acquired infection which decreases the specificity. The immunofluorescent antibody (IFA) test is more practical for initial diagnosis and initial management of leptospirosis.

The most sensitive and specific IFA test is a fourfold or greater increase in titers of acute and subsequent subsequent sera. However, this is often not possible in our hospital, where most patients are lost to follow-up once they are discharged. In addition, since it is the only tertiary unit in southern Thailand, our serology unit receives specimens, most of which are single, from many other hospitals.

We chose a cut-off titer for single sera of 1:400 based on a study done at our hospital showing that such a titer was the most specific and moderately sensitive.

Abnormal chest radiographs related to leptospirosis were found in 77% of our patients. Similar to other studies, the characteristic and most common chest radiographic abnormality in our patients was multiple, tiny, ill-defined nodules in both lungs, some of which subsequently became confluent and turned into patchy infiltration.


These nodules might have been acinar or alveolar nodules representing fluid in the terminal airspaces. They typically had rapid progression and resolution. No pathological study was done in our patients, but these nodules might have been hemorrhagic spots and the bilateral patchy infiltration could

have been pulmonary hemorrhage as mentioned in some reports.

The ultrastructural findings were uniform with consistent capillary lesions in hemorrhagic areas.

They suggested leptospirosis caused hemorrhagic pneumopathy and septal capillary lesions.

However, this chest radiographic finding is not pathognomonic for leptospirosis since it can be seen in other diseases, such as aspiration, bronchioloalveolar cell carcinoma, or pulmonary hemorrhage due to many entities. Correct determination of these diseases needs clinical correlation. The second most common chest radiographic abnormality was bilateral patchy infiltration, which may have simply been a late manifestation of the first group. These cases were unlikely to have been adult respiratory distress syndrome as mentioned in one report since they had rapid resolution. In adult respiratory distress syndrome, complete resolution of chest radiographic abnormalities often needs a much longer time. They were also unlikely to have been pulmonary edema due to their peripheral distribution and lack of other supportive evidence, such as Kerley's lines or pleural effusion. We also observed that most patients who had abnormal chest radiographs had higher morbidity. They had longer hospitalization and usually had renal and liver involvement. A small number of patients with renal or liver involvement had abnormal chest radiographs. This may indicate that pulmonary involvement could be either a late manifestation of the disease or correlate with disease severity. We believe that the patients in the third group who had chest radiographic abnormalities consistent with pulmonary edema were likely to have had impaired renal excretion and received excessive fluid replacement rather than having real pulmonary involvement from leptospirosis. Other nonspecific abnormal findings including subsegmental atelectasis (group 4) and pleural effusions (group 5), which occurred in a small number of patients. We could not make any firm conclusions concerning bilateral reticular infiltrations in one patient (group 6), since he was also diagnosed as being HIV-positive, and he improved after receiving treatment for *Pneumocystis carinii* pneumonia. Even though leptospirosis is prevalent in tropical regions, reports from Asia are few, and all large studies to date of radiographic findings from leptospirosis are from South America. Our current study confirms what is currently known. From 1953 to now, in various regions of the world, patterns of lung involvement in leptospirosis do not seem to have changed; most commonly pulmonary hemorrhage and much less common ARDS. Pulmonary hemorrhage is usually mild in the course of disease, with spontaneous resolution and no permanent damage to the lungs. However, pulmonary hemorrhage and ARDS are two of the most fatal conditions in leptospirosis.



The most common abnormal chest radiographs found in our leptospirosis patients (A) showed multiple ill-defined nodules in both lungs (A, with a magnified view of a left upper lobe, B) that subsequently became confluent and turned to bilateral patchy infiltration (C, with a magnified view of the left upper lobe, D)

4. Summary

There were 31 men and 14 women, with ages ranging from 9 to 71 years (mean = 45 years). Forty five patients were hospitalized, including 15 to the intensive care unit. The length of hospitalization ranged from 4 to 39 days (mean = 9 days). Four died, 2 from adult respiratory distress syndrome, 1 from acute renal failure, and 1 due to a myocardial infarction with pulmonary edema. Respiratory symptoms developed in 40/45 patients (89%); 28 of the 40 (72%) had cough, 18/40 (45%) had dyspnea and 4/40 (10%) had hemoptysis. A rising serum creatinine higher than 1.5 g% occurred in 23 patients; 20 patients had jaundice (serum bilirubin more than 2 g %).

Thus, in the end, there were 35/45 (77%) patients who had chest radiograph abnormalities which were considered leptospirosis related. These abnormalities could be categorized into 6 groups:

- 1) Fourteen patients (41%) presented with multiple tiny ill-defined nodules in both lungs, which, in 10 of the 14, subsequently became bilateral diffuse patchy infiltrates. One patient had a right pleural effusion.
- 2) Eight patients (24%) had bilateral diffuse patchy infiltration.
- 3) Five showed bilateral haziness of the hili with congested pulmonary vessels and Kerley's lines which were consistent with pulmonary edema.
- 4) Small pleural effusions were present in four patients.
- 5) Small but thick plates in the basal lungs, so-called plate atelectasis, were the main feature in three (8%) patients.
- 6) There was one patient who had perihilar reticular infiltration in both lungs. Because this patient also had HIV, he therefore received treatment for both leptospirosis and *Pneumocystis carinii* pneumonia.

5. Conclusion

All 35 of the patients who had abnormal chest radiographs related to leptospirosis were hospitalized, 15 (42%) of them in the intensive care unit. The average length of hospitalization was 12 days, versus 5 days for those who had a normal chest radiograph. Of the patients with abnormal chest radiographs, 30/35 (85%) needed oxygen therapy (including 11 with endotracheal intubation and artificial ventilation support), 23/35 (65%) had a rising serum creatinine, including 11 patients who needed dialysis, and 20/35 (57%) who had a rising serum bilirubin and liver enzymes.

Six of 35 (17%) patients had an abnormal chest radiograph without renal or liver impairment. All five hemoptysis patients had abnormal chest radiographs.

There were 22 patients who had follow-up chest radiographs. Complete resolution occurred within 3 days, 3-7 days and more than 7 days in 6/22 (29%), 13/22 (59%) and 3/22 (12%) patients, respectively. No permanent damage was seen.

References

- [1] Antonio C, Nicodemo, Maria Irma S. Lung lesions in human leptospirosis: microscopic immunohistochemical and ultrastructural features related to thrombocytopenia. *Am J Trop Med Hyg* 1997; 56: 181-7.
- [2] Felson B. The pulmonary airways. In: Felson B, ed. *Chest roentgenology*. Philadelphia: WB Saunders, 1973: 265-313.
- [3] Im JG, Yeon KM, Han MC, et al. Leptospirosis of the lung: radiographic findings in 58 patients. *AJR Am J Neuroradiol* 1989; 152: 955-9.
- [4] Lee REJ, Terry SI, Walker TM, Uraquhart AE. The chest radiograph in leptospirosis in Jamaica. *Br J Radiol* 1981; 54: 939-43.
- [5] Lomar AV, Diament D, Torres JR. Leptospirosis in Latin America. *Infect Dis Clin North Am* 2000; 14: 23-39.
- [6] McCrumb FR Jr, Stockard JL, Robinson CR, et al. Leptospirosis in Malaya. I. Sporadic cases among military and civilian personnel. *Am J Trop Med Hyg* 1957; 6: 238-56.
- [7] Niwattayakul K, Homvijitkul J, Niwattayakul S, et al. Hypotension, renal failure and pulmonary complication in leptospirosis. *Ren Fail* 2002; 24: 297-305.
- [8] Du Couedic L, Courtin JP, Poubeau P, et al. Patent and occult intra-alveolar hemorrhage in leptospirosis. *Rev Mal Respir* 1998; 15: 61-7.
- [9] Dupont H, Dupont-Perdrizet D, Perie JL, et al. Leptospirosis: prognostic factors associated with mortality. *Clin Infect Dis* 1977; 25: 720-4.
- [10] Im JG, Yeon KM, Han MC, et al. Leptospirosis of the lung: radiographic findings in 58 patients. *AJR Am J Neuroradiol* 1989; 152: 955-9.
- [11] Kiatboonsri S, Vathesatogit P, Charoenpan P. Adult respiratory distress syndrome in Thai medical patients. *Southeast Asian J Trop Med Public Health* 1995; 26: 774-80.
- [12] Lee REJ, Terry SI, Walker TM, Uraquhart AE. The chest radiograph in leptospirosis in Jamaica. *Br J Radiol* 1981; 54: 939-43.
- [13] Lomar AV, Diament D, Torres JR. Leptospirosis in Latin America. *Infect Dis Clin North Am* 2000; 14: 23-39.
- [14] McCrumb FR Jr, Stockard JL, Robinson CR, et al. Leptospirosis in Malaya. I. Sporadic cases among military and civilian personnel. *Am J Trop Med Hyg* 1957; 6: 238-56.