

Sonographic Diagnosis of Amniotic Band Syndrome and its Risk Factors

Dr Vidhi Sahni¹, Dr Deepak Sharma²

¹Senior Resident, Kalpana Chawla Medical College

²Consultant at Neocare, Jaipur

Abstract: *Objective:* The aim of this study was to describe our experience with amniotic band syndrome (ABS), define specific sonographic characteristics and common features. *Methods:* Patients diagnosed with ABS underwent detailed ultrasound evaluation at the time of diagnosis and during follow-up. Their ultrasound examinations and medical records concerning the current pregnancy and past medical records were analyzed. *Results:* Ten pregnancies were diagnosed with ABS. Most pregnancies were diagnosed at the beginning of the second trimester. Two cases were bichorionic twin pregnancies involving one of the fetuses and these were the only women who continued their pregnancies to term. The other eight cases with ABS chose to terminate their pregnancies. One pregnancy was conceived following trachelectomy. We found a significantly higher rate of prior uterine surgeries (p=0.008) in patient with ABS compared to control. *Conclusions:* ABS diagnosed in early pregnancy can be a sporadic event. However, there is a higher risk of ABS in pregnancies preceded by uterine procedures.

Keywords: amniotic band syndrome, ultrasound, previous uterine surgery

1. Introduction

Amniotic band syndrome (ABS) is a rare congenital disorder caused by entrapment of fetal parts by fibrous amniotic bands, resulting in devastating cosmetic and functional disability¹. Incidence varies from one in 1200 to one in 15000 live births². The hallmark of ABS is the presence of fibrous bands; however, the precise mechanism of amniotic band formation is still unknown³. Several risk factors for ABS have been suggested, such as smoking, drug use, maternal hyperglycemia, amniocentesis and high altitude^{4,5}. The aim of this study was to describe our experience with ABS diagnosis in a tertiary center ultrasound unit, define specific sonographic characteristics, risk factors and common features of this syndrome.

2. Methods

This retrospective study was conducted at a tertiary care centre over a period of 14 years. All patients with diagnosis of amniotic band syndrome were included in the study. Patient's past medical and obstetrical history was collected. Data details of ultrasound at the time of diagnosis and follow up was retrieved. All deliveries in the past eight years were used as control (86686 deliveries). Statistical analysis was performed by a two-tailed chi-square analysis with Yate's correction. The study was approved by the local Institutional Review Board committee.

3. Results

Out of all patients included in the study, 10 patients were diagnosed with amniotic band syndrome. Eight patients were diagnosed with ABS during early second trimester (13–15 weeks) at the first ultrasound scan. Two other cases were diagnosed during late second trimester and early third trimester (22 and 27 weeks). Eight patients discontinued the pregnancy, two of the 10 continued the pregnancy to term. Both of which were twin pregnancies

involving one of the fetuses. Most of the anomalies observed in this study were limb deformities. Other anomalies like cleft lip and cleft palate were also seen in some cases. We compared our study group to all 86686 deliveries in the previous eight years. In the study group, five out of the 10 patients had prior uterine surgery.

Table 1: Characterization of all cases included in this study

	Gestational week	Obstetrical history	Fetal findings	outcome	Previous uterine surgery
1.	14	G3P2	Amputation of hand	TOP	
2.	13	G1P0	Hands confined by amniotic band	TOP	
3.	14	G3P2	Foot and hand confined by amniotic band	TOP	CS
4.	15	G2P0	Deformation of hand	TOP	
5.	27	G6P5	Amputation of hand	NVD at 37 weeks.	
6.	14	G1P0	Absence of fingers 2-4 in both hands	TOP	
7.	15	G2P0A1	Cleft lip and palate, foot and hand confined by band	TOP	Laparoscopic myomectomy
8.	13	G2P1	Fetus confined by amniotic bands	TOP	Radical trachelectomy
9.	22	G3P2	Cleft lip and palate, amputated hands	TOP	Previous CS
10.	15	G4P2A1	Absence of distal phalanx in fingers 2-3 in left hand.	CS at 38 weeks	CS

Three patients had prior cesarean sections (CS), one patient had a laparoscopic myomectomy and another pregnancy was conceived following radical trachelectomy. The rate of previous uterine surgery in women with ABS was significantly higher than in the control group (50 versus 14.9%, $p=0.008$). In three cases, a small vestige in the distal part of the amputated limb was reported (Figure 1) and it was visible after delivery (Figure 2).




Figure 1: 3D ultrasound (27 wk) pointing small vestige at site of amputation.



Figure 2: Post delivery picture of vestige.

4. Discussion

Amniotic band syndrome can cause malformations that range from mild deformities to severe anomalies that are incompatible with life⁶. The most common defects associated with ABS are limb defects such as focal constrictions, amputations, pseudosyndactyly, and brachydactyly. An amputation with the distal bone protruding beyond the soft tissue at the site of the amputation is diagnostic of ABS.⁷ The two main theories regarding the pathogenesis of ABS are known as the extrinsic and intrinsic models. The most accepted etiology for ABS is the one proposed by Torpin. He proposed a sequence mechanism originating in rupture of the amniotic sac, followed by extrusion of fetal part into the chorionic cavity and resulting in fetal anomalies due to vascular compression by the amniotic bands.⁸ Other authors suggested that ABS is formed by vascular disruption⁹ and conditions associated with relative hypoxia, such as smoking and living in high altitude, have been identified as risk factors for ABS.⁴

Orioli *et al.* studied an increased incidence of ABS in the population living at a high altitude, in primipara, in women with a history of febrile illness in the antenatal period, and in women with a history of vaginal bleeding in the first trimester.⁶ The pathogenesis of ABS is thought to be disruption of amnion allowing the embryo to enter the chorionic cavity resulting in entanglement of fetal parts creating various malformations¹⁰. When any of the malformation involving craniofacial region is seen ABS should be suspected while doing the ultrasound. A search for amniotic band (although not diagnostic) should also be made in ultrasound.

In our study we found that history of prior uterine surgeries (50%) is associated with high risk of ABS in the subsequent pregnancy. Previous uterine surgery leading to uterine vascular compromise (placental hypoxia) may be the cause. This study also emphasizes the importance of ultrasound examination in first trimester specially in women with history of uterine surgery.

5. Conclusion

We found that there is a higher risk of ABS in previous uterine surgeries and also it can be diagnosed in ultrasound by characteristic fetal malformations involving craniofacial region, trunk, extremities alone or in combination.

References

- [1] Rushton DI. Amniotic band syndrome. *Br Med J (Clin Res Ed)* 1983;286:919–20.
- [2] Seeds JW, Cefalo RC, Herbert WN. Amniotic band syndrome. *Am J Obstet Gynecol* 1982;144:243–8.
- [3] Hunter AG, Seaver LH, Stevenson RE. Limb-body wall defect. Is there a defensible hypothesis and can it explain all the associated anomalies? *Am J Med Genet A* 2011;155A:2045–59.
- [4] Cignini P, Giorlandino C, Padula F, et al. Epidemiology and risk factors of amniotic band syndrome, or ADAM sequence. *J Prenat Med* 2012;6:59–63.

- [5] Luehr B, Lipsett J, Quinlivan JA. Limb-body wall complex: a case series. *J MaternFetal Neonatal Med* 2002;12:132-7.
- [6] Orioli IM, Ribeiro MG, Castilla EE. Clinical and epidemiological studies of amniotic deformity, adhesion, and mutilation (ADAM) sequence in a South American (ECLAMC) population. *Am J Med Genet A* 2003;118A: 135-45.
- [7] Chandran S, Lim MK, Yu VY. Fetal acalvaria with amniotic band syndrome. *Arch Dis Child Fetal Neonatal Ed* 2000;82:F11-3.
- [8] Torpin R. Amniochorionic mesoblastic fibrous strings and amniotic bands: Associated constricting fetal malformations or fetal death. *Am J Obstet Gynaecol* 1965;91:65-75.
- [9] Russo R, D'Armiento M, Angrisani P, Vecchione R. Limb body wall complex: a critical review and a nosological proposal. *Am J Med Genet* 1993;47:893-900.
- [10] Higginbottom MC, Jones KL, Hall BD, Smith DW. The amniotic band disruption complex: timing of amniotic rupture and variable spectra of consequent defects. *J Pediatr* 1979;95:544-549.

