

Isolation and Molecular Characterization of a Variant Infectious Bronchitis Virus in Kafrelsheikh, Egypt During 2016

Asmaa Magouz¹, Gabr Elbagoury², Shimaa Ayoub³

^{1,3}Department of Virology, Faculty of Veterinary Medicine, Kafrelsheikh University, 33516, Egypt

²Department of Virology, Faculty of Veterinary Medicine, Banha University

Abstract: *Infectious bronchitis (IB) is a major disease affecting chickens all over the world. The present study was designed for isolation and molecular characterization of IB viruses circulating among different regions in Kafrelsheikh province, Egypt. Twenty two field samples were collected from broiler chicken flocks. The collected samples were propagated in the allantoic cavity of embryonated chicken eggs. The isolated strains were characterized by polymerase chain reaction (PCR) through amplification of S1 gene followed by sequence analysis of one isolate. Egg inoculation revealed dwarfing with characteristic curling appearance. PCR assay shows 10 positive field samples and phylogenetic analysis revealed the closest relationship with the Israel variant strain IS/1494/06.*

Keywords: IBV, Egypt, Kafrelsheikh, S1 gene.

1. Introduction

Acute upper-respiratory diseases in chickens are major causes of economic losses throughout the world due to high mortality rates especially in poorly managed cases (Yashpal *et al.*, 2004). Infectious bronchitis (IB) is an acute highly contagious, worldwide distributed viral disease of chickens caused by infectious bronchitis virus (IBV) (Jahantigh *et al.*, 2013). IBV affects chickens of all ages, causing respiratory and renal symptoms (Cook *et al.*, 2012). The disease commonly causes respiratory manifestations such as coughing, sneezing, rales, and nasal discharge, (El-Mahdy *et al.*, 2011). In addition, some nephropathogenic strains of IBV lead to severe kidney lesions (Pohuang *et al.*, 2009).

In Egypt, IB infection was first recognized during 1950s by Ahmed, 1954 from birds with respiratory distress with the isolation of a variant resembling the Dutch variant D3128 (De Wit *et al.*, 2011). Later, isolates closely related to D274, Massachusetts, D-08880, 4/91, Egypt/Beni-Suef/ 01 and the Israeli variant were isolated from different Egyptian provinces (Abdel-Moneim *et al.*, 2006). The IB viral particles are corona shaped about 120 nm in diameter with club-shaped surface spikes. (El-Shafey 2008).

Viral genome is a positive-sense RNA of about 27.6 kb in overall size. IBV encodes four structural proteins, the spike (S) glycoprotein, the membrane (M) glycoprotein, the nucleocapsid (N) protein and the envelope (E) protein (Mo *et al.*, 2013). Many IBV serotypes have been isolated worldwide, e.g. Connecticut, Massachusetts (Mass), 4/91, D274, H120, Baudette and California with low or even no-cross protection between each other (Roussan *et al.*, 2008). The spike (S) glycoprotein gene consists of two subunits, S1 and S2. S1 is frequently used for PCR and sequencing due to its high variability which can reach up to 50 % in amino acid sequence between IBV serotypes (Patel *et al.*, 2015). Thus, accurate diagnosis of the exact serotype identity circulating in chicken farms in Egypt is very essential for

modifying the effective vaccination program to get over IB outbreaks in Egypt (Mahgoub *et al.*, 2010). Controlling IB disease is a great challenge because of the huge variations in the serotypes which emerge from time to time. (El-Mahdy *et al.*, 2010).

In this study we report the isolation and molecular characterization of a nephropathogenic IB virus isolated from chicken flocks in Kafrelsheikh province during the period from January 2015 till March 2016.

2. Material and Methods

2.1 Source and collection of samples

A total of 22 tissue samples (lung, trachea and kidneys) were collected from chicken flocks in Kafrelsheikh province experiencing respiratory manifestations and high mortality rates during the period from January 2015 till March 2016. The samples were stored at -80°C till processed.

2.2 Virus isolation

Samples from the same flocks were pooled and grinded to prepare 10% tissue suspensions in PBS (pH 7.2) solution containing Penicillin (1 million IU/L) and Gentamycin (250 mg/L). It was centrifuged at 10,000 for 15 min at 4°C and the supernatant fluid was inoculated into three 9-11- days old SPF embryonated chicken eggs via the allantoic cavity route. The eggs were incubated at 37°C for 4-7 days and were daily examined.

2.3 Identification of the isolated viruses by PCR

• RNA extraction

RNA was extracted from infected allantoic fluid using Trizol Easy-Red™ Total RNA extraction Kit (Intron Biotechnology) as per manufacturer's protocol. Briefly, 350 µl of allantoic fluid were mixed with 750 µl of Trizol

reagent then centrifuged at 13,000 rpm /5 min. The supernatant was transferred to a fresh DEPC treated tube and added with 200 µl of cold chloroform. Then centrifuged at 13,000 rpm /10 min at 4°C. 500 µl of chilled isopropanol was added to the upper aqueous phase then centrifuged at 13,000 rpm /5 min at 4°C. The supernatant was discarded and the pellet was washed with 70% cold ethanol. The pellet was re-suspended in 30 µl RNase free water and stored at -80°C till used

• **Reverse transcription of extracted RNA to cDNA**

The reverse transcription (RT) reaction was performed using Maxime™ RTPreMix Kit (Gene On) with specific primers as per manufacturer's protocol. Briefly, 5 µl of RNA, 1 µl of

specific primer and 2 µl of RNAs free water were added in a DEPC treated tube and incubated at 70 °C for 5 min. Then kept on ice for 5 min. Another mixture of 4 µl 5X reaction buffer, 1 µl dNTP mix ,1 µl RNAs inhibitor, 1 µl MMLV Reversease (200u/ µl) and 13 µl RNase free water. Mix I and II were combined together, vortexed and incubated at 50°C for 60 min, then at 70°C for 5 min. cDNA was stored at -20°C till used.

• **PCR reaction**

Primers that specifically amplify IBV (S1 gene) were commercially synthesized (Metabion international AG, Germany) and are shown in table 1.

Table 1: Nucleotide sequences of S1 gene primer used in this study.

Primer	Oligonucleotide sequence	gene	bp	Reference
Co1	TGACTCTTTTGKTGCACTAT	S1 gene	403bp	Farsang et.al 2002
Co2	AAATTATAATAACCACTCTGA			

The PCR reaction mixture consisted of 12.5 µl of (Taq DNA Polymerase 2X-preMix™, GeneON) 1 µl of forward and reverse primers, 3 µl of cDNA and 7.5 µl of PCR grade water. The thermal conditions consisted of initial denaturation at 95°C/5 min then 35 cycles of denaturation at 95°C/45 sec, annealing at 55°C/1min and extension at 72°C/1min then final extension at 72°C/10 min. A sample with only primers and PCR grade water was used as a negative control.

• **Detection of PCR product by gel electrophoresis**

Amplified PCR products were visualized under ultraviolet light after agarose gel electrophoresis using 1.5% agarose and staining with 0.1 µg/ml ethidium bromide.

2.4 Sequencing and phylogenetic analysis

The nucleotide sequence data of our isolate IBV/KFS/2016 was submitted to the GenBank sequence database and have been assigned the accession number KY072832.

Resulting sequences were analyzed using the nucleotide Basic Local Alignment Sequence Tools (BLASTn) and the multiple sequence alignment programs (ClustalW) using reference GenBank sequences. Phylogenetic tree was constructed via multiple alignments of 403 bp nucleotide sequence of our isolate (IBV/KFS/2016) using the neighbor-joining (N-J) tree method using the MEGA version 6 software.

3. Results

3.1 Virus isolation

The inoculated ECE were daily examined and embryo mortality was recorded. Embryonic deaths occurred between 3-5 days post inoculation with characteristic stunting and curling of embryos after applying three blind egg passages as shown in Fig.1.



Figure 1: Stunting and curling of embryos infected with IBV (right) compared to a normal non inoculated embryo (left).

3.2 Identification of the isolated viruses by PCR

The collected allantoic fluids were examined for the presence of IBV virus through amplification of the S1 gene using specific primers. Results revealed that 10 samples were positive for IBV as they show amplification of specific band (403bp) as shown in Fig 2.




Figure 2: Positive amplification of specific bands of S1 gene of 10 field samples (403bp).

3.3 Sequencing and phylogenetic analysis

S1 sequence analysis of our isolate IBV/KFS/2016 (KY072832) revealed its close relatedness to IBV-EG/13280F-SP1-2013(KU979008.1), Eg/CLEVB-2/IBV/012 (JX173488.1) serotypes and the Israeli strains IS/1494/06 (EU780077.2), IB VAR2-06 (JX027070.1) and Israel/720/99 (AY091552.2) as shown in table 2 and Fig. 3.

Table 2. Nucleotide and amino acid identities of the isolate IBV/KFS/2061 (KY072832) compared with other Genbank located reference strains.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1--IBV/KFS/2016	89	91	92	90	90	91	90	91	90	90	89	91	89	89	89	89	89	89	89	89	89	89	89
2-D1887/2/11/EG	89	99	97	97	99	98	98	98	99	96	96	96	96	83	82	82	81	81	81	82	81	85	81
3-IBV/EG/13280F-SP1-2013	90	92	92	97	97	99	98	98	99	96	96	96	96	83	81	81	81	81	81	82	81	85	81
4-Israel/7/20/99	89	95	97	96	99	97	96	96	97	97	98	98	97	83	82	82	81	82	81	82	82	86	82
5-isolate IS/885	88	96	96	97	97	95	96	96	97	98	98	97	83	82	82	82	82	82	82	82	82	87	82
6-IBV/EG/1212S-SP1-2012	89	99	99	96	96	96	96	99	98	99	96	96	96	82	81	81	80	80	80	82	81	85	81
7-02572/2/14_EG	90	98	99	96	95	98	96	97	97	95	95	94	82	82	81	81	81	81	81	82	86	82	
8-Eg/1494/06	90	99	100	97	96	99	95	97	99	95	95	85	82	82	81	81	81	81	82	85	85	99	
9-IBV/EG/1586CV-SP1-2015	90	97	98	97	96	97	97	98	98	98	98	97	81	80	80	80	80	80	80	80	85	80	
10-IBV/EG/1442F-SP1-2014	88	98	98	95	95	98	97	98	96	96	95	95	82	80	80	80	80	80	80	80	85	80	
11-Eg/CLEVB-2/IBV/012	89	95	95	96	95	95	95	97	97	95	95	95	81	81	80	82	82	81	86	81	86	91	
12-IS/1494/06	89	95	95	96	95	95	95	97	94	100	99	81	81	81	80	82	83	81	86	81	86	91	
13-IB VAR2-06	87	93	94	95	94	93	93	94	95	92	98	98	81	81	80	82	82	81	85	81	85	96	
14-Egypt/F/03	79	82	82	81	82	81	83	83	80	82	82	81	81	80	82	82	81	81	86	81	82	15	
15-strain H120	79	82	82	81	82	81	83	83	80	82	82	81	81	100	98	95	80	99	81	99	82	15	
16-MA5 strain	79	82	82	81	82	81	83	83	80	82	82	81	81	99	98	95	80	99	81	99	82	16	
17-strain MA1-UPM2013	77	81	81	80	80	82	82	80	80	81	81	80	95	96	96	96	81	97	80	97	81	17	
18-Connecticut	80	81	81	82	83	80	82	82	81	80	83	83	82	92	93	93	92	82	95	80	94	81	
19-strain 4/91	86	84	85	85	86	84	86	85	84	87	87	86	80	80	79	81	82	80	84	80	81	19	
20-D4 strain	78	83	83	82	83	82	84	84	81	83	83	82	98	99	99	96	92	79	80	99	87	20	
21-isolate IB VAR2	83	85	85	87	88	85	87	86	85	83	86	86	85	85	85	83	84	85	89	81	86	21	
22-Eg10574/2010	78	80	80	80	80	80	81	81	79	79	80	80	80	99	99	95	92	80	98	83	82	22	
23-01795/2/7/11_EG	90	99	100	97	96	99	99	100	98	98	95	95	94	83	83	82	82	85	84	86	81	23	




Figure 3: Phylogenetic analysis of IBV/KFS/2061 (KY072832) isolated from Kafrelsheikh province, Egypt, based on partial S1 gene nucleotide sequences with Genbank located reference sequences.

4. Discussion

Infectious bronchitis disease is one of the main causes of respiratory disease in chicken worldwide leading to severe economic losses to the poultry section (Broomand et al., 2012). IBV is endemic in most of poultry sites, with a mortality rate reaching up to 100% while mortalities differs from 20% to 30% or even more and these are commonly due to mixed infections with other pathogens such as Mycoplasma and E coli. Mortalities of up to 25% are reported in young chicks affected nephritis caused by other IBV serotypes (Ignjatovic and Sapats 2000).

In the current study, we report the isolation of an Egyptian IBV isolate; IBV/KFS/2016 from chicken flocks experiencing respiratory and renal symptoms with relatively high mortality rates. The virus was isolated from Kafrelsheikh province in the Northern part of Egypt. All the collected samples were inoculated into ECE and confirmed by PCR analysis.

Inoculation of tissue suspensions of the collected samples in SPF chicken eggs resulted in stunted growth of the embryos which agreed with Cook et al., 2012 who showed that the best method of IBV isolation is to passage a sample in 9-11-day-old embryonated SPF chicken eggs via the allantoic

cavity route. The virus causes characteristic stunting and curling of the embryo after 5-7 days post inoculation. Due to the great similarity in the symptoms of upper respiratory diseases, rapid and accurate diagnostic methods for identification of the causative pathogens should be applied (Rafiei et al., 2009). PCR test is the most sensitive and fast test for the identification of infections from clinical specimens which may show negative results due to improper sampling technique or loss of viral infectivity during transportation (Vapret et al., 2000). The positive samples were confirmed as IBV through PCR assay which resulted in a 403 bp product by using specific S1 primers. Only 10/22 of the samples were confirmed as IBV viruses. Although, the increased mortality that associated with the disease described here, might be due to additional factors which increased the severity such as mixed infection with other pathogens as AIV, NDV, E coli or Mycoplasma.

Nucleotide sequence variation in S1 gene is commonly used for recognizing different IBV serotypes as the emergence of new variants is the result of few amino acid changes in the S1 glycoprotein leading to severe outbreaks (Abdel-Moneim et al 2006).

In Egypt, IBV variants have been recognized firstly in 1950s by isolation of a variant closely related to the Dutch variant D3128 (A. Ballal et al., 2005). While in 2006 Abdel-moneim et al., 2006 isolated an Egyptian strain of IBV 'EgyptF/03' which was closely related to the Massachusetts, H120, Beaudette-US, and M41 serotypes.

S1 sequence analysis of our isolate IBV/KFS/2016 (KY072832) revealed its close relatedness to IBV-EG/13280F-SP1-2013 (KU979008.1) (92% nucleotide and 90% amino acid identities), Eg/CLEVB-2/IBV/012 (JX173488.1) (90% nucleotide and 89% amino acid identities) serotypes and the Israeli strains IS/1494/06 (EU780077.2) (90% nucleotide and 89% amino acid identities), IB VAR2-06 (JX027070.1) (90% nucleotide and 87% amino acid identities) and Israel/720/99 (AY091552.2) (90% nucleotide and 89% amino acid identities). While it but was clearly different from that of the H120 (KR605489.1), MA5 (KU736747.1) (80% nucleotide and 79% amino acid identities), and 4/91 (AF093793.1) (82% nucleotide and 86% amino acid identities) vaccinal strains commonly used in Kafrelsheikh province which explains the frequent vaccination failure. These results are in agreement with Susan et al., 2010 and Selima et al., 2012, who isolated an IBV strain closely related to the Israeli Variant-2 and IS/1494/06 strains. In addition, Awad et al., 2014 have characterized seven IBV field isolates which exhibited high nucleotide sequence similarity to variant 2 like strains; IBV-CU-2-SP1 and Eg/12120s/2012. These isolates come close to a new variant group which is suspected to be the primary cause for IBV outbreaks in Egypt.

Conclusion from this study is that IBV variant 2 emergences has been reported to be circulating in Egypt and playing an important role in IBV outbreaks in the Middle East and Egypt. To provide sufficient protection against IBV, vaccination should be applied with the homologous serotype, as vaccination with a live attenuated heterologous

strain will provide insufficient and incomplete protection (Xie *et al.*2011). Thus, it is necessary to apply scheduled diagnosis of newly originating IBV strains for modifying a successful vaccination program to avoid and control new IBV outbreaks.

References

- [1] **Abdel-Moneim, A. S., El-Kady, M.F.,Ladman, B.S., Jack Gelb, J.,2006.**S1 gene sequence analysis of a nephropathogenic strain of avian infectious bronchitis virus in Egypt *Virology Journal*, 3, 78.
- [2] **Ahmed, H.N.,1954.**Incidence and treatment of some infectious viral respiratory diseases of poultry in Egypt.Ph.D.Thesis, Fac. Vet. Med. Cairo University, Giza, Egypt.
- [3] **Awad,A. M.,SediekM.E.,El-Yamany M. E.,2014.**Isolation and Molecular Characterization of Novel IBV Isolates from Broiler Chicken Farms in Egypt.*Alexandria journal of veterinary Sciences*, 42, 74-82.
- [4] **Ballal, A., Karrar,A., El-Hussein, A.,2005.**Isolation and characterization of infectious bronchitis virus strain 4/91 from commercial layer chickens in the Sudan *Journal of Animal VeterinaryAdvanced*, 4, 910–912.
- [5] **Broomand,C.H., Aini,I., Abdulrahman, O., Khatijah, Y.,Shrifah, S.H.,2007.**Development of multiplex RT-PCR for rapid detection and subtyping viruses type A .*Proceedings of the 12th International Conference of the Association of Institutions of Tropical Veterinary Medicine - Montpellier, France 2007*.
- [6] **Cook, J.K.,Jackwood, M., Jones, R. C., 2012.**The long view: 40 years of infectious bronchitis research, *Avian Pathology*, 41, 239-250.
- [7] **De Wit,S. J., Cook, J. K.,Van der Heijden H. M.,2011.** Infectious bronchitis virus variants: a review of the history, current situation and control measures *Avian Pathology* 40, 223-235.
- [8] **El-Mahdy, S.S., Ekram, S., Ahmed, A.,2010.** Efficacy of some living classical and variant infectious bronchitis vaccines against local variant isolated from Egypt, *Natural and Science*, 10, 292–299.
- [9] **El-Shafey,A.M.,2008.**Advanced study on infectious bronchitis of chicken in Egypt.PhD Thesis, Zagazig University.
- [10] **Farsang, A., Ros C., RenströmL.H., M., Baule, C., Soós, T.,Belák S.,2002.**Molecular epizootiology of infectious bronchitis virus in Sweden indicating the involvement of a vaccine strain, *Avian Pathology*, 31, 229-236.
- [11] **Ignjatovic, J. , Sapats, S.,2000.**Avian infectious bronchitis virus. *RevueScientifiqueEt Technique*.19, 493-508.
- [12] **Jahantigh, M.,Salari, S.,Hedayati, M., 2013.** Detection of infectious bronchitis virus serotypes by reverse transcription polymerase chain reaction in broiler chickens *Springerplus* 2, 36.
- [13] **Mahgoub, K.M., Bassiouni, A.A. ,Afify,M.A. ,Rabie,N.S.,2010.**The Prevalence of Infectious Bronchitis (IB) Outbreaks in Some Chicken Farms. I. Spotlight on the Status of IB Outbreaks in Some Chicken Flocks.*Journal of American Science* ,6, 57-70.
- [14] **Mo, M.L., Hong, S.M., Kwon, H.J., Kim, I.H., Song, C.S., Kim, J.H., 2013.**"Genetic Diversity of Spike, 3a, 3b and E Genes of Infectious Bronchitis Viruses and Emergence of New Recombinants in Korea. *Viruses* 5, 550-567.
- [15] **Patel,B., Bhimani,M., Bhanderi, B., Jhala,M., 2015.** Isolation and molecular characterization of nephropathic infectious bronchitis virus isolates of Gujarat state, India. *Virusdisease*, 26, 42–47.
- [16] **Pohuang, T., Chansiripornchai, N.,Tawatsin, A. , Sasipreeyajan, J.,2009.**Detection and molecular characterization of infectious bronchitis virus isolated from recent outbreaks in broiler flocks in Thailand. *Journal of Veterinary Science* 10, 219-223.
- [17] **Rafiei, M.M.,Vasfi-Marandi, M.,Bozorgmehri-Fard, M.H. , Ghadi, S.,2009.**Identification of Different Serotypes of Infectious Bronchitis Viruses in Allantoic Fluid Samples with Single and Multiplex RT-PCR.*Iranian Journal of Virology* 3, 24-29.
- [18] **Roussan, D. A.,Totanji, W. S. ,Khawaldeh, G. Y.,2008.**Molecular Subtype of Infectious Bronchitis Virus in Broiler Flocks in Jordan. *Poultry Science* 87, 661–664.
- [19] **Selima,K., Arafa, A.S., Hussein, H.A., El-Sanousi, A.A. ,2012.**Molecular characterization of infectious bronchitis viruses isolated from broiler and layer chicken farms in Egypt during 2012.*International journal of Veterinary Science and Medicine* 1, 102-108
- [20] **Vabret, A., Sapin, G., Lezin, B., Mosnier, A., Cohen, J., Burnouf, L .,Petitjean, J., Gouarin, S., Campet, M. , Freymuth, F.,2000.**Comparison of three non-tested RT-PCR for detection of influenza American *Journal of Clinical Virology*, 17, 167-75 .
- [21] **Xie,Q., Ji, J., Xie,J., Chen,F.,Cai,M.,Xue,C.,Ma,J., Bi,Y.,2011.**Epidemiology and immunoprotection of nephropathogenic avian infectious bronchitis virus in southern China *Virology Journal*, 8, 484.
- [22] **Yashpal, S.M., Devi, P.P., Sagar, M.G., 2004.** Detection of three avian respiratory viruses by single-tube multiplex reverse transcription polymerase chain reaction assay. *Journal of Veterinary Diagnostic Investigation*, 16, 244–248.