

Review of the Level of Awareness' Towards Measures of Patients Safety among Restorative Staff at Selected Dental Clinics, Saudi Arabia

Dr. Abdulrahim Abdullah Alshehri

Ministry of Defense, Armed Forces Hospital Southern Region, Saudi Arabia

Abstract: ***Background:** acquiescence with policies and procedures of measures of patients' safety need to be extremely respected and practices by all the health care practitioners, though decisive staff adherence towards all the standards of patients' safety. **Objective:** To assess the level of awareness Towards Measures of Patients Safety among Restorative Staff at Selected Dental clinics, Saudi Arabia. **Method:** A cross-sectional survey was conducted among restorative staff at selected dental clinics, Saudi Arabia.. A 28item self-administered questionnaire was provided to 150 restorative staff in the research setting to assess their level of awareness towards measures of patients' safety. **Results:** Approximately more than one third of the participants (40.0%) were not aware about measures of patients' safety. Furthermore, majority of the participants (78.7%)** which revealed a highly statistically significantly difference ($P<0001$) were replied that they were adequacy of protective equipments within the current research setting. Concerning the assessment for the level of awareness towards measures of patients' safety. The current study results showed that majority of the participants were showed a high level of awareness towards measures of patients' safety. (86.0%) ** which represent highly significance differences ($P<0.0001$). **Conclusions:** The current study results revealed that there were high levels of awareness Towards Measures of Patients Safety among Restorative Staff at Selected Dental clinics, Saudi Arabia.*

Keywords: Restorative Staff; Awareness; Measures of Patients Safety

1. Introduction

Measures of patients' safety are meant to reduce the risk of transmission of blood borne and other pathogens from both recognized and unrecognized sources. They are the basic level of infection control precautions which are to be used, as a minimum, in the care of all patients. (Siegel et al., 2007). According to National Communicable Disease Center, universal precautions is designed to prevent the transmission of blood borne diseases such as human immune deficiency virus, hepatitis B, and other blood borne pathogens when first aid or health care is provided. Under Universal Precautions, blood and certain body fluids of all patients are considered potentially infectious.

Measures of patients' safety were initially developed in 1987 by the Centers for Disease Control and Prevention in the United States and in 1989 by the Bureau of Communicable Disease Epidemiology in Canada. The Precautions include specific recommendations for use of gloves, gowns, masks, and protective eyewear when contact with blood or body secretions containing blood is anticipated. (Al-Saigul, Fontaine, Haddad, 2002)

Health care workers in particular dental restorative staff is at risk of acquiring infection through professional exposure to Infectious diseases. The minority studies have reported on dental restorative staff adherence towards universal precautions and reported lack of adequate practices in relation to compliance towards the personnel protective equipments. (Janjua, et al., 2007).

Exposure to particular health hazards are expected to influence definite high-risk for all the health care providers. All the health care workers especially the medical staff who

are working in surgical units and Operation Theater are more required to have a reason of a better understanding in adherence with PPE usage which is significant as it provides an assessment of the efficacy of accessible preventative strategies. This could then assist to recognize the preventive variables which are likely to improve the compliance and decrease the risk of infection transmission. Then, it is possible to integrate these preventative approaches into the strategies of health care setting. (Colodner, et al., 2003 & Taneja, ., 2010)

Measures of patients' safety is the only approach so that all these infections could be prevented. Inadequate experience of dental restorative staff in performing invasive procedures, they are at particular risk of exposure to blood-borne pathogens (Chopra, et al., 2008). Dental restorative staff should have reasonable knowledge and performance in relation to adherence to personnel protective equipments. Additionally, Elliott et al. (2005), reported that dedicated training must be conducted before a dental restorative staff caring for any patient procedure particularly the ones concerning sharp devices. Physicians' compliance towards the personnel protective equipments has been reported to be with low rate. (Jawaid, Iqbal & Shahbaz, 2009).

Hazards caused by non adherence to universal precautions by the health care providers, statistics reported by the Central Register of Occupational Diseases in Poland indicates that among 314 new cases of occupational diseases in HCWs in 2005, HBV and HCV represented 42.6% of all cases. Despite the substantial reduction in HBV infection since vaccination was introduced in 1989, the incidence of HCV hepatitis in Poland is still on the increase in this occupational group. (Wacawik, siorowski & Inglot, (2003) & Wilczyn, et al., (2005).

Volume 6 Issue 1, January 2017

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

Measures of patients' safety consciousness education has not been prominent among health care workers especially the category of dental restorative staff, particularly in developing countries. To the best of our knowledge, the awareness and standardized practices with universal precautions among dental restorative staff. We, therefore, conducted this study to assess the levels of knowledge and awareness towards universal precautions among dental restorative staff during their duties at the University Hospital of the King Abdul-Aziz University, Saudi Arabia.

2. Participants and Methods

This study was conducted in February, 2016 at the Selected Dental clinics, Saudi Arabia.. The study was granted ethical approval by the Administrative Committee within the selected setting. The participants were selected from the Departments of Dental Restorative after signing an informed written consent form; the questionnaire was given to each participant. Before administration of the questionnaire, the purpose of the study was explained to each respondent and confidentiality of the information assured.

The research was carried out by one of the author who were appropriately trained in administering the informed consent and the self-report questionnaire to the health care workers. In this cross-sectional study, a structured questionnaire prepared by the authors, was administered to the participants. A 28-item self-administered structured questionnaire about awareness of measures of patient safety was devised de novo and tested. It included a full range of response options, designed to identify the practitioner's level of awareness of measures of patient safety in the selected setting. Prior to distribution of the questionnaire, a pilot study was done on a selective group of health care workers who were asked to fill out the questionnaire and return it back with their comments and criticism. Minor changes were then made to the final instrument.

The initial part of the questionnaire consisted of demographic information such as occupation, age, gender, and the marital status. The second part of the questionnaire comprised of questions regarding their awareness of measures of patient safety. This part also assessed awareness of policies regarding awareness of measures of patient safety. It took approximately 15 minutes to complete each questionnaire.

The participants' results concerning the assessment of the levels of awareness of measures of patient safety, there were three levels for answers high level, medium level and high level. The level of awareness of measures of patient safety by examining questions about: use of protective barriers such as gloves and gown, mask and protective goggles.

The data were coded and analyzed by SPSS[®] for Windows[®] ver. 12.0. Strict confidentiality was maintained. All the data were stored in computers at a secured location, with access provided only to the researchers involved in the study. The χ^2 test was used to test association between categorical

variables. A p value <0.05 (two-tailed) was considered statistically significant

3. Results

Table 1: Percentage of the level of Awareness of Measures of Patient Safety

levels of awareness of measures of patient safety	Low level	Medium Level	High Level
Awareness of measures of patient safety	60 (40.0%)	5 (3.4%)	85(57.0%)*
Awareness of Measures of Patient Safety concerning Provision of Protective Equipments	18 (12.0%)	14 (9.3%)	118 (78.7%)**
Level of awareness towards Universal Precaution	23 (15.3%)	7 (4.7%)	120 (80.0%)**
Significantly different: **p<0.0001; *p<0.01			

Table 2 showed that approximately more than one third of the participants (**40.0%**) were not aware about measures of patients' safety. Furthermore, majority of the participants (**78.7%**)** which revealed a highly statistically significantly difference (**P<0.0001**) were replied that they were adequacy of protective equipments within the current research setting. Concerning the assessment for the level of awareness towards measures of patients' safety. The current study results showed that majority of the participants were showed a high level of awareness towards measures of patients' safety. (**86.0%**) ** which represent highly significance differences (**P<0.0001**).

4. Discussion

The current study findings represent that majority of the participants were showed a high Level of adherence towards Awareness of Measures of Patient Safety concerning Provision of Protective Equipments (86.0%) ** which represent highly significance differences (P<0.0001). This results consistent with the study results carried out by Sadoh, et al, (2006) who reported that more than two third of his study sample had a high level of compliance in relation to protective equipments. Moreover, these study results is congruent with the study findings of Odujurin, Adegoke & Clin (2013), which revealed that, reported frequent use of protective gears such as gloves, eyewear, masks and aprons. More women than men reported using protective gears most times with significantly more nurses reported frequent use followed by medical technologists and medical doctors. Although, this study results is incongruent with the study results of According to Jawaid, Iqbal & Shahbaz, (2009) who reported that among medical doctors working in a tertiary care hospital in Pakistan, compliance for hand washing was 86%, for wearing gloves was 79%, masks 46%, eye goggles 25% and for using gowns/plastic aprons was 45%.35 However, there is sometimes a high rate of non-compliance among health care workers and this may be due to a lack of understanding among health care workers of how to properly use protective barriers. Although, according to Danchaivijitr, Tantiwatanapaiboon & Chokloikaew (2005), who found that the use of personal protective equipment was somewhat favorable, the concern among most of the health

care workers, particularly the porters, was that the provision of protective gears was inadequate which will interfere with the health care providers to provide personal protective equipment and the appropriate training for the correct use. Furthermore, improvement in safety equipment is needed to better protect health care workers from exposure to blood-borne pathogens.

Furthermore the study results revealed that majority of the participants(78.7%)** were showed that there was a high level concerning the adequacy in relation to the provision of protective equipments which represent highly significance differences ($P < 0.0001$). According the standards, policies & procedures of infection control, recommendations for protection against viral hepatitis(1985), and the recommendations for preventing transmission of infection with human T-lymphotropic virus type III/lymphadenopathy-associated virus in the workplace(1985), it was stated that availability of supplies and awareness programs for these standard precautions are among the main rules and regulations for better compliance.

In this study, Concerning the assessment for the level of awareness towards universal precautions the current study results showed that majority of the participants were showed a high level of awareness towards universal precautions (86.0%) ** which represent highly significance differences ($P < 0.0001$). These results congruent with the study results carried out by Beltrami (2000) and the study done by Wang, Chen & Liu (2010), who concluded that respondents who were had a high level of awareness towards universal precautions. Furthermore, these study results consistent with the findings reported by Orji (2002), who concluded that non-compliance among medical doctors and nurses are associated with insufficient knowledge, workload, forgetfulness, workplace safety and the insight that colleagues also failed to follow.

5. Summary

A cross-sectional survey was conducted among restorative staff at selected dental clinics, Saudi Arabia. A 28-item self-administered questionnaire was provided to 150 restorative staff in the research setting to assess their level of awareness towards measures of patients' safety. : Approximately more than one third of the participants (40.0%) were not aware about measures of patients' safety. Furthermore, majority of the participants (78.7%) ** which revealed a highly statistically significantly difference ($P < 0.0001$) were replied that they were adequacy of protective equipments within the current research setting. Concerning the assessment for the level of awareness towards measures of patients' safety. The current study results showed that majority of the participants were showed a high level of awareness towards measures of patients' safety. (86.0%) ** which represent highly significance differences ($P < 0.0001$). The current study results revealed that there were high levels of awareness Towards Measures of Patients Safety among Restorative Staff at Selected Dental clinics, Saudi Arabia.

6. Acknowledgements

Appreciation is hereby all the participants engaged in the study.

References

- [1] Lacerda RA. Infecção hospitalar e sua relação com a evolução das práticas de assistência à saúde. In: Lacerda RA, ed. Controle de infecção em centro cirúrgico. São Paulo: Atheneu, 2003: 9-23.
- [2] Lacerda RA, Egry EY. As infecções hospitalares e a sua relação com o desenvolvimento da assistência hospitalar: reflexões para análise práticas atuais de controle. Rev Latinoam Enfermagem 1997;5:13-23.
- [3] Beltrami EM, Williams IT, Shapiro CN, Chamberland ME. Risk and management of blood-borne infections in health care workers. Clin Microbiol Rev 2000;13(3):385-407.
- [4] Gerberding JL. Incidence and prevalence of human immunodeficiency virus, hepatitis B virus, hepatitis C virus, and cytomegalovirus among health care personnel at risk for blood exposure: final report from a longitudinal study. J Infect Dis 1994;170(6):1410-7.
- [5] Ruben FL, Norden CW, Rockwell K, Hruska E. Epidemiology of accidental needle-puncture wounds in hospital workers. Am J Med Sci 1983;286(1):26-30.
- [6] Pruss-Ustun A, Rapiti E, Hutin Y. Estimation of the global burden of disease attributable to contaminated sharps injuries among health-care workers. Am J Ind Med 2005;48(6):482-90.
- [7] Khuri-Bulos NA, Toukan A, Mahafzah A, et al. Epidemiology of needlestick and sharp injuries at a university hospital in a developing country: a
- [8] Wang FD, Chen YY, Liu CY. Analysis of sharp-edged medical-object injuries at a medical center in Taiwan. Infect Control Hosp Epidemiol 2010;21(10):656-8.
- [9] Pruss-Ustun A, Rapiti E, Hutin Y. Sharp injuries: global burden of disease from sharp injuries to health care workers Geneva, Switzerland. World Health Organization, 2003.
- [10] Orji EO, Fasubaa OB, Onwudiegwu U, Hutin Z. Occupational health hazards among health care workers in an obstetrics and gynaecology unit of a Nigerian teaching hospital. J Obstet Gynaecol 2002;22(1):75-8.
- [11] Recommendations for protection against viral hepatitis. MMWR Morb Mortal Wkly Rep 1985;34(22):313-24, 329-35.
- [12] Recommendations for preventing transmission of infection with human T-lymphotropic virus type III/lymphadenopathy-associated virus in the workplace. MMWR Morb Mortal Wkly Rep 1985;34(45):681-6, 691-5.
- [13] McCarthy GM. Universal Precautions J Can Dent Assoc 2000;66:556-7.
- [14] Update: human immunodeficiency virus infections in health-care workers exposed to blood of infected patients. MMWR Morb Mortal Wkly Rep 1987;36(19):285-9.
- [15] Acquired immunodeficiency syndrome (AIDS): precautions for health-care workers and allied

- professionals. *MMWR Morb Mortal Wkly Rep* 1983;32(34):450-1.
- [16] Garner JS. Hospital Infection Control Practices Advisory Committee. Guideline for isolation precautions in hospitals. *Infect Hosp Epidemiol* 1996;17:53-80.
- [17] Spire B, Barre-Sinoussi F, Montagnier L, Chermann JC. Inactivation of lymphadenopathy associated virus by chemical disinfectants. *Lancet* 1984;2(8408):899-901.
- [18] Martin LS, McDougal JS, Loskoski SL. Disinfection and inactivation of the human T lymphotropic virus type III/Lymphadenopathy-associated virus. *J Infect Dis* 1985;152(2):400-3.
- [19] McDougal JS, Martin LS, Cort SP, Mozen M, Heldebrant CM, Evatt BL. Thermal inactivation of the acquired immunodeficiency syndrome virus, human T lymphotropic virus-III/lymphadenopathy-associated virus, with special reference to antihemophilic factor. *J Clin Invest* 1985;76(2):875-7.
- [20] Olowu O, Oluaje E, Kehinde O. Knowledge and practice of universal precautions among final year medical and dental students in the University College of Ibadan. *Dokita* 2001;28:6-9.
- [21] Odjurin OM, Adegoke OA & Clin ,S: Awareness and blood handling practices of health care workers in Lagos. *Nig J Epidemiol* 2013;11(4):425-30.
- [22] Palenick C. Strategic planning for infection control. *J Canadian Dental Association* 2000;66:556-7.
- [23] Danchaivijitr S, Tantiwatanapaiboon Y & Chokloikaew S. Universal precautions: knowledge, compliance and attitudes of doctors and nurses in Thailand. *J Med Assoc Thai* , 2005;78 Suppl 2:S112S117.
- [24] Twitchell K. Bloodborne pathogens: what you need to know-Part I. *Journal of the American Association of Occupational Health Nurses* 2003;51:46-7.
- [25] Godin G, Naccache H, Morel S, Ebacher MF. Determinants of nurses' adherence to universal precautions for venipunctures. *Am J Infect Control* 2000;28(5):359-64.
- [26] Stein AD, Makarawo TP, Ahmad MFR. A survey of doctors' and nurses' knowledge, attitudes and compliance with infection control guidelines in Birmingham teaching hospitals. *Journal of Infection Control* 2003;54:68-73.
- [27] Trim JC, Adams D, Elliott TS. Healthcare workers' knowledge of inoculation injuries and glove use. *Br J Nurs* 2003;12(4):215-21.
- [28] Rana JS, Khan AR, Haleem AA, et al. Hepatitis C: knowledge, attitudes and practices among orthopedic trainee dental restorative staff in Pakistan. *Ann Saudi Med* 2000;20(5-6):477-9.
- [29] Wang H, Fennie K, He G, et al. A training programme for prevention of occupational exposure to bloodborne pathogens: impact on knowledge, behaviour and incidence of needle stick injuries among student nurses in Changsha, People's Republic of China. *J Adv Nurs* 2003;41(2):187-94.
- [30] Heinrich J. Occupational Safety: Selected cost and benefit implications of needle stick prevention devices for hospitals (letter to House of Representatives from US General Accounting Office), 2000.
- [31] Gerberding JL. Clinical practice. Occupational exposure to HIV in health care settings. *N Engl J Med* 2003;348(9):826-
- [32] Connington A. Has the point been made? a needlestick injury awareness survey. *Safe Gard Medical* 2002.
- [33] Alam M. Knowledge, attitude and practices among health care workers on needle-stick injuries. *Ann Saudi Med* 2002;22(5-6):396-9.
- [34] Sadoh WE, Fawole AO, Sadoh AE & Heinrich J.. Practice of universal precautions among healthcare workers. *J Natl Med Assoc* 2006;98(5):722-6.
- [35] Jawaid M, Iqbal M, Shahbaz S. Compliance with standard precautions: a long way ahead. *Journal of Public Health* 2009;38:85-8.
- [36] Evanoff B, Kim L, Mutha S, et al. Compliance with universal precautions among emergency department personnel caring for trauma patients. *Ann Emerg Med* 1999;33(2):160-5.
- [37] Gershon RR, Karkashian CD, Grosch JW, et al. Hospital safety climate and its relationship with safe work practices and workplace exposure incidents. *Am J Infect Control* 2000;28(3):211-21.
- [38] Janjua NZ, Razaq M, Chandir S, et al. Poor knowledge-predictor of nonadherence to universal precautions for blood borne pathogens at first level care facilities in Pakistan. *BMC Infect Dis* 2007;7:81.
- [39] Cutter J, Jordan S. Uptake of guidelines to avoid and report exposure to blood and body fluids. *J Adv Nurs* 2004;46(4):441-52.
- [40] Nelsing S, Nielsen TL, Nielsen JO. Noncompliance with universal precautions and the associated risk of mucocutaneous blood exposure among
- [41] Danish physicians. *Infect Control Hosp Epidemiol* 1997;18(10):692-8.
- [42] Brooks AJ, Phipson M, Potgieter A, et al. Education of the trauma team: video evaluation of the compliance with universal barrier precautions in resuscitation. *Eur J Surg* 1999;165(12):1125-8.
- [43] Richman G, Dorsey A, Stayer S, Schwartz R. Compliance With Standard Precautions Among Pediatric
- [44] Anesthesia Providers. *The Internet Journal of Anesthesiology* 2000 4(4):1-8.
- [45] Siegel JD, Rhinehart E, Jackson M, Chiarello L., The Healthcare Infection Control Practices Advisory Committee, 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings
- [46] National Communicable Disease Center. *Isolation Techniques for Use in Hospitals*. 1st ed. Washington, DC: US Government Printing Office; PHS publication no 2054 1970.
- [47] Drusin LM, Ross BG, Rhodes KH, Krauss AN, Scott RA. Nosocomial ringworm in a neonatal intensive care unit: a nurse and her cat. *Infect Control Hosp Epidemiol* 2000;21(9):605-7.
- [48] Uemura T, Kawashita T, Ostuka Y, Tanaka Y, Kusubae R, Yoshinaga M. A recent outbreak of adenovirus type 7 infection in a chronic inpatient

facility for the severely handicapped. Infect Control Hosp Epidemiol 2000;21(9):559-60.

[49] Al-Saigul AM, Fontaine RE, Haddad Q. Nosocomial malaria from contamination of a multidose heparin container with blood. Infect Control Hosp Epidemiol 2002;21(5):329-30.

[50] Colodner R, Sakran W, Miron D, Teitler N, Khavalevsky E, Kopelowitz J. *Listeria monocytogenes* cross-contamination in a nursery. Am J Infect Control 2003;31(5):322-4.