Assessment of the Prothotodontic Staff Knowledge towards the Methods of Transmission of the Communicable Diseases at Selected Prothotodontic Units, Saudi Arabia

Dr. Abdulmajed Mansour Saad Alqarni¹, Dr. Mastour Safar Alshahrani²

^{1, 2}Dental Intern, Dental College - King Khalid University, Abha

Abstract: Communicable Diseases (CD) is one of the frequently widespread infections affecting the elderly patients. The aim of this article was to assess prothotodontic staff knowledge towards the methods of transmission of the CD. Looking forwards prevention strategies for improving the management of CD in elderly prothotodontic inpatients.

Keywords: Communicable Diseases, Knowledge & Prothotodontic Units

1. Methods

A descriptive research design was utilized in the current research. A total of 150 prothotodontic staff at selected Prothotodontic units was given a structured questionnaire On June, 2016. They were asked to indicate their knowledge towards the methods of transmission of the CD. For each item concerning the knowledge towards the methods of transmission of the CD the proportion of Prothotodontic staff perspectives towards measuring their level of knowledge about the methods of was calculated. Descriptive statistics like percentage was used to describe the findings using SPSS 20.

2. Results

The current research study objectives were to assess the Prothotodontic staff perspective toward methods of transmission of the CD.Mean knowledge about of items pertaining to Prothotodontic staff towards the methods of transmission of the CD within the study setting was (73.1%).

3. Conclusions

Concerning the Prothotodontic staff knowledge toward the methods of transmission of the CD, contained by the study setting, the data findings showed that; there was a highest level of prothotodontic staff knowledge towards the modes of transmission of Communicable Diseases within the selected setting.

4. Introduction

Today, infectious diseases is consider as one of the most common evidence based concepts to be investigated and involved in most of the prothotodontic researches .Communicable Diseases(CD) is one of the frequently widespread infections affecting the elderly patients. Nevertheless; CD is over diagnosed and over treated in most of the health care setting all over the world. (Rao & Patel ,2009). According to McMurdo & Gillespie, (2000), hospitalization put patient at associated risks, including risk of infection. Nosocomial infections, or hospital-associated infections, are estimated to occur in Five percent of all acute care hospitalizations, or two million cases per year. Hospitalassociated infections have been identified as one of the most serious patient safety issues in health care. Postponing urination, pregnancy, menopause, frequent intercourse, records on urinary infection during childhood are considered as predisposing factor to the infection. A quarter of women who reported urinary infection for the first time would be reinfected in less than 3 months. Early diagnose of urethral infection was considered very important to prevent relapse and indication of its complications such as rental failure, adhesions among pregnant women, abortion and early delivery

Woodford & George, (2009), mentioned that infections that be converted into clinically obvious after forty eight hours of hospitalization are well thought-out hospitalassociated. Risks factors for hospital-associated infections are generally categorized into three areas: iatrogenic, organizational, or patient-related. Iatrogenic risk factors include invasive procedures like intubation, indwelling vascular lines, urine catheterization and antibiotic use and prophylaxis. Organizational risk factors include such things as contaminated air-conditioning systems, contaminated water systems, staffing, and physical layout of the facility as open beds close together). Examples of patient-related risk factors include severity of illness, immunosuppressant, and length of stay.

Attributable to increased risk of infection associated with urinary catheters, a number of practices have been evaluated in an attempt to reduce the incidence of urinary catheterrelated infections. These include alternative approaches to use of urinary catheters and antimicrobial urinary catheters. (CD) is known as a common infection among women; in more than 95 percent of cases urinary infection is created by a certain group of bacteria; the major pathogen of urinary infection is E.coli bacteria, which is a normal bacterial flora in intestines. It seems that the reasons for increased prevalence of such infections among women are proximity of their urethral and anus, short length of urethral and warm and wet environment of perinea (Boscia, et al., 2012).

5. Methods

A descriptive research design was utilized in the current research. A total of 150 prothotodontic staff of selected prothotodontic units, Saudi Arabia, were given a structured questionnaire during June 2016. This number constitute, more than two third of the Prothotodontic staff of the time of data collection. All wards representing units concerning all the subspecialties' were consider as a setting for the current research. The Prothotodontic staff at each unit was contacted to discuss the importance of the research and the study protocol. An anonymous 20-items questionnaire was adopted from valid & reliable tool using the guidelines from a Taxonomy for Augustine (1993). All questions had fixed answer categories.

The questionnaire was pilot-tested and dealt with the following topics: demographic data; data concerning

knowledge towards the methods of transmission of the CD.. Descriptive statistics like percentage was used to describe the findings using SPSS 20. All categories data were analyzed using the Chi-squared test with or without Yates' correction; Statistical significance for all analyses was presumed for P at 0.05.

6. Results

The current research study objectives were to assess the prothotodontic staff perspective toward methods of transmission of the CD. The Mean knowledge about of items pertaining to prothotodontic staff towards the methods of transmission of the CD within the study setting was (73.1%).

Regarding the prothotodontic staff knowledge toward the methods of transmission of the CD, contained by the study setting, the data findings showed that; there was a highest level of prothotodontic staff knowledge towards the modes of transmission of Communicable Diseases within the selected setting.

Table 1: Frequency and Percentage of items pertaining to Prostodontic Staff towards the methods of transmission of the
Communicable Diseases

		Frequency of	Percentage of
Sl. No.	Knowledge of Items	correct Response	correct Responses
	Hospital acquired infections are the result of self infection, cross infection &		
1	environmental infection	20	40
2	The single most important measure for preventing HAI is hand washing	43	86
3	The common causative organism of CD is E. Coli	37	74
4	HAI's are transmitted through body fluids, staff hands and reusable equipment	45	90
5	Immunization is not a universal precaution	44	88
6	Hemoglobin less than 11 gm % is not a sign of infection.	48	96
7	Patients receiving immunosuppressive therapy are more susceptible to HAI	44	88
8	In case of CD, bacterial count of 10 CFU/ml of Urine c/s is significant	22	44
9	Sterilization is a process of killing of microorganism including spores	43	86
10	Glutehyde is to be changed every 28 days	45	90
11	More than 15 CFU organisms are grown in an agar plate	9	18
12	Sterile technique is not necessary for nose gastric feeding.	41	82
13	The most important factor involved in hand washing is friction	24	48
14	HAI is synonymous to nosocomial infection	39	78
15	Moisture enhances the trans-mission of micro- organisms	48	96
16	Boiling is a method of sterilization	16	32
17	Chemical disinfection is the best method of sterilizing surgical instruments	38	76
18	If a person doesn't show sign and symptoms of disease, he can't transfer a disease	49	98
19	Alcohol is an effective disinfectant when rubbed in skin.	39	78
20	CD's are one of the most common forms of HAI	37	74

Mean knowledge about of items pertaining to Prostodontic Staff towards the methods of transmission of the Communicable Diseases within the study setting was (73.1%).

7. Discussion

The response rate to the current research was excellent, suggesting that this was an area of importance for prothotodontic staff. In this research, more than two third of the prothotodontic staff were replied that there was a high level of knowledge towards the transmission of Communicable Diseases compared to a study conducted by Ciavarelli, Figlock & Sengupta who studied the factors in aviation of the staff awareness in relation to methods of CD transmission it revealed that the hospital staff reported (46%) of the prothotodontic staff in relation to organizational culture safety. Moreover, In Saudi Arabia, it was reported that there was a lack of skills in relation to communicable diseases of the universal measures to prevent the transmission of CD Communicable Diseases are a common infection among women and is followed by several complications. The aim of this study was determine the prevalence and factors influencing prevention of Communicable Diseases based on health belief model. A cross-sectional study was conducted among 166 female students in Shadegan city, Iran; which was randomly selected with the proportional to size among different girls high school at Shadegan city for participation in this study. A structured questionnaire was applied for collecting data

and data were analyzed by SPSS version 20 using bivariate correlations and logistic regression statistical tests. Almost 22.3% of the participants had history of experience Communicable Diseases at last one in lifelong. The best predictor for Communicable Diseases was perceived susceptibility with odds ratio estimate of 0.810 [95% CI: 0.664, 0.987]. Based on our result, it seems that designing and implementation of educational programs to increase susceptibility about chance of getting Communicable Diseases among the female students may be useful in order to prevent Communicable Diseases.

Prevalence of CD between women is high; in this regard, Eriksson et al in their study reported 29.6% of women were diagnosed as having CD at least one. Furthermore, Sorto et al reported the CD incidence among the women in their study was of 35.8%. Studies showed the prevalence of CD in different parts of the world, which made it necessary to determine and control its predisposing factors. Along this, personal health behaviors were considered as the major predisposing factors to relapse of urinary infection. Various factors are reflected in human behavior and it is vital to know the causality network to affect impressive factors on behavior, which has been investigated through years by behaviorists. Knowing effective factors on behavior showed to be operational in designing educational programs.

On the other hand, studies on designing education programs showed that the most effective educational programs were based on theory-based approaches rooted in behavior change patterns. The very first step to plan an education program is to choose a proper pattern or theory in health education while effective health education directly depends on the dominance on using best theories and proper approaches in each event. Several studies, also, reported that it was essential to focus on mental factors in health education comprehensive preventive programs as mediating and predicators of behavior; as the result health care providers should be aware of various effective factors on fulfilling preventive and health improving behaviors to be able to design and implement advantageous programs in the field. Hereof, it is worth to mention that health belief model (HBM) is an analytical behavior theory, which has been used in several studies on different fields of preventive behaviors.

8. Conclusions

The current research aimed to assess the prothotodontic staff knowledge towards the methods of transmission of the CD. Looking forwards prevention strategies for improving the management of CD in elderly prothotodontic inpatients. The response rate to the current research was excellent, suggesting that this was an area of importance for prothotodontic staff. In this research, more than two third of the prothotodontic staff were replied that there was a high level of knowledge towards the transmission of Communicable Diseases

The current study recommends utilization of continuous educational measures to enhance the prothotodontic and other Health care providers to increase their knowledge towards the methods of transmission of the CD. Furthermore applying performance feedback and increased availability of specialist equipments are imperative for improving management of CDs in the elderly.

9. Acknowledgements

Thanks to all the interns helps in performing the current research

References

- [1] Rao GG, Patel M. Communicable Diseases in hospitalized elderly patients in the United Kingdom: the importance of making an accurate diagnosis in the post broad-spectrum antibiotic era. J Antimicrob Chemother 2009;63:5–6
- [2] McMurdo MET, Gillespie ND. Communicable Diseasesin old age: over-diagnosed and over-treated. Age Ageing 2000;29:297–8.
- [3] Woodford HJ, George J. Diagnosis and management of Communicable Diseasesin hospitalized older people. J Am Geriatr Soc 2009;57:107–14.
- [4] Boscia JA, Kobasa WD, Knight RA, et al. Lack of association between bacteriuria and symptoms in the elderly. Am J Med 2012;81:979-82.
- [5] Scottish Intercollegiate Guidelines Network (SIGN). Management of suspected bacterial Communicable Diseasesin adults. Edinburgh: SIGN publication no. 88. July 2012. http://www.sign.ac.uk/guidelines/fulltext/88/index.html (accessed 20 January 2014).
- [6] Scottish Intercollegiate Guidelines Network (SIGN). Algorithm: suspected CD in older people, management of suspected bacterial Communicable Diseasesin adults. Edinburgh: SIGN publication no. 88. July 2012. http://www.sign.ac.uk/pdf/sign88_algorithm_older.pdf (accessed 20 January 2014.
- [7] Levit K, Ryan K, Elixhauser A, et al. HCUP facts and figures: statistics on hospital-based care in the United States in 2005. Rockville, MD: Agency for Healthcare Research and Quality; 2007. Available at: <u>www.hcupus.ahrq.gov/reports.jsp</u>.
- [8] Nguyen QV. Hospital acquired infections. Emedicine. 2004. [Accessed March 10, 2006]. <u>http://www.emedicine.com/PED/topic1619.htm</u>.
- [9] Stone PW, Larson E, Kawar LN. A systematic audit of economic evidence linking nosocomial infections and infection control interventions: 1990–2000. Am J Infect Control. 2002;30(3):145–52. [PubMed].
- [10] de Jonge E. Effects of selective decontamination of digestive tract on mortality and antibiotic resistance in the intensive-care unit. Curr Opin Crit Care. 2005;11(2):144–9. [PubMed].
- [11] Centers for Disease Control and Prevention. Guidelines for preventing health-care associated pneumonia, 2003: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. MMWR. 2004;53:1–35.
- [12] Safdar N, Dezfulian C, Collard HR, et al. Clinical and economic consequences of ventilator-associated pneumonia: a systematic review. Crit Care Med. 2005;33:2184–93. [PubMed].
- [13] Healthcare Infection Control Practices Advisory Committee and Hand-Hygiene Task Force, Society for

Healthcare Epidemiology of America, Association for Professionals in Infection Control, Epidemiology, & Infection Diseases Society of America. Guideline for hand hygiene in healthcare settings. J Am Coll Surg. 2004;198:121–7. [PubMed].

- [14] Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings: recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Infect Control Hos Epidemiol. 2002;23:S3– 40. [PubMed]
- [15] Kac G, Podglajen I, Gueneret M, et al. Microbiological evaluation of two hand hygiene procedures achieved by healthcare workers during roCDne patient care: a randomized study. J Hosp Infect. 2005;60:32– 9. [PubMed].
- [16] Thomas JA, McIntosh JM. Are incentive spirometry, intermittent positive pressure breathing, and deep breathing exercises effective in the prevention of postoperative pulmonary complications after upper abdominal surgery? A systematic overview and metaanalysis including commentary by Dean E with author response. Phys Ther. 1994;74(1):3–16. [PubMed].
- [17] Overend TJ, Anderson CM, Lucy SD, et al. The effect of incentive spirometry on postoperative pulmonary complications: a systematic review. Chest. 2001;120:971–8. [PubMed].
- [18] Chumillas S, Ponce JL, Delgado F, et al. Prevention of postoperative pulmonary complications through respiratory rehabilitation: a controlled clinical study. Arch Phys Med Rehabil. 1998;79:5– 9. [PubMed].
- [19] Hall JC, Tarala RA, Tapper J, et al. Prevention of respiratory complications after abdominal surgery: a randomised clinical trial. BMJ. 1996;312:148– 52. [PMC free article] [PubMed].
- [20] Westerdahl E, Lindmark B, Eriksson T, et al. Deepbreathing exercises reduce atelectasis and improve pulmonary function after coronary artery bypass surgery. Chest. 2005;128:3482–8. [PubMed].
- [21] Mackay MR, Ellis E, Johnston C. Randomised clinical trial of physiotherapy after open abdominal surgery in high risk patients. Aust J Physiother. 2005;51:151– 9. [PubMed].
- [22] Craven DE, Kunches LM, Kilinsky V, et al. Risk factors for pneumonia and fatality in patients receiving continuous mechanical ventilation. Am Rev Respir Dis. 1986;133:792–6. [PubMed].
- [23] Kollef MH. Ventilator-associated pneumonia. A multivariate analysis. JAMA. 1993;270:1965– 70. [PubMed].
- [24] Kollef MH. The identification of ICU-specific outcome predictors: a comparison of Prothotodontic, Prothotodontic, and cardiothoracic ICUs from a single institCDon. Heart Lung. 1995;24:60–6. [PubMed].
- [25] Torres A, Serra-Batlles J, Ros E, et al. Pulmonary aspiration of gastric contents in patients receiving mechanical ventilation: The effect of body position. Ann Intern Med. 1992;116:540–3. [PubMed].
- [26] Orozco-Levi M, Torres A, Ferrer M, et al. Semirecumbent position protects from pulmonary aspiration but not completely from gastroesophageal

reflux in mechanically ventilated patients. Am J Respir Crit Care Med. 1995;152:1387–90. [PubMed]