

Carotid Body Tumour Radiological Diagnosis and Surgical Management - Our Experience

Dr. Aditya Nutakki¹, Dr. Harivadan Lukka², Dr. Manitha Kondru³

¹Associate Professor of Radio-diagnosis, GayatriVidyaParishad Health Care and Medical Technology, Visakhapatnam, India

²Assistant Professor of Cardio-thoracic Surgery, Andhra Medical College, Visakhapatnam, India
Corresponding author Email: [dr_harivadan\[at\]yahoo.com](mailto:dr_harivadan[at]yahoo.com)

³Associate Professor of Cardio-thoracic Surgery, Andhra Medical College, Visakhapatnam, India

Abstract: ***Background:** Carotid Body Tumours are rare. Despite over a century of describing surgery, controversy remains surrounding details of their treatment. Most complications are considered technically preventable. **Patients and Methods:** This retrospective study included 13 cases diagnosed as Carotid Body Tumours. Patients were subjected to clinical evaluation, carotid duplex ultrasound, computed tomography angiography scan, and sometimes MR angiography. All patients underwent CBT resection and when needed vascular reconstruction. Patients were classified into three groups depending on Shamblin classification. **Results:** Mean operative time was longer in patients of Shamblin class 3 (5.4±0.77 h; range, 4.5–6 h; P<0.001). Regarding vascular reconstruction, all patients of Shamblin class 3 were reconstructed by interposition vein graft and two patients of Shamblin class 1 were reconstructed through primary repair, but patients of Shamblin class 2 were reconstructed by all means. In postoperative follow-up complications, bleeding was more significantly in Shamblin class 3 (P=0.032). Cranial nerve deficit was observed in four (33.3%) patients, and all of them were transient. The mean size of excised tumor was 5.17±1.4 cm, with range of 3–7.5 cm. **Conclusion:** Early surgical excision by a cardiovascular surgeon is the only proven cure for CBTs, because they are small and easy to remove. However, if neglected, there will be many challenges, as most tumors can become locally invasive. Meticulous periadventitial (and sometimes subadventitial) dissection can greatly reduce the rates of complications.*

Keywords: Carotid Body tumours (CBT), Shamblin Classification

1. Introduction

The carotid body is the largest chemoreceptor tissue mass located within the periadventitia of posterior surface of carotid bifurcation. Normal carotid body is ovoid in shape and ~5 mm in its longest dimension. It usually receives arterial supply from external carotid artery and derives sensory innervation from glossopharyngeal nerve [1,2]. Carotid body tumors (CBTs) have many names, such as chemodectomas, paragangliomas, and glomustumors. They represent neoplasia of chemoreceptive tissue. These tumors belong to paragangliomas, which occur along autonomic ganglion chain. In head and neck, CBTs are the most common paragangliomas [3]. Macroscopically, they are reddish brown and rubbery. Although they lack a true capsule, they are highly vascular and can invade adventitia of carotid vessels. As the tumors enlarge, they splay carotid artery branches, giving 'lyre sign' on angiography [4].

Most carotid chemodectomas are benign slowly growing tumors. Patients usually present with pulsatile neck masses. The exact incidence of malignant transformation is unknown. Malignancy here is based mainly on clinical behavior more than histology. Even benign tumors are capable of aggressive invasion into adjacent structures. If carotid body cells are found in lymph nodes biopsied during surgery, this is diagnostic for malignancy. Some cases have been reported with hypertension, resolved after tumor resection, which may be caused by tumor-related catecholamine release [1,5–7]. Conventional angiography was traditionally the mainstay and provides excellent details especially about vascularity of the lesion. Duplex ultrasound has been considered the most important, noninvasive

method to examine carotid lesions with no discomfort or risk. Recently, computed tomography angiography and MR angiography (MRA) have been used instead of conventional angiography. These methods provide high levels of anatomic and vascular details [1, 8, 9]. Shamblin and colleagues at Mayo Clinic developed a classification system for carotid chemodectomas that depends on tumor extent and neurovascular involvement. Group I tumors are smaller, so can be dissected easily. Group II tumors are larger, adherent to adventitia and partially surround vessels. Group III tumors are more adherent and encase internal and external carotid arteries, so dissection is difficult in the periadventitial plane [10,11]. Surgical resection is the gold standard treatment for carotid chemodectomas. Although these lesions tend to be small, they are best treated early. The first successful carotid chemodectoma resection was done by Scudder in the United States. He was able to preserve carotid system, and there were no significant complications [8, 12, 13]. Surgeons must plan their operative approach according to the tumor characteristics, and attempts must be made to reduce blood loss and preserve vital structures as these tumors are highly vascular and carry high risk of hemorrhage during surgery. The lesions may also adhere to or envelop neurovascular structures [1,14]. Radiation therapy is another treatment modality for CBTs, but it remains controversial owing to conflicting studies and lack of long-term follow-up. There is currently no effective chemotherapy for these tumors [1, 12].

2. Patients and Method

After approval by the local ethical committee of King George Hospital and GayatriVidyaParishad Health care and Medical Technology and obtaining written fully informed consent from the patients, this retrospective was conducted on patients diagnosed with Carotid Body Tumour at the Cardio-Vascular and thoracic Surgery Department of King George Hospital and GayatriVidyaParishad Health care and Medical Technology from August 2015 till January 2020; the enrollment period was 42 months, and the patient follow-up period was 12 months.

All patients were subjected to detailed clinical evaluation, laboratory assessment, carotid duplex ultrasound, computed tomography angiography scan, and sometimes MRA examination. All patients underwent CBT resection followed by vascular reconstruction if needed. Patients were classified depending on Shamblin classification into three groups. Patients included in this study were fit for general anesthesia with resectable tumor or if tumors demonstrated airway compression. However, patients excluded from this study were patients with significant comorbidities, with limited life expectancy, with contralateral cranial nerve dysfunction, or with irresectable tumor.

3. Operative Procedure

The procedure was performed general anesthesia. Cerebral monitoring and protection were done with internal carotid artery occlusion or reconstruction using routine shunting by Pruitt-Inahara shunt.

Patient was kept supine with head rotated to the contralateral side. Shoulder roll was placed to extend the neck, and table head was elevated 10–15° to reduce venous pressures and incisional blood loss. A longitudinal skin incision was made along anterior border of sternomastoid muscle. If needed, the incision was extended proximally toward sternal notch and distally toward mastoid process. The dissection continued until carotid sheath appears. Then we opened this sheath along anterior border of internal jugular vein. Common facial vein typically joins internal jugular vein near the carotid bifurcation and was a useful landmark; we divided this vein. Here, laryngeal nerves were protected. Proximal common carotid artery was circumferentially mobilized in the periadventitial plane, with sufficient length for bypass shunt. We proceeded dissection distally on the lateral aspect of carotid system until beyond the upper end of the mass where we circumferentially mobilized internal carotid artery. Here hypoglossal nerve was carefully separated and protected. Then dissection was continued down along medial aspect of internal carotid artery till bifurcation. At this point, we circumferentially mobilized the external carotid artery.

For small tumors (Shamblin group I), we conserved the external carotid artery; branches of external carotid artery supplying the tumor were divided and ligated. Once the posterior, lateral, and medial aspects of the tumor were mobilized, the superior extent was divided between the ligature and the mass removed. Primary repair was done for small tears.

For larger tumors that were adherent to the external carotid artery (Shamblin group II), we tried to conserve the External Carotid Artery (ECA) first, but if tedious, it was divided at its origin, and proximal stump was oversewn to reduce blood loss and provide a handle to rotate the tumor helping dissection. Patients of Shamblin class 2 were reconstructed by all means.

For tumors that could not be dissected from the bifurcation (Shamblin group III), both internal and external carotid arteries were removed with the tumor. Vascular reconstruction for the internal carotid was performed with autologous vein interposition graft using shunt through the graft and removing the shunt before the second anastomosis was completed. In addition, if clamping of carotid arteries was performed, the patient received systemic heparin.

Follow-up

A preliminary neurologic assessment was performed when patient awakens from anesthesia. Patient was brought to recovery room where blood pressure and neurologic function could be carefully monitored. Care was done for neck hematomas. Drain was removed after 24 h. Patients were followed closely for 12 months.

Statistical analysis

The study data were tabulated and analyzed using SPSS, version 16 soft ware (SPSS Inc., Chicago, Illinois, USA). Categorical data were presented as number and percentages, and analyzed using Fisher's exact test. Continuous data were expressed as mean \pm SD and range. Differences between groups were tested using analysis of variance (F test) for variables or Kruskal–Wallis test for nonparametric ones ($P < 0.05$ was considered significant)

4. Results

No mortality related to surgery was reported. One patient missed follow-up, and this patient was excluded from the study data analysis. So at completion of the study, clinical data were available for 12 patients. Their age range was 35–69 years old.

Overall, seven (58.3%) patients were females and five (41.7%) patients were males. All patients were fit for general anesthesia depending on American Society of Anesthesiologists (ASA I and ASA II). Patients were classified depending on Shamblin classification into three groups: seven (58.3%) patients were class 1, three (25%) patients were class 2, and two (16.7%) patients were class 3.

Neck mass was the main presenting complaint in 11 (91.7%) patients; only one (8.3%) patient was incidental finding. Sensory changes (pulsatile tinnitus, numbness, and face or neck pain) were observed in five (41.7%) patients. Duration of symptoms was 1.25 ± 0.46 years. Bilateral CBT was reported only in one (8.3%) patient

All patients passed uneventful intraoperative course without complications. Mean operative time was longer in patients of Shamblin class 3 (mean, 5.4 ± 0.77 h; range, 4.5–6 h; $P < 0.01$). Intraoperative blood loss was also more in patients of

Shamblin class 3. Mean duration of postoperative hospital stay (in days) was longer in the patients of Shamblin class 3 (mean, 9.3 ± 4.1 ; range, 6–14 days; $P=0.003$)

Regarding vascular reconstruction, all patients of Shamblin class 3 were reconstructed by interposition vein graft and two patients of Shamblin class 1 were repaired primarily, but three patients of Shamblin class 2 were reconstructed by all means. The mean size of excised tumor was 5.17 ± 1.4 , with range of 3–7.5 cm. Most tumors were benign paraganglioma [10 (83.4%)], and only one (8.3%) tumor was malignant, with positive surgical margin and lymph node invasion. There was a histological surprise in one (8.3%) patient, having lymph node hyperplasia.

Challenges in the postoperative follow-up were as follows: bleeding was observed in three (25%) patients in the form of hematoma or ecchymosis; only one (8.3%) patient needed re-exploration to ligate slipped ligature of common facial vein, and others were managed conservatively. Stroke was reported in two (16.7%) patients; both patients were transferred to neurological ICU for management. Cranial nerve deficit was observed in four (33.3%) patients; all of them were transient and treated by neurotonic drugs for up to 6 months and consisted of superior laryngeal nerve in one patient in the form of choking, recurrent laryngeal nerve in one patient in the form of hoarseness of voice, hypoglossal nerve in one patient in the form of deviated tongue to the same side of injury, and marginal mandibular nerve in one patient in the form of wry lip. Infection was treated in one (8.3%) patient by proper antibiotics

5. Discussion

Surgery of carotid chemodectoma is the gold standard for cure. These tumors carry a high risk of massive hemorrhage during resection because they are highly vascular. The lesions may also adhere to or envelop neurovascular structures, increasing the risk of complications [8].

These tumors are best treated early for several reasons: smaller tumors are easier to remove; most tumors can become locally invasive; lesions may grow distally into the skull or involve cranial nerves, complicating resection; and these tumors may develop malignant behavior [1, 12, 13].

Because complications are common, these morbidities dramatically can be decreased by meticulous dissection and adequate control of carotid system [1]. In the current study, most patients' age was more than 40 years old; despite the exact age incidence of CBT being unknown, they are usually common in the age of 30–50 years old. There was near-equal sex distribution in this series despite sex predominance not being established [1]. This age and sex distribution was comparable to that mentioned by Ma et al. [15], who reported their study on 18 patients (female: eight and male: 10, mean age: 52 years). The current study patients were more than that mentioned by Zhu et al. [16], who reported 11 cases, and they mentioned that CBT is reported at 50–70 years old, with higher female incidence. Patients can be anesthetized either cervical block or general anesthesia; in this study, general anesthesia was preferred owing to better airway management and reduced patient

movement during the operation as mentioned by Warren et al. [1]. Patients were classified depending on Shamblin classification into three groups; seven (58.3%) patients were class 1, three (25%) patients were class 2, and two (16.7%) patients were class 3. This was near similar to that mentioned by Ma et al. [15], who reported 10 (55.6%) cases could be classified into Shamblin I, five (27.8%) cases were Shamblin II, and only three (16.6%) cases were Shamblin III. Most reported carotid chemodectomas are symptomatic mass, which is similar to the current study. A neck mass was the main symptom in 11 (91.7%) patients. Duplex ultrasound imaging, computed tomography angiography, or MRA allowed for adequate operative planning [8,9,17]. Regarding treated side in this study, bilateral CBT was reported only in one (8.3%) patient, right CBTs were reported in four (33.3%) patients, and left CBTs were reported in seven (58.3%) patients. This was comparable to that mentioned by Ma et al. [15], who observed unilateral mass in 17/18 (94.4%) cases, with 13 masses in the left and five masses in the right side and bilateral in 1/18 (5.6%) case.

By reviewing operative events, mean operative time was longer in patients of Shamblin class 3 (5.4 ± 0.77 h; range, 4.5–6 h; $P < 0.001$) Mean intraoperative blood loss was more also in patients of Shamblin class 3 (963.3 ± 464.7 ; range, 700–1500 ml; $P=0.004$). Mean duration of postoperative hospital stay (in days) was longer in the patients of Shamblin class 3 (9.3 ± 4.1 ; range, 6–14 days; $P=0.003$). This was similar to that mentioned by Warren et al. [1] and Power and Hallet [18] but less than that mentioned by Davila et al. [8], where the mean time for resected carotid chemodectoma was 224.5 min (52–696 min), and the mean blood loss was 143.9 ml (10–2000 ml).

Regarding vascular reconstruction, all patients of Shamblin class 3 were reconstructed by interposition vein graft and two patients of Shamblin class 1 were repaired primarily, but the three patients of Shamblin class 2 were reconstructed by all means. This result was comparable to that observed by Davila et al. [8], where 20 (10.9%) patients underwent reconstruction: primary closure in eight cases, saphenous interposition graft in six cases, and patch closure in six cases. The mean size of excised tumor was 5.17 ± 1.4 , with range of 3–7.5 cm. Most histological types were benign paraganglioma in 32 (94.2%) and only one type was malignant one (2.9%) and another case one (2.9%) was histological surprise, that is, lymph node hyperplasia. This tumor size was comparable to that reported by Ma et al. [15], where the resected tumor was larger than 3.5 cm and smaller than 8 cm. Risk of malignancy of carotid chemodectoma is less than 5% [19]. This was reported also by Davila et al. [8]. One case developed pulmonary secondaries after resection by 2 years. However, Ma et al. [15] found neither recurrence nor metastasis in 11-year follow-up. This study confirms that patients can safely undergo excision, and most complications are considered technically preventable, and adequate planning by the surgeon can greatly reduce the rates of adverse events. The current study complication percentage was more as compared with that observed by Davila et al. [8] Postoperative complications were observed in 36 (19.7%) patients: cranial nerve deficit in 28 (15.2%), stroke in two

(1.0%), and hematoma in three (1.6%). However, Ma et al. [15] reported no nerve injuries, except hoarse voice, which recovered later on. Although recurrence was rare, patients should be followed closely after discharge [1, 18].

6. Conclusion

Early surgical excision by a cardio-vascular surgeon is the only proven cure for CBTs because they are small and easy to remove. However, if neglected, there will be many challenges as most tumors can become locally invasive. Meticulous periadventitial (and sometimes subadventitial) dissection can greatly reduce the rates of complications.

7. Financial Support and Sponsorship

Nil

8. Conflicts of Interest

There are no conflicts of interest.

References

- [1] Warren BC, Wesley SM, Glenn ML. Carotid body tumors' in rutherford's vascular surgery and endovascular therapy. In: Anton NS, Bruce AP, eds. Rutherford's Vascular Surgery. 9th ed. Philadelphia, PA: Elsevier; 2018. 1: 4193–4215.
- [2] Abu-Ghanem S, Yehuda M, Carmel N, Abergel A, Fliss D. Impact of preoperative embolization on outcomes of carotid body tumor surgery: a meta-analysis and review of the literature. *Head Neck* 2016; 38(Suppl 1): E2386–E2394.
- [3] Wieneke J, Smith A. Paraganglioma: carotid body tumor. *Head Neck Pathol* 2009; 3:303–306.
- [4] Chinwe I, Valeria P. Enhancing carotid space mass. *J Am Osteopath CollRadiol* 2015; 4:24–26.
- [5] Junior NM, Silva RS, Ribeiro JH, Batista LC, Bringhentti EM, Brunini de Souza BB, Cabral LC. Carotid body tumor (paraganglioma): report of two cases treated surgically. *J Vasc Bras* 2016; 15:158–164.
- [6] Zhang TH, Jiang WL, Li YL. Perioperative approach in the surgical management of carotid body tumors. *Ann VascSurg* 2012; 26:775–782.
- [7] Zeng G, Zhao J, Ma Y, Huang B, Yang Y, Feng H. Comparison between the treatments of functional and nonfunctional carotid body tumors. *Ann VascSurg* 2012; 26:506–510.
- [8] Davila VJ, Chang JM, Stone WM, Richard J, Fowl RJ, Bower TC, et al. Current surgical management of carotid body tumors. *J VascSurg* 2016; 6:1704–1708.
- [9] Tong Y. Role of duplex ultrasound in the diagnosis and assessment of carotid body tumor: a literature review. *Intractable Rare Dis Res* 2012; 1:129–133.
- [10] Arya S, Rao V, Juvekar S, Dcruz A. Carotid body tumors: objective criteria to predict Shamblin group on MR imaging. *Am J Neuroradiol* 2008; 29:1349–1354.
- [11] Shamblin WR, ReMine WH, Sheps SG, Harrison EG. Carotid body tumor (chemodectoma). Clinicopathologic analysis of ninety cases. *Am J Surg* 1971; 122:732–739.
- [12] Künzel J, Koch M, Brase C, Fietkau R, Iro H, Zenk J. Treatment of cervical paragangliomas: is surgery the only way? *Am J Otolaryngol* 2014; 35:186–191.
- [13] Fruhmann J, Geigle J, Konstantiniuk P, Cohnert T. Paraganglioma of the carotid body: treatment strategy and SDH-gene mutations. *Eur J VascEndovascSurg* 2013; 45:431–436.
- [14] Ozay B, Kurc E, Orhan G, Yucel O, Senay S, Tademir M. Surgery of carotid body tumor: 14 cases in 7 years. *ActaChirBelg* 2008; 108:107–111.
- [15] Ma D, Liu M, Yang H, Ma X, Zhang C. Diagnosis and surgical treatment of carotid body tumor: a report of 18 cases. *J Cardiovasc Dis Res* 2011; 1:122.
- [16] Zhu J, Han D, Zhou J. Retrospective analysis of the diagnosis and treatment of 11 cases of carotid body tumor. *Lin Chuang Er Bi Yan HouKeZaZhi* 2005; 19:817–819.
- [17] Pellitteri PK, Rinaldo A, Myssiorek D, Gary Jackson C, Bradley PJ, Devaney KO. Paragangliomas of head and neck. *Oral Oncol* 2004; 40:563–575.
- [18] Power AH, Hallett JW. Carotid body tumors. In Stanley JC, Veith FJ, Wakefield TW, eds. *Current therapy in vascular and endovascular surgery*. 5th ed. Philadelphia: Elsevier; 2014. 141–145
- [19] Liapis CD, Evangelidakis EL, Papavassiliou VG, Kakisis JD, Gougoulakis AG, Polyzos AK. Role of malignancy and preoperative embolization in management of carotid body tumors. *World J Surg* 2000; 24:1526–1530