

Emulgel: A Review

Fenil Vanpariya¹, Milan Shiroya², Mitesh Malaviya³

^{1, 2, 3}Shree Dhanvantary Pharmacy college, Kim- Surat, India

¹Correspondence author's Email: [fenilvanpariya35316\[at\]gmail.com](mailto:fenilvanpariya35316[at]gmail.com)

Abstract: *Many advantages of gels a major limitation is in the delivery of hydrophobic drugs. So to overcome this limitation an emulsion based approach is being used so that even a hydrophobic therapeutic moiety can enjoy the unique properties of gels. When gels and emulsions are used in combined form the dosage form are referred as emulgel. In recent years, there has been great interest in the use of novel polymers. A unique aspect of dermatological pharmacology is the direct accessibility of the skin as a target organ for diagnosis and treatment. The combination of hydrophilic cornified cells in hydrophobic intercellular material provides a barrier to both hydrophilic and hydrophobic substances. Within the major group of semisolid preparations, the use of transparent gels has expanded both in cosmetics and in pharmaceutical preparations. Polymer can function as emulsifiers and thickeners because the gelling capacity of these compounds allows the formulation of stable emulsions and creams by decreasing surface and interfacial tension and at the same time increasing the viscosity of the aqueous phase. In fact, the presence of a gelling agent in the water phase converts a classical emulsion into an emulgel. These emulgel are having major advantages on novel vesicular systems as well as on conventional systems in various aspects. Various permeation enhancers can potentiate the effect; Soemulgels can be used as better topical drug delivery systems over present systems. The use of emulgels can be extended in analgesics and antifungal drugs.*

Keywords: Emulgels, Topical drug delivery

1. Introduction

Topical drug administration is a localized drug delivery system anywhere in the body through ophthalmic, rectal, vaginal and skin as topical routes. These are apply a wide spectrum of preparations for both cosmetic and dermatological, to their healthy or diseased skin.¹ These formulations range in physicochemical nature from solid through semisolid to liquid. Drug substances are seldom administered alone, but rather as part of a formulation, in combination with one or more non medicated agents that serve varied and specialized pharmaceutical function. Drugs are administered topically for their action at the site of application or for systemic effects.² Drug absorption through the skin is enhanced if the drug substance is in solution, if it has a favourable lipid/water partition coefficient and if it is a nonelectrolyte. For the most part, pharmaceutical preparations applied to the skin are intended to serve some local action and as such are formulated to provide prolonged local contact with minimal systemic drug absorption. Drug applied to the skin for their local action include antiseptics, antifungal agent, skin emollients and protectant. The main advantages of topical delivery system are to bypass first pass metabolism. Avoidance of the risks and inconveniences of intravenous therapy and of the varied conditions of absorption like pH changes, presence of enzymes, gastric emptying time are other advantages of topical preparations.³ ⁴ The topical drug delivery system is generally used where the others system of drug administration fails or it is mainly used in fungal infection. Human skin is a uniquely engineered organ that permit its terrestrial life by regulating heat and water loss from the body whilst preventing the ingress of noxious chemicals or microorganisms. It is also the largest organ of the human body, providing around 10% of the body mass of an average person, and it covers an average area of 1.7 m². Whilst such a large and easily accessible organ apparently offers ideal and multiple sites to administer therapeutic agents for both local and systemic actions, human skin is a highly efficient self-repairing barrier designed to keep the inside in and the outside out.⁵

Gels are a relatively newer class of dosage form created by entrapment of large amounts of aqueous or hydroalcoholic liquid in a network of colloidal solid particles, which may consist of inorganic substances, such as aluminum salts or organic polymers of natural or synthetic origin.⁶ They have a higher aqueous component that permits greater dissolution of drugs, and also permit easy migration of the drug through a vehicle that is essentially a liquid, compared with the ointment or cream base.⁷ These are superior in terms of use and patient acceptability. In spite of many advantages of gels a major limitation is in the delivery of hydrophobic drugs. So to overcome this limitation, emulgels are prepared and used so that even a hydrophobic therapeutic moiety can enjoy the unique properties of gels.

In fact, the presence of a gelling agent in the water phase converts a classical emulsion into an emulgel.¹² Both oil-in-water and water-in-oil emulsions are used as vehicles to deliver various drugs to the skin. Emulgels for dermatological use have several favorable properties such as being thixotropic, greaseless, easily spreadable, easily removable, emollient, nonstaining, long shelf life, bio-friendly, transparent & pleasing appearance.¹

Use of topical agents requires an appreciation of the factors that influence percutaneous absorption.¹⁴ Molecules can penetrate the skin by three routes: through intact stratum corneum, through sweat ducts, or through sebaceous follicle. The surface of the stratum corneum presents more than 99% of the total skin surface available for percutaneous drug absorption.¹⁵

Passage through this outer most layer is the rate limiting step for percutaneous absorption. The major steps involved in percutaneous absorption include the establishment of a concentration gradient, which provides the driving force for drug movement across the skin, release of drug from the vehicle (partition coefficient), and drug diffusion across the layers of the skin (diffusion coefficient). Preferable characteristics of topical drugs include low molecular mass

Volume 10 Issue 3, March 2021

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

(600 Da), adequate solubility in oil and water, and a high partition coefficient. Except for very small particles, water soluble ions and polar molecules do not penetrate intact stratum corneum. Topical formulation can be used to manipulate the barrier function of the skin, for example, topical antibiotics and antibacterials help a damaged barrier toward off infection, sun screening agents and the horny layer protect the viable tissues from Ultraviolet radiation and emollient preparations restore pliability to a desiccated horny layer.¹⁶

During development of semi-solid preparations for cutaneous application whose formulation contains an antimicrobial preservative, the need for and the efficacy of the chosen preservative shall be demonstrated to the satisfaction of the competent authority. A suitable test method together with criteria for judging the preservative properties of the formulation are provided in efficacy of antimicrobial preservation. Sterile semi-solid preparations for cutaneous application are prepared using materials and methods designed to ensure sterility and to avoid the introduction of contaminants and the growth of microorganisms.¹⁷

The efficacy of an antimicrobial preservative may be enhanced or diminished by the active constituent of the preparation or by the formulation in which it is incorporated or by the container and closure used. Preparation for topical use should have microbiological quality and it is checked with test for sterility. Total viable aerobic count should not be more than 10² micro-organisms (aerobic bacteria plus fungi) per gram. It should not have more than 10¹ enterobacteria, certain other gram-negative bacteria per gram and completely devoid of *Pseudomonas aeruginosa* and *Staphylococcus aureus*.¹⁸⁻¹⁹

This project is to reveal the material and method used doesn't imparts any microbial contamination and the methyl paraben 0.2% used is sufficient to maintain its sterility. (micro biology)

2. Rationale

Many widely used topical agents like ointment, cream, lotion have many disadvantages. They have very sticky causing uneasiness to the patient when applied. Moreover they also have lesser spreading coefficient and need to apply with rubbing. And they exhibit the problem of stability also. Due to all these factors within the major group of semisolid preparations, the use of transparent gels has expanded both in cosmetics and in pharmaceutical preparations.

A gel is colloid that is typically 99% wt liquid, which is immobilized by surface tension between it and a macromolecular network of fibers built from a small amount of a gelating substance present. In spite of many advantages of gels a major limitation is in the delivery of hydrophobic drugs. So to overcome this limitation an emulsion based approach is being used so that even a hydrophobic therapeutic moiety can be successfully incorporated and delivered through gels.²⁰

Drug delivery across the skin

The epidermis is the most superficial layer of the skin and is composed of stratified keratinised squamous epithelium which varies in thickness in different parts of the body. It is thickest on with elastic fibres. The skin forms a relatively waterproof layer that protects the deeper and more delicate structures. Blood vessels are distributed profusely beneath the skin. Especially important is a continuous venous plexus that is supplied by inflow of blood from the skin capillaries. In the most exposed areas of the body-the hands, feet, and ears blood is also supplied to the plexus directly from the small arteries through highly muscular arteriovenous anastomoses. A unique aspect of dermatological pharmacology is the direct accessibility of the skin as a target organ for diagnosis and treatment. The skin acts as a two-way barrier to prevent absorption or loss of water and electrolytes. There are three primary mechanisms of topical drug absorption: transcellular, intercellular, and follicular. Most drugs pass through the torturous path around corneocytes and through the lipid bilayer to viable layers of the skin. The next most common (and potentially under recognized in the clinical setting) route of delivery is via the pilosebaceous route. The barrier resides in the outermost layer of the epidermis, the stratum corneum, as evidenced by approximately equal rates of penetration of chemicals through isolated stratum corneum or whole skin. Creams and gels that are rubbed into the skin have been used for years to deliver pain medication and infection fighting drugs to an affected site of the body. These include, among others, gels and creams for vaginal yeast infections, topical creams for skin infections and creams to soothe arthritis pain. New technologies now allow other drugs to be absorbed through the skin (transdermal). These can be used to treat not just the affected areas (for example, the skin) but the whole body. (systemic)

Factors Affecting Topical Absorption of Drug²¹⁻²²

Physiological Factors

- 1) Skin thickness.
- 2) Lipid content.
- 3) Density of hair follicles.
- 4) Density of sweat glands.
- 5) Skin pH.
- 6) Blood flow.
- 7) Hydration of skin.
- 8) Inflammation of skin

Physiochemical Factors

- 1) Partition coefficient.
- 2) Molecular weight (<400 dalton).
- 3) Degree of ionization (only unionized drugs gets absorbed well).
- 4) Effect of vehicles

Factors to be considered when choosing a Topical Preparation²³⁻²⁴

- 1) Effect of the vehicle e.g. an occlusive vehicle enhances penetration of the active ingredient and improves efficacy. The vehicle itself may have a cooling, drying, emollient or protective action.

- 2) Match the type of preparation with the type of lesions. For example, avoid greasy ointments for acute weepy dermatitis.
- 3) Match the type of preparation with the site.(e.g., gel or lotion for hairy areas)
- 4) Irritation or sensitization potential. Generally, ointments and w/o creams are less irritating, while gels are irritating. Ointments do not contain preservatives or emulsifiers if allergy to these agents is a concern.

Method to Enhance Drug Penetration and Absorption ²⁵

- 1) Chemical enhancement
- 2) Physical enhancement
- 3) Biochemical enhancement
- 4) Supersaturation enhancement

Advantages ²⁶⁻²⁷

- 1) Hydrophobic drugs can be easily incorporated into gels using d/o/w emulsions. Most of the hydrophobic drugs cannot be incorporated directly into gel base because solubility act as a barrier and problem arises during the release of the drug. Emulgel helps in the incorporation of hydrophobic drugs into the oil phase and then oily globules are dispersed in aqueous phase resulting in o/w emulsion. And this emulsion can be mixed into gel base. This may be proving better stability and release of drug than simply incorporating drugs into gel base.
- 2) Better stability: Other transdermal preparations are comparatively less stable than emulgels. Like powders are hygroscopic, creams shows phase inversion or breaking and ointment shows rancidity due to oily base.
- 3) Better loading capacity: Other novel approaches like niosomes and liposomes are of nano size and due to vesicular structures may result in leakage and result in lesser entrapment efficiency. But gels due to vast network have comparatively better loading capacity.
- 4) Production feasibility and low preparation cost: Preparation of emulgels comprises of simpler and short steps which increases the feasibility of the production. There are no specialized instruments needed for the production of emulgels. Moreover materials used are easily available and cheaper. Hence, decreases the production cost of emulgels.
- 5) No intensive sonication: Production of vesicular molecules need intensive sonication which may result in drug degradation and leakage. But this problem is not seen during the production of emulgels as no sonication is needed.
- 6) Controlled release: Emulgels can be used to prolong the effect of drugs having shorter t_{1/2}.

Important Constituents of Emulgel Preparation

- 1) **Aqueous Material:** This forms the aqueous phase of the emulsion. Commonly used agents are water, alcohols.²⁸
- 2) **Oils:** These agents form the oily phase if the emulsion. For externally applied emulsions, mineral oils, either alone or combined with soft or hard paraffins, are widely used both as the vehicle for the drug and for their occlusive and sensory characteristics. Widely used oils in oral preparations are nonbiodegradable mineral and castor oils that provide a local laxative effect, and fish liver oils or various fixed oils of vegetable origin

(e.g., arachis, cottonseed, and maize oils) as nutritional supplements.²⁹⁻³⁰

- 3) **Emulsifiers:** Emulsifying agents are used both to promote emulsification at the time of manufacture and to control stability during a shelf life that can vary from days for extemporaneously prepared emulsions to months or years for commercial preparations.eg Polyethylene glycol 40³¹ stearate, Sorbitan monooleate³² (Span 80), Polyoxyethylenesorbitan monooleate (Tween 80)³³, Stearic acid³⁴, Sodium stearate.³⁵
- 4) **Gelling Agent:** These are the agents used to increase the consistency of any dosage form can also be used as thickening agent.³⁶⁻³⁷
- 5) **Permeation Enhancers:** These are agents that partition into and interact with skin constituents to induce a temporary and reversible increase in skin permeability.³⁸

Emulgel Preparation

Emulgel was prepared by the method reported by Mohammad et al (2004) with minor modification. The Gel in formulations were prepared by dispersing Carbopol 934 in purified water with constant stirring at a moderate speed and Carbopol 940 in purified water with constant stirring at a moderate speed then the pH are adjusted to 6 to 6.5 using Tri ethanol amine (TEA).

The oil phase of the emulsion were prepared by dissolving Span 20 in light liquid paraffin while the aqueous phase was prepared by dissolving Tween 20 in purified water. Methyl and Propyl paraben was dissolved in propylene glycol whereas drug was dissolved in ethanol and both solutions was mixed with the aqueous phase. Both the oily and aqueous phases were separately heated to 70° to 80°C; then the oily phase were added to the aqueous phase with continuous stirring until cooled to room temperature. And add Glutaraldehyde in during of mixing of gel and emulsion in ratio 1:1 to obtain the emulgel.³⁹

Characterization of Gellified Emulsion

Physical appearance: The prepared Emulsion formulations were inspected visually for their color, homogeneity, consistency and pH. The pH values of 1% aqueous solutions of the prepared Gellified Emulsion were measured by a pH meter (Digital pH meter DPH 115 pm).⁴⁰

Spreadability: Spreadability is determined by apparatus suggested by Mutimer et al (1956) which is suitably modified in the laboratory and used for the study. It consists of a wooden block, which is provided by a pulley at one end. By this method, spreadability is measured on the basis of 'Slip' and 'Drag' characteristics of emulgels. A ground glass slide is fixed on this block. An excess of emulgel (about 2 gm) under study is placed on this ground slide. The emulgel is then sandwiched between this slide and another glass slide having the dimension of fixed ground slide and provided with the hook. A 1 Kg weight is placed on the top of the two slides for 5 minutes to expel air and to provide a uniform film of the emulgel between the slides. Excess of the emulgel is scrapped off from the edges. The top plate is then subjected to pull of 80 gms. With the help of string attached to the hook and the time (in seconds) required by the top

slide to cover a distance of 7.5 cm be noted. A shorter interval indicate better spreadability. Spreadability was calculated by using the formula,

$$S = M.L/T$$

Where, S = spreadability,

M = Weight tied to upper slide,

L = Length of glass slides

T = Time taken to separate the slides completely from each other.

Extrudability study

It is a usual empirical test to measure the force required to extrude the material from tube. The method applied for determination of applied shear in the region of the rheogram corresponding to a shear rate exceeding the yield value and exhibiting consequent plug flow. In the present study, the method adopted for evaluating emulgel formulation for extrudability is based upon the quantity in percentage of emulgel and emulgel extruded from lacquered aluminum collapsible tube on application of weight in grams required to extrude at least 0.5 cm ribbon of emulgel in 10 seconds. More quantity extruded better is extrudability. The measurement of extrudability of each formulation is in triplicate and the average values are presented. The extrudability is then calculated by using the following formula:

Extrudability = Applied weight to extrude emulgel from tube (in gm) / Area (in cm²)

Globule size and its distribution in emulgel

Globule size and distribution was determined by Malvern zetasizer. A 1.0 gm sample was dissolved in purified water and agitated to get homogeneous dispersion. Sample was injected to photocell of zetasizer. Mean globule diameter and distribution was obtained.⁴³

Rheological Study

The viscosity of the different emulgel formulations is determined at 25°C using a cone and plate viscometer with spindle 52 (Brookfield Engineering Laboratories,) and connected to a thermostatically controlled circulating water bath.

Swelling Index

To determine the swelling index of prepared topical emulgel, 1 gm of gel is taken on porous aluminum foil and then placed separately in a 50 ml beaker containing 10 ml 0.1 N NaOH. Then samples were removed from beakers at different time intervals and put it on dry place for some time after it reweighed. Swelling index is calculated as follows:

$$\text{Swelling Index (SW) \%} = [(W_t - W_0) / W_0] \times 100$$

Where, (SW) % = Equilibrium percent swelling, W₀ =

Original weight of emulgel at zero time after time t, W_t =

Weight of swollen emulgel

Ex-vivo Bioadhesive strength measurement of topical emulgel

(MICE SHAVEN SKIN): The modified method is used for the measurement of bioadhesive strength. The fresh skin is cut into pieces and washed with 0.1 N NaOH. Two pieces of skin were tied to the two glass slide separately from that one glass slide is fixed on the wooden piece and other piece is tied with the balance on right hand side. The right and left

pan were balanced by adding extra weight on the left-hand pan. 1 gm of topical emulgel is placed between these two slides containing hairless skin pieces, and extra weight from the left pan is removed to sandwich the two pieces of skin and some pressure is applied to remove the presence of air. The balance is kept in this position for 5 minutes. Weight is added slowly at 200 mg/ min to the left-hand pan until the patch detached from the skin surface. The weight (gram force) required to detach the emulgel from the skin surface gave the measure of bioadhesive strength. The bioadhesive strength is calculated by using following:

$$\text{Bioadhesive Strength} = \text{Weight required (in gms)} / \text{Area (cm}^2\text{)}$$




Figure 1: Setup for bioadhesive test

Drug Content Determination

Drug concentration in Gellified Emulsion was measured by spectrophotometer. Drug content in Gellified Emulsion was measured by dissolving known quantity of Gellified Emulsion in solvent (methanol) by Sonication. Absorbance was measured after suitable dilution in UV/VIS spectrophotometer (UV-1700 CE, Shimadzu Corporation, Japan).⁴⁶

In Vitro Release Study

Franz diffusion cell (with effective diffusion area 3.14 cm² and 15.5 ml cell volume) was used for the drug release studies. Gellified Emulsion (200 mg) was applied onto the surface of egg membrane evenly. The egg membrane was clamped between the donor and the receptor chamber of diffusion cell. The receptor chamber was filled with freshly prepared PBS (pH 5.5) solution to solubilize the drug. The receptor chamber was stirred by magnetic stirrer. The samples (1.0 ml aliquots) were collected at suitable time interval. Samples were analyzed for drug content by UV visible spectrophotometer after appropriate dilutions. Cumulative corrections were made to obtain the total amount of drug release at each time interval. The cumulative amount of drug released across the egg membrane was determined as a function of time.⁴⁷

Microbiological assay

Ditch plate technique was used. It is a technique used for evaluation of bacteriostatic or fungistatic activity of a compound. It is mainly applied for semisolid formulations. Previously prepared Sabouraud's agar dried plates were used. Three grams of the Gellified Emulsion are placed in a ditch cut in the plate. Freshly prepared culture loops are streaked across the agar at a right angle from the ditch to the edge of the plate. After incubation for 18 to 24 hours at 25°C, the fungal growth was observed and the percentage inhibition was measured as follows.

$$\% \text{ inhibition} = L_2 / L_1 \times 100$$

Where L₁ = total length of the streaked culture, and

L₂ = length of inhibition.

Skin irritation test

A 0.5 gm sample of the test article was then applied to each site (two sites per rabbit) by introduction under a double gauze layer to an area of skin approximately 1" x 1" (2.54 x 2.54 cm²). The Gellified Emulsion is applied on the skin of rabbit. Animals were returned to their cages. After a 24 hour exposure, the Gellified Emulsion is removed. The test sites were wiped with tap water to remove any remaining test article residue.

Accelerated stability studies of Gellified Emulsion

Stability studies were performed according to ICH guidelines. The formulations were stored in hot air oven at 37 ± 2°, 45 ± 2° and 60 ± 2° for a period of 3 months. The samples were analyzed for drug content every two weeks by UV-Visible spectrophotometer. Stability study was carried out by measuring the change in pH of gel at regular interval of time.⁴⁸

3. Conclusion

In the coming years, topical drug delivery will be used extensively to impart better patient compliance. Since emulgel possesses an edge in terms of spreadibility, adhesion, viscosity and extrusion, they will become a popular drug delivery system. Moreover, they will become a solution for loading hydrophobic drugs in an water soluble gel bases.

References

- [1] Kshirsagar N A. Drug Delivery Systems. Ind. J. Pharmacol. 2000; 32:S54- S61.
- [2] Rashmi M. Topical gel: A review august vol. 2008; available from [http:// www.pharmainfo.com](http://www.pharmainfo.com)
- [3] Sharma S. Topical preparations are used for the localized effects at the site of their application by virtue of drug penetration into the underlying layers of skin or mucous membranes. Pharmaceutical reviews 2008; 6:1
- [4] Laithy HM. and El shaboury KMF. The development of Cutinalipogels and gel microemulsion for topical administration of fluconazole. Ame Pharm Sci. PharmSciTech. 2003; 3:10 25.
- [5] McGrath JA, Eady R & Pope Fm.chapter 3 anatomy and organization of human skin, p 3.1 3.15
- [6] Kumar L, Verma R. *In vitro* evaluation of topical gel prepared using natural polymer. International Journal of Drug Delivery 2010; 2:58-63.
- [7] Gennaro AR, ed. Remington: the Science and Practice of Pharmacy. Easton, Mack Publishing Company 19th ed.; 1995.
- [8] Ansel HC, Allen LV Jr., Popovich NG. Pharmaceutical Dosage Forms and Drug Delivery Systems. New York Lippincott Williams and Wilkins 7th ed.; 1999.
- [9] Topical Emulsion- Gel Composition Comprising Diclofenac Sodium. Patent no. WO/2004/017998).
- [10] Mohamed MI. Optimization of ChlorphenesinEmulgel Formulation. AAPS J. 2004; 6 (3).
- [11] Gupta A, Mishra AK, Singh AK, Gupta V, Bansal P. Formulation and evaluation of topical gel of diclofenac sodium using different polymers. Drug Invention Today 2010; 2:250-253.
- [12] Rieger MM, Lachman L, Lieberman HA, Kanig JL. The Theory and Practice of Industrial Pharmacy. 3rd ed., PA Lea and Febiger, Philadelphia; 1986. pp. 502-533
- [13] Stanos SP. Topical Agents for the Management of Musculoskeletal Pain. J Pain Symptom Manage 2007; 33.
- [14] Stanos SP. Topical Agents for the Management of Musculoskeletal Pain. J Pain Symptom Manage March 2007; 33.
- [15] Jain A, Deveda P, Vyas N, Chauhan J et al. Development Of Antifungal Emulsion Based Gel For Topical Fungal Infection(S). IJPRD 2011; 2(12).
- [16] Bruton L, Keith P, Blumenthal D, Buxton Z. Goodman & Gillman's Manual of Pharmacology and Therapeutics. Mc Graw's Hill. 2008. pp.1086-1094.
- [17] Kohei Kyuki, Tomohisa Shibuya, Kaito Tsurumi Hajime Fujimura, "Anti-Inflammatory effect of diclofenacsodium ointment (cream) in topical application" by Japan J. Phamacol. 33 121-132 (1983.)
- [18] <http://www.dermweb.com/therapy/common.htm> "Principle of Skin Therapy"
- [19] British Lee, Fang-Yu (Tw) Chen, ShanChiung (Tw) Chen, Topical formulation comprising a NSAID, preferably diclofenac, for alleviating pain/inflammation caused by herpes virus infection published on September 2004, European Patent Application EP1457202 Kind Code: A2.
- [20] Cecv G. Preclinical characterisation of NSAIDs in ultradeformable carriers or conventional topical gels. International journal of pharmaceutics; 2008.
- [21] Kalia YN, Guy RH. Modeling transdermal drug release. Adv Drug Deliv Rev. 2001, 48:159-72.
- [22] Ayub, CA, Gomes ADM, Lima MVC, Vianna- Soares CD, FerreiraLMA. Topical Delivery of Fluconazole: In Vitro Skin Penetration and Permeation Using Emulsions as Dosage Forms Drug. Dev. Ind. Pharm. 2007; 33:273- 280.
- [23] Gaur PK, Mishra S, Purohit S, Dave K. Transdermal Drug Delivery System: A Review. AJPCR 2009; 2: 14-20.
- [24] Subranayam N, Ghosal SK, Moulik SP. Enhanced In Vitro Percutaneous Absorption and In Vivo Anti-Inflammatory Effect of a Selective Cyclooxygenase Inhibitor Using Microemulsion. Drug Dev. and Industrial Pharm., 2005.
- [25] Pathan, I.B.; Setty, C.M. Chemical penetration enhancers for transdermal drug delivery systems. Trop J Pharm Res. April 2009; 8:173-179.
- [26] Rashmi, MS. Topical Gel: A Review, 2008. Available from: [http://www.pharmainfo.net/reviews/topic algel-review](http://www.pharmainfo.net/reviews/topic%20algel-review).
- [27] Djordjevic J, Michniak B, Uhrich, Kathryn E AAPS PharmSciTech 2003; 5(4):1-12.
- [28] Lachman, L.; Lieberman, H.A. The Theory and Practice of Industrial Pharmacy. 3rd Ed. Varghese Publishing house; 1990. pp. 534.
- [29] Vyas, S.P.; Khar, R.K. Controlled Drug Delivery. 1st Ed. Vallabh Prakashan; 2002. pp. 416-417.
- [30] Bonacucina G, Cespi M, Palmieri GF. Characterization and Stability of Emulsion Gels Based on

- Acrylamide/Sodium AcryloyldimethylTaurate Copolymer AAPS PharmSciTech. June 2009; 10 (2).
- [31] Curr AEB. Transdermal Drug Delivery: Penetration Enhancement Techniques Heather. Drug Deliv. 2005; 2:23-33.
- [32] Rutrer N. Drug absorption through the skin: a mixed blessing .Arch Dis Child 1987; 62:220-221.
- [33] Zhang XL, Zhao R, Qian W. Preparation of an emulgel for treatment of aphthous ulcer on the basis of carbomers. Chin. Pharm. J. 1995; 30:417-418.
- [34] Swarbrick, J. Encyclopedia of pharmaceutical technology, 3rd ed., 1551.
- [35] Gibson, M. Pharmaceutical formulation and preformulation ,Interpharm 2004.
- [36] Mortazavi SA, Aboofazeli R. An Investigation into the Effect of Various Penetration Enhancers on Percutaneous Absorption of Piroxicam. Iranian Journal of Pharmaceutical Research 2003; 135-140.
- [37] Kumar, L.; Verma, R. Int. J Drug Delivery 2010,58-63.
- [38] Jacob SW, Francone CA. Structure and Function of Man, (2). Mohamed MI. Optimization of Chlorphenesin emulgel formulation. *Aaps*, 6:1-7, 2004. WB Saunders Co. Philadelphia, 1970, 55-60.
- [39] Kasliwal N, Derle D, Negi J, Gohil J. Effect of permeation enhancers on the release and permeation kinetics of meloxicam gel formulations through rat skin. Asian Journal of Pharmaceutical Sciences 2008, 3 (5): 193-199
- [40] Sanjay, Jain BD, Padsalg A, Patel K, Mokale V, Formulation, development and evaluation of Fluconazole gel in various polymer bases, Asi. J. Pharm., 2007; 1: 63 –68
- [41] Gondaliya DP and Pundarikakshudu K. *Indian drugs*, 39: 465-473, 2002.
- [42] Gupta GD, Gound RS. Release rate of nimesulide from different gellants. Indian J Pharm Sci., 61, 1999, 229-234.
- [43] Jones DB, Woolfson AD, Brown AF. Textural, Masmoudi H, Piccerelle P, Le Dréau Y, Kister J. A rheological method to evaluate the physical stability of highly viscous pharmaceutical oil-in-water emulsions. Pharm Res 2006;23 8:1937–47 ts
- [44] viscoelastic and mucoadhesive properties of pharmaceutical gels composed of cellulose polymers. Int J Pharm 1997; 151:223–33.
- [45] Chaudhari P, Ajab A, Malpure P, Kolsure P, Sanap D, Development and in-vitro evaluation of thermo reversible nasal gel formulations of Rizatriptan benzoate, Indian J. Pharm. Edu. Res., 2009; 43: 55-62.