

Review on Herblaxmitaru

Mahesh Jaywant Patil¹, Dr. Mrunal Shirsat²

¹Research Scholar, Madhav University, Pindwara Sirohi. Rajasthan, India
Mail id: mpatilphd.madahvuniversity[at]gmail.com

²Dean, Madhav University, Pindwara Sirohi Rajasthan, India

Abstract: *Greatest of drugs typically used now are of herbal origin. Generally known as Lakshmitaru in Kannada Language belongs from family Simaroubaceae, it is known as Paradise tree, Bitter ash, dysentery-bark, in English. Pharmacognostical studies on the leaves of a well-known anti-cancerous drug, commonly known as Lakshmi taru in India is botanically associated to Simarouba glauca DC. The bark and leaf extract of Simarouba is well known for its varied types of pharmacological assets such as hemostatic, anthelmintic, antiparasitic, anti-dysenteric, anticancer. The bark is used to cure fever, malaria, stomach and bowel disorders.*

Keywords: Simarouba glauca, Laxmitaru, Paradise, Anticancer

1. Introduction

Western medicine however treats a very minor share of the people on earth. Eighty percent of world population be contingent on totally on local medicines made closely from plants. It is assessed that among 35,000 and 70,000 diverse species of plants have been used by various people of the world. Really, into the 20th century much of the Pharmacopoeia of technical medicine was subsequent from the herbal wisdom of native populations. In emergent world, and utility of ordinary product of medicinal elements is better in last decade over synthetic medicine, since of toxic effects. WHO now optional stimulates traditional herbal remedies in NHC programs since herbal drug easily available at low cost and comparatively safe. Greatest of drugs typically used now are of herbal origin. Simarouba glauca DC (SG), even sized tree exotic to India is generally known as Lakshmitaru in Kannada Language belongs from family Simaroubaceae, it is known as Paradise tree, Bitter ash, dysentery-bark, in English, Aceituno in Spanish, and so on. Pharmacognostical studies on the leaves of a well known anti-cancerous drug, commonly known as Lakshmi taru in India is botanically associated to Simarouba glauca DC. Its place to the family Simbouraceae, innate of South, Central America, native to the Amazon rain forest and other humid areas of South America like Mexico, Cuba, Haiti, Jamaica, Brazil, Bahamas, Puerto Rico, El-Salvador, Guatemala, while it is unusual to India, Sri Lanka, Philippines and Myanmar. Common name: Paradise, Aceituno, Bitter wood tree, Malayalam: Lakshmi tharu, Tamil: Shorgum Maram, Hindi: Luxmitaru, Scientific Classification, Kingdom: Plantae, Family: Simaroubaceae, Genus: Simarouba, Species: S. Glauca.

Cultivation and collection:

It was a regular sized tree regularly influences a height about 20 m and trunk diameter about 50-80 cm and life about 70 years. It grows below a wide range of agroclimatic situations similar warm, humid and tropical regions. Its farming depends upon rainfall distribution at around 400 mm, water holding capacity of the soil. Simarouba glauca one of the vital herbal drugs used in inconsistency of dysentery hence its bark is also known as dysentery bark..

Chemical constituents:

Seeds of S. glauca are opulent in palatable fat that is cast off for cooking in tropical countries. Fruit is good basis of vegetable oil which is improbable source of fat soluble vitamins like A and E, The hard fraction rich in steric acid, palmitic acids can be used as coco-butter substitutes. S. glauca has the expense of quassinoids, an alkaloid 8-hydroxyl canthin-6-one. Stated that the seeds contain 40% kernels, the kernels comprise 60% fat, a good source of fat. Protein content in the Simarouba kernels was 18.2g/100g which augmented to 47.7g/100g in a defatted meal of Simarouba. Studies of stated that deoiled meal of S. glauca is a rich source in-vitro protein digestibility (88%) and amino acid-based total nutritive indices.

Seeds include 50-65% oil that can be captivated out by traditional methods. Major components are oleic (52-54%), stearic (27-33%) and palmitic (11-12%). Protein values (45.6-56.8g/100g; regular, 51.8g/100g) in their deoiled meal cake. Throughout the study of Simarouba from dissimilar sources has described a range of protein standards in their deoiled meal cake. Similarly, crude fiber content of Simarouba kernel augmented to 11.8g/100g in its deoiled meal. Major mechanisms are oleic, stearic and palmitic are sticky matter, volatile oil, malic acid, gallic acid in actual small amount, an ammoniacal salt calcium malate, oxalate, ferric oxide, silica, ulmin, and lignin, afforded six canthin-6-one type alkaloid offshoots, canthin-6-one (1), 2-methoxycanthin-6-one (2), 9-methoxycanthin-6-one (3), 2-hydroxycanthin-6-one, 4,5-dimethoxycanthin-6-one and 4,5-dihydroxycanthin-6-one, a limonoid, melianodiol, an acyclic squalene-type triterpenoid, 14-deacetylurylene, two coumarins, scopoletin and fraxidin, and two triglycerides, triolein and trilinolein. 14-Deacetylurylene. Selectively active in illogicality of the human lung cancer cell line but was inactive in an in-vivo deep fiber assay using this same cell type roots, stems and fruits of Simarouba. Isolated quassinoids, triterpenoids, a mixture of steroids, flavonoid, and kaempferol.

Pharmacological and Clinical Aspects:

Pancreatic cancer is one of the most lethal human malignancies. Nearly 100% of cases of pancreatic cancer. Glauca rubinone was originally developed as an

antimalarial drug and has more recently been recognized as an anticancer agent. Some studies determine whether glaucarubinone, alone or in combination with the front-line chemotherapeutic agent gemcitabine. Glaucarubinone, a natural product first isolated from the seeds of the tree *Simarouba glauca*, was originally developed as an antimalarial drug and has more recently been recognized as an anticancer agent. Dichloromethane parts were screened for their cytotoxicities on brine shrimp larvae, and 50% inhibitory concentrations were resolute for *Plasmodium falciparum* in-vitro cultures. *Simarouba glauca* twigs collected and showed significant effect on tumor cell line.

References

- [1] Kaul S, Kumar A, Bhatnagar AK, Goyal HB and Gupta AK: Biodiesel: a clean and Sustainable fuel for the future. In: All India seminars on national policy on -non-edible oils as biofuels, II SC Bangalore 2003.
- [2] Rakshit D and Bhagya S: Journal of Science of Food and Agriculture 2008; 88: 911-919.
- [3] Neelavathi A, Sekhar KB, Babu CR and Jayaveera KN: Removal of toxic Cr (VI) by the adsorption of activated carbons prepared from *Simarouba* shells. J Environ SciEng 2004; 46(2): 137-42.
- [4] Joshi SG, Late SL, Joshi S and Hiremath S: Ramachandra Int J Biotech Biochem 2008; 4:Patil Manasi S, Gaikwad D K. A Critical Review on Medicinally Important Oil yielding Plant Laxmitaru (*Simarouba glauca* DC.). J. Pharm. Sci. & Res. 2011; 3(4): 1195-1213.
- [5] Rajurkar B M. A Comparative study on Antimicrobial activity of *Clerodendrum infortunatum*, *Simarouba glauca* and *Psoralea corylifolia*. International Journal of research and Review in pharmacy and Applied Sciences 2011; 1(4): 278-282.
- [6] Santhana Laxmi K, Sangeetha D, Sivamani S, Tamilarasan M, rajesh T P, Anandraj B. In Vitro Antibacterial, Antioxidant, haemolytic, Thrombolytic activities and phytochemical Analysis of *Simarouba glauca* leaves extracts. International Journal of Pharmaceutical Sciences and Research 2014; 5(2): 432-437.
- [7] Shankara Sharma, Sriram N. Anti-Ulcer activity of *Simarouba glauca* against Ethanol and Indomethacin induced ulcer in rats. Int. J. of Res. In Pharmacology & Pharmacotherapeutics 2014; 3(2): 85-89.
- [8] Sneha Kochath Santhosh, Aswathy venugopal, Meera Chakrappully Radhakrishnan. Study on the phytochemical, antibacterial and antioxidant activities of *Simarouba glauca*. South Indian Journal of Biological Sciences 2016; 2(1): 119-124.
- [9] Chase & Pratt R. Fluorescence of Powdered Vegetable drugs with particular reference to development of a system of identification, J. Am. Pharm. Assoc. Sci. Ed; 1949; 38: 324-331.
- [10] Rajurkar B M. A Comparative study on Antimicrobial activity of *Clerodendrum infortunatum*, *Simarouba glauca* and *Psoralea corylifolia*. International Journal of research and Review in pharmacy and Applied Sciences 2011; 1(4): 278-282.
- [11] Neelavathi, A., Chandra Sekhar, K. B., Ramesh Babu ,C., Jayaveera, K .N. J Environ Sci Eng., 2004, 46 :(2) 137-142
- [12] Joshi, S, Hiremath, S.R. (2000) *Simarouba* A potential oil tree seed. curr.sci.78 (6): 694-697. Severen, M. L., Journal of the American Oil Chemists Society.1953, 30:124-126.
- [13] Rao, K.V.S.A., Lakshminarayana, G., Journal of Food Science and Technology.1983.20:176-177
- [14] Krishnaiah, D., Devi, T., Bono, A., Sarbatly, R. 2009. Study on phytochemical constituents of six Malasian medicinal plants. J. Med. Plants Res., 3(2): 67-72.
- [15] Kumar, A., Tyagi, G., Sharma, S., Kumar,V., Pundir, R. 2014. Current Review on Biotechnological and Pharmacological Investigations of *Simarouba Glauca*-an oil Yielding Plant'. Int. J. Pharmacognosy, 1(12):100-122.
- [16] Manasi PS, Gaikwad DK. A critical review on medicinally important oil yielding plant laxmitaru (*Simarouba glauca* DC.). J Pharm Sci Res. 2011; 3(4):1195.
- [17] Kirby GC, O'Neill MJ, Phillipson JD, Warhurst DC. In vitro studies on the mode of action of quassinoids with activity against chloroquine-resistant *Plasmodium falciparum*. Biochem Pharmacol. 1989;38(24):4367-74.
- [18] Kaur K, Jain M, Kaur T, Jain R. Antimalarials from nature. Bioorg Med Chem. 2009;17(9):3229-56.
- [19] Lima RK, Guerreiro-Tanomaru JM, Faria-Júnior NB, Tanomaru-Filho M. Effectiveness of calcium hydroxide-based intracanal medicaments against *Enterococcus faecalis*. Int Endod J 2012 Apr;45(4):311-316.
- [20] Abida P., De Britto A.J., Antoney J., Stephan Raj T.L. Evaluation of in vitro anticancer activity of *Symplocos cochinchinensis* (Lour.) S. Moore bark. Int J Herb Med. 2016; 4:117-119.
- [21] Umesh T.G. In vitro antioxidant potential, free radical, scavenging and cytotoxic activity of *Simarouba glauca* leaves. Int J Pharm and Pharmaceutical Sci,7(2):411-416, 2015.
- [22] Renuka K., Vellai R.D, Sorimuthu P.S. Phytochemical screening and in vitro evaluation of antioxidant potential of immature *Palmyra* palm (*Borassus flabellifer* Linn) fruits. Int J Pharm Sci 10(8):77-83, 2018.
- [23] Jose, A., Kannan, E., Kumar, P. R. A. V., and Madhunapantula, S. V. (2019). Therapeutic Potential of Phytochemicals Isolated from *Simarouba glauca* for Inhibiting Cancers: A Review. Systematic Reviews in Pharmacy, 10(1); 73-80.
- [24] Joshi S, and Joshi S. (2002). Oil Tree *Laxmitaru Simarouba glauca*. University of Agricultural sciences, Bangalore and Indian council of Agricultural Research, New Delhi, India, ,86
- [25] Meshram, M. S., Itankar, P. R., and Patil, A. T. (2019). Journal of Pharmacognosy and Phytochemistry 8(3): 4224-4227.