

Cardiac Autonomic Neuropathy in Diabetes: A Predictor of Cardiometabolic Events: Overview

Premshanker Singh*, Ritu Karoli**, Shobhit Shakya

*FMR Prof of Medicine, **Addl Prof of Medicine, ***Assoc Prof of Medicine

Dr RML Institute of Medical Sciences, Lucknow-226010 **Corresponding Author:** Premshanker Singh mobile: +919412520054

Email: premshanker0354[at]gmail.com **Address:** N-244, Sector-N, Ashiana Colony, Lucknow-226003, U.P., India

1. Introduction

Autonomic nervous system (ANS) imbalance manifesting as cardiac autonomic neuropathy in the diabetic population is an important predictor of cardiovascular events. Symptoms and signs of ANS dysfunction, such as resting heart rate elevations, diminished blood pressure responses to standing, and altered time and frequency domain measures of heart rate variability in response to deep breathing, standing, and the Valsalva maneuver, should be elicited from all patients with diabetes and prediabetes. With the recognition of the presence of ANS imbalance or for its prevention, a rigorous regime should be implemented with lifestyle modification, physical activity, and cautious use of medications that lower blood glucose. Rather than intensifying diabetes control, a regimen tailored to the individual risk of autonomic imbalance should be implemented. New agents that may improve autonomic function, such as SGLT2 inhibitors, should be considered and the use of incretins monitored. One of the central mechanisms of dysfunction is disturbance of the hypothalamic cardiac clock, a consequence of dopamine deficiency that leads to sympathetic dominance, insulin resistance, and features of the metabolic syndrome. An improvement in ANS balance may be critical to reducing cardiovascular events, cardiac failure, and early mortality in the diabetic population.

As the population grows older and more obese, one third of people over the age of 65 will develop diabetes¹ and a host of comorbidities; and almost two thirds of these patients will experience cardiometabolic events. Unfortunately, the existence of comorbidities like metabolic syndrome, obesity, hyperlipidemia, and hypertension predict only about 24% of the risk of such events. Neuropathy (of all types) is one of the most prevailing complications of diabetes and a considerable source of morbidity and mortality; it includes DPN and AN². AN, particularly when it involves cardiac autonomic function (CAN), independently predicts risk of cardiovascular death and myocardial infarction (MI) resulting in substantial morbidity and mortality³.

It is likely that dysfunction of the ANS is an underdiagnosed cause of excessive morbidity and mortality in adults with diabetes. In fact, CAN results in cardiovascular (CV) dysfunction and is often accompanied by the progression to myocardial ischemia (which is often silent in diabetes), stroke, CAD, perioperative morbidity, and overall morbidity and mortality⁴. In combination with signs of DPN, the odds

ratio increases to 4.55 for mortality and CVD⁵. This is a more robust indicator of CVD risk, exceeding lipoprotein profile, BP, as well as adenosine scans⁶. Thus, CAN assessments may be useful to predict cardiovascular risk.

Cardiac Autonomic Neuropathy and Risk of Cardiovascular Events and Mortality

The prevalence of diabetes-related AN may be as high as 90% depending on the diagnostic methods and the cohort studied⁷. CAN prevalence increases substantially with diabetes duration in both T1DM (30% after 20 years)⁸ and T2DM (up to 60% after 15 years)⁸. In addition, CAN is present in some patients with prediabetes. CAN is strongly associated with risk of arrhythmias, major cardiovascular events, myocardial dysfunction, and cardiovascular mortality^{9, 10}. Consequently, early diagnosis of cardiac autonomic dysfunction is important. Several studies including ADVANCE, VADT, and ACCORD) have shown that cardiac autonomic dysfunction may predict the risk of cardiovascular events and sudden death seen with intensification of glycemic control in subjects with T2DM. Significant escalations in cardiovascular events and all causes of mortality were independently correlated with resting baseline and average HR in a *post hoc* analysis of the ONTARGET/TRANSCEND studies, two large cohort studies of patients with stable, chronic CVD^{10, 11}. The influence of elevated resting HR and blunted HRV, two measures of ANS imbalance, on the development and progression of CVD, diabetes, and early mortality was assessed¹³. In this Framingham Heart Study offspring cohort 2 these measures, in addition to smoking, age, and gender, were shown to be significant predictors for developing CVD, DM, and early mortality within 12 years¹⁴ also showed that resting tachycardia (HR > 100 beats/minute), chronotropic incompetence (inability of the heart to increase its rate commensurate with increased activity or demand), and reduced HR recovery after treadmill exercise were independently correlated with stroke, MI, and overall mortality in T2DM patients (hazard ratios of 1.97, 1.89, and 1.77, respectively), regardless of established coronary heart disease. In addition, the EURODIAB Prospective Cohort Study revealed that CAN was the most distinctive predictor for mortality in a large group of patients with T1DM, and increased mortality risk with deteriorating CAN was shown in a meta-analysis of several trials^{14, 15}.

Detection of Cardiac Autonomic Neuropathy

The most frequent clinical symptoms of CAN include dizziness, heart palpitations, lightheadedness, and

fragility, all of which are consequences of dysregulation of the cardiovascular system secondary to malfunction of the sympathetic and parasympathetic nervous systems. Early autonomic dysfunction may exhibit no symptoms and only be detected by abnormal indices of HRV. Resting tachycardia (>100 bpm) and/or a fixed HR, as well as, orthostatic hypotension, intolerance to exercise, and syncope are present in patients with advanced CAN^{16, 17}. In advanced stages diagnosis can be done clinically, but HRV tests may be necessary to detect early asymptomatic cardiac autonomic dysfunction. This can be easily performed in the office by utilizing an electrocardiogram recording during either 1–2 min of deep breathing, or as the subject begins to rise from a seated position, with calculation of HRV indices^{18, 19}.

ADA recommendations regarding screening and diagnosis of CAN are as follows²⁰

- “Symptoms and signs of AN should be assessed in patients with microvascular and neuropathic complications.
- In the presence of symptoms or signs of cardiovascular AN, tests excluding other comorbidities or drug effects/interactions that could mimic cardiovascular AN should be performed.
- Consider assessing symptoms and signs of cardiovascular AN in patients with hypoglycemia unawareness.”

The Toronto Consensus Panel, the European Society of Cardiology, the North American Society of Pacing and Electrophysiology, and the ADA Position Statement on Diabetic Neuropathy recommend the following regarding CAN assessments for clinical trials measuring a targeted intervention or for prognostication^{21, 22}

- “Standardized CARTs: simple, sensitive, specific, and reproducible tests that assess changes in the R–R interval on electrocardiogram recordings in response to simple clinical maneuvers (deep breathing, Valsalva, and standing);
- Indices of HRV including sdNN and rMSSD;
- Resting HR and measurement of QTc interval on ECG recording”.

Cardiovascular autonomic reflex tests assess cardiovascular autonomic function through time-domain HR response to deep breathing, Valsalva maneuver and postural change, and by measuring the consequent changes in HR and BP. Although indirect autonomic measures, the following CARTs are considered the gold standard in autonomic testing: HR response to deep breathing (E: I ratio), standing (30: 15 ratio) and Valsalva maneuver, and BP response to standing. The presence of one abnormal cardiovagal test result identifies possible or early CAN, to be confirmed over time. At least two abnormal cardiovagal results are required for a definite or confirmed diagnosis of CAN. The time-domain HR tests and the BP response to postural change have the reproducibility necessary for clinical trials.

Time-Domain Measures of Heart Rate Patterns

Analysis of time-domain measures under resting conditions offers an accurate assessment of the sympathetic and parasympathetic regulation of the heart beat (the R–R interval on an electrocardiogram) documented at baseline conditions and during deep breathing, Valsalva, and standing from a sitting position maneuvers. The sdNN is an evaluation of both sympathetic and parasympathetic activity on HRV, and the rMSSD is a primary indicator of parasympathetic activity. Expiratory to inspiratory ratio (E/I ratio) measures HRV during deep breathing, which is parasympathetic predominant. Postural 30: 15 ratio is evaluated at beats 15 and 30 after standing up and is regarded as reflective of sympathetic response and baroreflex function demonstrated that cardiac parasympathetic withdrawal mediated by the carotid cardiac baroreflex is the principal trigger for tachycardia within milliseconds of a postural change, while sympathetic adrenergic control sustains tachycardia during extended periods of orthostasis. The Valsalva ratio is calculated by the longest R–R interval during the procedure to the shortest R–R interval throughout the duration or immediately following the maneuver. This reaction is facilitated by the interspersed activity of parasympathetic and sympathetic nerve fibers^{21, 22, 23}.

Frequency-Domain Measures

Frequency-domain analysis can identify underlying periodicities in HR patterns. Rfa (also termed HF power) is calculated as the area under the HR spectral curve over a frequency range fixed on the fundamental Rfa (0.15–0.4 Hz), which is defined by the peak mode of the respiratory activity spectrum. LF is computed as the area under the HR spectral curve over the frequency range from 0.04 to 0.10 Hz^{23, 24}.

Despite its use in research for decades, the concept that LF and HF bands fully reflect separate influences of the sympathetic and parasympathetic branches has been recently under debate mainly due to their simultaneous action in the LF power^{24, 25, 26}. Therefore, there has been increasing interest in using non-linear analyses of HRV, as these may be clinically more relevant by providing a better interpretation of the pathophysiological behavior of HRV under various conditions and by enhancing its prognostic value^{26, 27}. Non-linear analysis methods do not assess the magnitude of variability but rather the quality, scaling, and correlation properties of the signals; these analyses allow a more subtle characterization of autonomic balance and have been shown to be more reliable markers of morbidity and mortality in patients with CVD. A number of studies have shown that abnormal non-linear HRV indices are associated with diabetes or an elevated risk of developing diabetes^{27, 28} that non-linear HRV indices in diabetic populations may have diagnostic and prognostic potential for identifying asymptomatic CAN and cardiovascular events^{28, 29}. The technical complexity of these analyses, however, has made interpretation and understanding of variability challenging for common clinical use. Further research is

needed to demonstrate conclusively that these refinements in the analysis enhance the sensitivity for prediction of cardiovascular events^{28, 29}

Sudorimetry Measures

Sudomotor nerves are thin unmyelinated C-fibers, with largely cholinergic neurotransmission, where the ganglion neurotransmitter is acetylcholine, the primary parasympathetic nervous system neurotransmitter. However, epinephrine, norepinephrine, vasoactive intestinal peptide (VIP), atrial natriuretic peptide, calcitonin gene related polypeptide (CGRP), galanin, ATP, and substance P have been identified in periglandular nerves and thus may be contributing to the electrical response. The addition of sudomotor function assessments, combined with CART, may present a more precise and well-defined early diagnosis of ANS dysfunction.

Although quantitation of intraepidermal nerve fiber density on skin biopsies remains the gold standard and is the most recognized technique to diagnose small nerve fiber dysfunction, sudorimetry has the ability to produce diagnostic information on the evaluation of the small somatosensory nerves, detection and progression of disease, and responsiveness to therapeutic intervention. Sudorimetry technology has advanced rapidly as a non-invasive and precise tool to assess small fibers that can potentially be incorporated into clinical practice.

Current sudorimetry assessments can be performed using Sudoscan™, which measures electrochemical skin conductance (ESC) of hands and feet. This technology is founded on the electrochemical theories of reverse iontophoresis and chronoamperometry to measure sudomotor function, which makes it an affordable, practical, and precise tool generating accurate profiles for routine clinical use and a viable research tool on the integrity of this complex system of control. This testing has assumed greater significance now that there are medications that can prevent the development or progression of ANS dysfunction. The American Association of Clinical Endocrinologists (AACE) endorses the use of current procedural technology (CPT) code 95923 for simplified sudomotor testing and the code 95943 for evaluation of cardiac autonomic function testing. AACE would urge that sudomotor function testing be authorized for all practitioners seeing patients with diabetes, including primary care, endocrinology, and podiatry. It is a non-invasive objective test, takes a mere 2 min, and it has been compared with other reference tests [including HRV indices, intraepidermal nerve fiber density, and quantitative sudomotor axon reflex testing (QSART)] and been demonstrated to be useful in the detection of small nerve fiber neuropathy in those with and without T2DM with a sensitivity of 77 to 87% and a specificity of 67 to 92% (AUC 0.77 to 0.88)^{29, 30}

Medical Treatment of Cardiac Autonomic Neuropathy

Cardiac autonomic neuropathy therapies are typically focused on mitigating symptoms and should be directed to

specified clinical manifestations. Exercise, volume repletion, low dose fludrocortisone and midodrine are among the most frequently used therapies. Recommendations for the treatment of CAN include the following^{30, 31}

- Early optimization of blood glucose regulation to avoid or delay the development of CAN in people with T1DM;
- Multifaceted approach to treat hyperglycemia and additional risk factors (e.g., dyslipidemia, hypertension) to prevent CAN in T2DM;
- Lifestyle modifications to improve CAN in patients with prediabetes.

Most recently there has been great interest in the action and effects of the sodium-glucose cotransporter-2 (SGLT2) inhibitors on reducing cardiovascular events. Empagliflozin is a highly selective inhibitor of the SGLT2 in the kidney. Glucose reduction occurs by decreasing renal glucose reabsorption and thereby increasing urinary glucose elimination in patients with diabetes, leading to significant reductions in glycated hemoglobin (HbA1c), weight loss, and reductions in BP without increases in HR^{31, 32}.

The EMPA-REG Outcome trial recruited 1, 000 patients with T2DM, 700 of whom were enrolled and randomized to placebo or one of two different doses of empagliflozin (10 and 25 mg) daily in addition to standard care³³. Empagliflozin was similar to other oral antihyperglycemic agents in HbA1c reduction (0.6–1%) and decreased both fasting and postprandial glucose, with a modest weight loss of ~3 kg at 26 weeks vs. placebo, slightly greater weight loss at 52 weeks, modest BP reduction of 2–7 mmHg vs. placebo, and no intrinsic increased risk of hypoglycemia. The initial report demonstrated a significant reduction in cardiovascular events (predominantly mortality and admission to hospital for heart failure) by 35% which fast tracked the drug to be approved for both diabetes and CVD. A subsequent report showed that the reduction in cardiovascular deaths were significant in Southeast Asia and Latin America, but not as much in America and Europe³⁴. Despite these different findings, the fall in BP without an increase in HR implies a reduction in sympathetic tone with its use.

Liraglutide, a GLP-1 receptor agonist, was also found to reduce CV events, but not as robustly as empagliflozin³⁵. GLP-1 has widespread properties in the human body and targets receptors diffusely³⁶. Liraglutide improves HbA1c and compared with other medication classes has similar or greater efficacy, even compared to basal insulin. Its use has been shown to lead to a modest improvement in BP but, in contrast to empagliflozin, with an increase in HR³⁷. The FDA recently approved the use of liraglutide for management of CVD in diabetes³⁸. The actions of liraglutide on HRV and daily variation of HR in newly diagnosed, overweight patients with T2DM and stable CAD have been investigated. Diurnal HR fluctuations and sympathovagal balance evaluated by rMSSD in NN intervals and HF and LF power were assessed. Liraglutide decreased sdNN in some subjects; decreased rMSSD; and

increased mean, daytime, and nighttime HR compared to placebo. Liraglutide reduced HF power without any change in LF/HF ratio. Thus, in overweight patients with CAD and newly diagnosed T2DM, liraglutide increased HR and reduced HRV despite significant weight loss and improvement in metabolic parameters; the increase in nightly HR and decrease in parameters of parasympathetic activity (rMSSD and HF power) suggest that this medication may negatively affect sympathovagal balance.³⁹The authors hypothesize that the chronotropic effect of liraglutide, which may be mediated through the GLP-1 receptor on the sinoatrial node, cannot explain the worsening of HRV measures; instead, the impaired HRV may be due to a direct influence on sympathovagal balance, as reflected by the increase in night-time HR in conjunction with the significant decrease in sdNN and rMSSD suggesting an impairment of parasympathetic activity. The addition of a cholinergic agent to a GLP-1 analog might recapture the loss of cholinergic activity induced by a GLP-1 analog. This might even be a useful strategy to further enhance the cardiac protection afforded by the SGLT-2 inhibitors.

Non-Medical Treatment of Cardiac Autonomic Neuropathy

A number of researchers have demonstrated that autonomic balance can be restored using simple lifestyle interventions, potentially reversing CAN.⁴⁰ showed that elderly women experienced improved HRV while walking a dog by enhancing parasympathetic function. Removing the dog resulted in reversal of this benefit with sympathetic overactivity.⁴¹ There is strong evidence indicating that individuals with greater aerobic capacity exhibit enhanced HRV.⁴¹ Furthermore several studies have shown significant improvements in HRV measures after different training programs including cycling, walking, jogging and water aerobic exercise training in subjects with CAD. inspiratory muscle training in elderly individuals and high-intensity interval training in young, healthy individuals We have documented that falls and fractures in older diabetics were often the result of loss of organized^{42,43,44} variability, strength, and reaction times. Very simple strength and balance training can significantly reduce falls risk.⁴⁵For patients with orthostatic hypotension, volume repletion with both fluids and salt is central to management, but physical activity and exercise are essential to prevent deconditioning, which is known to exacerbate orthostatic intolerance.⁴⁶The relationship between HRV and different psychiatric disorders, as well as stress and trauma, has also been extensively studied. Subjects with depression and anxiety disorders exhibit abnormal HRV patterns compared with non-psychiatric controls. Reduced HRV characterizes emotional dysregulation, decreased psychological flexibility and defective social engagement, which in turn are linked to prefrontal cortex hypoactivity^{47, 48, 49}. High occupational stress has also been associated with lowered HRV, specifically with reduced parasympathetic activation. There is limited evidence that use of biofeedback with relaxation and meditation approaches may result in increased HRV and parasympathetic activity⁵⁰. A more detailed review on this topic is beyond the scope of this

article and the reader can refer to recent reviews on the subject.

Prevention and Reversibility of Cardiac Autonomic Neuropathy

Prevention of CAN should be a primary focus of lifestyle and other clinical interventions. Intense glycemic control⁵¹ utilizing a step-by-step progressive lowering of hyperglycemia, lipids, and BP, in addition to the use of antioxidants⁵²and ACE inhibitors reduces the odds ratio for CAN^{53, 54}. In the EDIC study, patients with T1DM showed continuously favorable effects of past glucose control on microvascular complications in spite of the loss of glycemic separation^{54, 55}. CAN progressed in both treatment groups during the EDIC follow-up, but the prevalence and incidence continued to be decreased in the previous intensive group compared to the standard group despite comparable levels of glycemic control. To diminish the development of CAN, intense glucose control of T1DM ought to be started as soon as possible⁵⁶. However, in patients with established CAN, glycemic control may need to be less stringent to avoid hypoglycemia and adverse drug effects⁵⁷The American Diabetes Association also recommends that individuals with CAN have a cardiac evaluation before starting or increasing physical activity for safety reasons⁵⁸ Pathogenesis-oriented interventions may promote some degree of reversal of established CAN⁵⁹Lifestyle interventions, increased physical activity, β -adrenergic blockers, aldose reductase inhibitors, ACE inhibitors, ARBs, and potent antioxidants such as α -lipoic acid have all been shown to restore autonomic balance. Enhanced glycemic control with a reduced HbA1c from 9.5 to 8.4 improved HRV in patients with T1DM with minimal autonomic abnormalities, but not in those with advanced autonomic irregularities⁶⁰The Veterans Administration Cooperative Study showed no impact on the occurrence of CAN after 2 years of intense glycemic control in patients with T2DM; however, people with T2DM receiving rigorous multifactorial therapies that targeted hypertension, hyperglycemia, dyslipidemia, and microalbuminuria, along with secondary CVD preventative measures like aspirin use, experienced a 63% reduction in ANS dysfunction in the Steno-2 Study. Although glucose-lowering agents exerted the least benefit in comparison with antihypertensive treatments, lipid-lowering agents, aspirin, and vitamin-mineral supplements^{61, 62}, this favorable effect continued after ~13 years of follow-up⁶³. Early identification of CAN also may allow for the well-timed initiation of antioxidant α -lipoic acid therapies that slow or reverse advancement of CAN⁶⁵.

Certain medications hold promise for the prevention and reversal of CAN. Early therapeutic intervention with ACE inhibition or ARBs improved both CAN and left ventricular diastolic dysfunction after 1 year of treatment in patients with no symptoms and long-term diabetes. The combined therapies were slightly superior to monotherapies, auguring well for patients with established CAN⁶⁶. Treatment with fluvastatin improves cardiac sympathetic neuropathy in the diabetic rat heart in relation

to attenuation of increased cardiac oxidative stress⁶⁷. Alternatively, selective inactivation of cyclooxygenase-2 (COX-2) guards against sympathetic denervation in experimental diabetes by decreasing intramyocardial oxidative stress and inflammation⁶⁸. Consequently, statins and COX-2 inactivation may assist in attenuating cardiac sympathetic dysfunction. In addition, it has been shown that early mortality is a consequence of beat-to-beat variability loss with MI, but can be reduced by 33% with the immediate administration of insulin. Successful pancreas transplantation showed improvements in epinephrine response and normalized hypoglycemia symptom awareness in patients with established diabetes as well as evidence of a reversible metabolic component in patients with early CAN^{69, 70, 71}.

Weight loss and weight-reducing surgeries may also potentially reduce CAN. ANS dysfunction and increased sympathetic activity have been directly correlated with obesity⁷². Moreover, weight reduction significantly improves HRV and reduces ANS imbalances^{71, 72, 73}. To evaluate the ability to reverse autonomic imbalance, we examined sudomotor function and HRV measurements in obese patients undergoing bariatric surgery. Patients were assessed at baseline, 4, 12, and 24 weeks after vertical sleeve gastrectomy or Roux-en-Y gastric bypass. Seventy subjects completed at least 24-weeks of follow-up. Sudorimetry results of ESC of feet improved significantly trending toward normal in T2DM patients. HRV improved significantly, as did many other metabolic parameters. Improvements in feet ESC were shown to be independently associated with HbA1c, insulin, and HOMA2-IR levels at baseline, as well as HbA1c at 24 weeks. Additionally, improvement in basal HR had an independent association with HbA1C, insulin and HOMA2-IR levels. These positive results suggest that bariatric surgery can return both cardiac and sudomotor autonomic C-fiber dysfunction in those with diabetes to normal, possibly positively influencing morbidity and mortality⁷⁴.

Future Directions of Research to Prevent and Reverse Cardiac Autonomic Neuropathy

The host of targets that are potential candidates for reduction of cardiovascular risk have been addressed in the previous paragraphs. For years we were confronted with glycemic control as the only measure by the glucocentric majority and those who believed in the lipid hypothesis who have now carried this to the extreme of need for even lower LDL-C in high risk patients. The entry of SGLT2 inhibitors and the incretins shed new light on the challenge armed with new ammunition and also created an avenue of adventure for those interested in novel pathways. However the initial inroad into reduction of CV events was a discovery of the power of resetting a biologic clock and targeting the brain rather than other members of the dreadful dektet!

It has been established that there is a brain dopamine deficiency in obese diabetic patients present in the early hours of the morning⁷⁵. The working hypothesis is that in early morning, decreased dopaminergic tone in the

hypothalamus unbridles sympathetic activation with all its consequences, as illustrated in Figure 1. Restoring the morning peak in dopaminergic activity by dopamine D2 receptor-mediated activities may, therefore, restore ANS balance.

Figure 1

Enhanced activity of the paraventricular nucleus (PVN), which increase autonomic tone and the paraventricular nucleus to raise levels of corticotrophin releasing hormone (CRH). The consequences include activation of the sympathetic nervous system (SNS) hypothalamic and glucose sensitization reducing parasympathetic (vagal drive) to the liver and resistance to both leptin and insulin. CRH, corticotrophin releasing hormone; eNOS, endothelial nitric oxide synthase; FFAs, free fatty acids; NPY, neuropeptide Y; PVN, paraventricular nucleus; SCN, suprachiasmatic nucleus; SNS, sympathetic nervous system; TGs, triglycerides; VMH, ventromedial hypothalamus⁷⁶

It may also be possible to reset the biologic hypothalamic clock and ANS function using bromocriptine QR to restore morning dopaminergic activity. It sensitizes the body to insulin and reduces sympathetic tone thereby reducing HR⁷⁷. Bromocriptine QR has also demonstrated a favorable effect on CV outcomes in clinical trials, suggesting a future direction for pathogenesis-oriented therapies^{78, 79, 80}. Our current quest is to determine if any of the novel discoveries in cardiovascular outcome studies (CVOTs) are indeed working through rebalancing the ANS thereby creating a wonderful opportunity for taking a fork in the road^{81, 82}.

2. Conclusion

An improvement in ANS balance may be critical to reducing cardiovascular events and early mortality. Symptoms and signs of autonomic dysfunction, including resting HR, BP responses to standing, and time and frequency measures of HRV in response to deep breathing, standing and Valsalva maneuver, should be elicited from all patients with diabetes to allow for early detection and intervention. With the recognition of the presence of ANS imbalance or for its prevention, a rigorous regime should be implemented with lifestyle modification, physical activity, and cautious use of medications that lower blood glucose. Rather than intensifying diabetes blood glucose management, a regimen tailored to the individual risk of ANS dysfunction should be constructed. The advent of new agents that may have the potential to improve ANS function, such as the SGLT2 inhibitors and the GLP-1 agonists, should be considered. However, it is not clear how these compounds work and what the mechanism of reduction of major adverse cardiovascular events is. An overlooked mechanism is a resetting of the biologic clock with correction of the dopamine deficiencies in the brainstem of obese people with diabetes, restoring the functioning of the ANS with its potential for significant reduction of cardiovascular events.

Conflict of Interest Statement: None

Funding: None

Ethical Clearance: Taken by Ethical Committee

References

- [1] American Diabetes Association (2017). Standards of medical care in diabetes - 2017. *Diabetes Care* 40(Suppl. 1), S1–S135.
- [2] Athyros, V. G., Didangelos, T. P., Karamitsos, D. T., Papageorgiou, A. A., Boudoulas, H., and Kontopoulos, A. G. (1998). Long-term effect of converting enzyme inhibition on circadian sympathetic and parasympathetic modulation in patients with diabetic autonomic neuropathy. *Acta Cardiol.* 53, 201
- [3] Azad, N., Emanuele, N. V., Abaira, C., Henderson, W. G., Colwell, J., Levin, S. R., et al. (1999). The effects of intensive glycemic control on neuropathy in the VA cooperative study on type II diabetes mellitus (VA CSDM). *J. Diabetes Complications* 13, 307–313. doi: 10.1016/S1056-8727(99)00062-8
- [4] Bernardi, L., Spallone, V., Stevens, M., Hilsted, J., Frontoni, S., Pop-Busui, R., et al. (2011). Methods of investigation for cardiac autonomic dysfunction in human research studies. *Diabetes Metab. Res. Rev.* 27, 654–664. doi: 10.1002/dmrr.1224
- [5] Boyle, J. P., Thompson, T. J., Gregg, E. W., Barker, L. E., and Williamson, D. F. (2010). Projection of the year 2050 burden of diabetes in the US adult population: dynamic modeling of incidence, mortality, and prediabetes prevalence. *Popul. Health Metr.* 8:29. doi: 10.1186/1478-7954-8-29
- [6] Brownlee, M., Aiello, L. P., Cooper, M. E., Vinik, A. I., Plutzky, J., and Boulton, A. J. M. (2016). “Complications of diabetes mellitus,” in *Williams Textbook of Endocrinology*, eds S. Melmed, K. S. Polonsky, P. R. Larsen, and H. M. Kronenberg (Philadelphia, PA: Elsevier), 1484–1581.
- [7] Burger, A. J., Weinrauch, L. A., D’Elia, J. A., and Aronson, D. (1999). Effects of glycemic control on heart rate variability in type I diabetic patients with cardiac autonomic neuropathy. *Am. J. Cardiol.* 84, 687–691. doi: 10.1016/S0002-9149(99)00417-8
- [8] Carnethon, M. R., Prineas, R. J., Temprosa, M., Zhang, Z. M., Uwaifo, G., and Molitch, M. E. (2006). The association among autonomic nervous system function, incident diabetes, and intervention arm in the Diabetes Prevention Program. *Diabetes Care* 29, 914–919. doi: 10.2337/diacare.29.04.06.dc05-1729
- [9] Casellini, C. M., Parson, H. K., Hodges, K., Edwards, J. F., Lieb, D. C., Wohlgemuth, S. D., et al. (2016). Bariatric surgery restores cardiac and sudomotor autonomic C-fiber dysfunction towards normal in obese subjects with type 2 diabetes. *PLoS One* 11:e0154211. doi: 10.1371/journal.pone.0154211
- [10] Casellini, C. M., Parson, H. K., Hodges, K., et al. (2015). “Abstract: bariatric surgery restores somatic and autonomic nerve function towards normal in obese patients with and without diabetes,” in *Proceedings of the AACE 24th Annual Scientific Clinical Congress*, Nashville, TN.
- [11] Casellini, C. M., Parson, H. K., Richardson, M. S., Nevoret, M. L., and Vinik, A. I. (2013). Sudoscan, a noninvasive tool for detecting diabetic small fiber neuropathy and autonomic dysfunction. *Diabetes Technol. Ther.* 15, 948–953. doi: 10.1089/dia.2013.0129
- [12] Cincotta, A. H., Meier, A. H., and Cincotta, J. M. (1999). Bromocriptine improves glycaemic control and serum lipid profile in obese Type 2 diabetic subjects: a new approach in the treatment of diabetes. *Expert Opin. Investig Drugs* 8, 1683–1707. doi: 10.1517/13543784.8.10.1683
- [13] Convertino, V. A. (2014). Neurohumoral mechanisms associated with orthostasis: reaffirmation of the significant contribution of the heart rate response. *Front. Physiol.* 5:236. doi: 10.3389/fphys.2014.00236
- [14] Cowie, C. C., Rust, K. F., Ford, E. S., Eberhardt, M. S., Byrd-Holt, D. D., Li, C., et al. (2009). Full accounting of diabetes and pre-diabetes in the U.S. population in 1988-1994 and 2005-2006. *Diabetes Care* 32, 287–294. doi: 10.2337/dc08-1296
- [15] De Godoy, M. (2016). Nonlinear analysis of heart rate variability: a comprehensive review. *J. Cardiol. Ther.* 3, 528–533. doi: 10.17554/j.issn.2309-6861.2016.03.101-4
- [16] De Sousa, A. F. M., Medeiros, A. R., Benitez-Flores, S., Del, R. S., Stults-Kolehmainen, M., and Boulosa, D. A. (2018). Improvements in attention and cardiac autonomic modulation after a 2-weeks sprint interval training program: a fidelity approach. *Front. Physiol.* 9:241. doi: 10.3389/fphys.2018.00241
- [17] Diabetes Control and Complications Trial Research Group (1993). The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin dependent diabetes mellitus. *N. Engl. J. Med.* 329, 977–986
- [18] Didangelos, T. P., Arsos, G. A., Karamitsos, D. T., Athyros, V. G., Georga, S. D., and Karatzas, N. D. (2006). Effect of quinapril or losartan alone and in combination on left ventricular systolic and diastolic functions in asymptomatic patients with diabetic autonomic neuropathy. *J. Diabetes Complications* 20, 1–7. doi: 10.1016/j.jdiacomp.2005.05.002
- [19] Drucker, D. J. (2016). Never waste a good crisis: confronting reproducibility in translational research. *Cell Metab.* 24, 348–360. doi: 10.1016/j.cmet.2016.08.006
- [20] Duckworth, W., Abaira, C., Moritz, T., Reda, D., Emanuele, N., Reaven, P. D., et al. (2009). Glucose control and vascular complications in veterans with type 2 diabetes. *N. Engl. J. Med.* 360, 129–139. doi: 10.1056/NEJMoa0808431
- [21] Fathizadeh, P., Shoemaker, W. C., Wo, C. C., and Colombo, J. (2004). Autonomic activity in trauma patients based on variability of heart rate and respiratory rate. *Crit. Care Med.* 32, 1300–1305. doi: 10.1097/01.CCM.0000127776.78490.E4
- [22] Freedman, B. I., Smith, S. C., Bagwell, B. M., Xu, J., Bowden, D. W., and Divers, J. (2015). Electrochemical skin conductance in diabetic kidney

- disease. *Am. J. Nephrol.* 41, 438–447. doi: 10.1159/000437342
- [23] Gaede, P., Lund-Andersen, H., Parving, H. H., and Pedersen, O. (2008). Effect of a multifactorial intervention on mortality in type 2 diabetes. *N. Engl. J. Med.* 358, 580–591. doi: 10.1056/NEJMoa0706245
- [24] Gaede, P., Vedel, P., Parving, H. H., and Pedersen, O. (1999). Intensified multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: the Steno type 2 randomized study. *Lancet* 353, 617–622. doi: 10.1016/S0140-6736(98)07368-1
- [25] Gaziano, J. M., Cincotta, A. H., Vinik, A., Blonde, L., Bohannon, N., and Scranton, R. (2012). Effect of bromocriptine-QR (a quick-release formulation of bromocriptine mesylate) on major adverse cardiovascular events in type 2 diabetes subjects. *J. Am. Heart Assoc* 1:e002279. doi:10.1161/JAHA.112.002279
- [26] Green, J. B., Bethel, M. A., Armstrong, P. W., Buse, J. B., Engel, S. S., Garg, J., et al. (2015). Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N. Engl. J. Med.* 373, 232–242. doi: 10.1056/NEJMoa1501352
- [27] Heathers, J., and Goodwin, M. (2017). *Dead Science in Live Psychology: A Case Study from Heart Rate Variability (HRV)*. doi: 10.31234/osf.io/637ym
- [28] Inzucchi, S. E., Bergenstal, R. M., Buse, J. B., Diamant, M., Ferrannini, E., Nauck, M., et al. (2015). Management of hyperglycemia in type 2 diabetes, 2015: a patient-centered approach: update to a position statement of the American diabetes association and the European association for the study of diabetes. *Diabetes Care* 38, 140–149. doi: 10.2337/dc14-2441
- [29] Karason, K., Molgaard, H., Wikstrand, J., and Sjoström, L. (1999). Heart rate variability in obesity and the effect of weight loss. *Am. J. Cardiol.* 83, 1242–1247. doi: 10.1016/S0002-9149(99)00066
- [30] Kellogg, A. P., Converso, K., Wiggin, T., Stevens, M., and Pop-Busui, R. (2009). Effects of cyclooxygenase-2 gene inactivation on cardiac autonomic and left ventricular function in experimental diabetes. *Am. J. Physiol. Heart Circ. Physiol.* 296, H453–H461. doi: 10.1152/ajpheart.00678.2008
- [31] Kendall, D. M., Rooney, D. P., Smets, Y. F., Salazar Bolding, L., and Robertson, R. P. (1997). Pancreas transplantation restores epinephrine response and symptom recognition during hypoglycemia in patients with long-standing type I diabetes and autonomic neuropathy. *Diabetes* 46, 249–257. doi: 10.2337/diab.46.2.249
- [32] Khandoker, A. H., Jelinek, H. F., and Palaniswami, M. (2009). Identifying diabetic patients with cardiac autonomic neuropathy by heart rate complexity analysis. *Biomed. Eng. Online* 8:3. doi: 10.1186/1475-925X-8-3
- [33] Kishi, T. (2012). Heart failure as an autonomic nervous system dysfunction. *J. Cardiol.* 59, 117–122. doi: 10.1016/j.jcc.2011.12.006
- [34] Kumarathurai, P., Anholm, C., Larsen, B. S., Olsen, R. H., Madsbad, S., Kristiansen, O., et al. (2017). Effects of liraglutide on heart rate and heart rate variability: a randomized, double-blind, placebo-controlled crossover study. *Diabetes Care* 40, 117–124. doi: 10.2337/dc16-1580
- [35] Kumarathurai, P., Anholm, C., Nielsen, O. W., Kristiansen, O. P., Mølvi, J., Madsbad, S., et al. (2016). Effects of the glucagon-like peptide-1 receptor agonist liraglutide on systolic function in patients with coronary artery disease and type 2 diabetes: a randomized double-blind placebo-controlled crossover study. *Cardiovasc. Diabetol.* 15:105. doi: 10.1186/s12933-016-0425-2
- [36] Laing, S. T., Gluckman, T. J., Weinberg, K. M., Lahiri, M. K., Ng, J., and Goldberger, J. J. (2011). Autonomic effects of exercise-based cardiac rehabilitation. *J. Cardiopulm. Rehabil. Prev.* 31, 87–91. doi: 10.1097/HCR.0b013e3181f1fda0
- [37] Lambert, E., Sari, C. I., Dawood, T., Nguyen, J., McGrane, M., Eikelis, N., et al. (2010). Sympathetic nervous system activity is associated with obesity-induced subclinical organ damage in young adults. *Hypertension* 56, 351–358. doi: 10.1161/HYPERTENSIONAHA.110.155663
- [38] Lambert, G. W., Straznicki, N. E., Lambert, E. A., Dixon, J. B., and Schlaich, M. P. (2010). Sympathetic nervous activation in obesity and the metabolic syndrome—causes, consequences and therapeutic implications. *Pharmacol. Ther.* 126, 159–172. doi: 10.1016/j.pharmthera.2010.02.002
- [39] Liakos, A., Karagiannis, T., Athanasiadou, E., Sarigianni, M., Mainou, M., Papatheodorou, K., et al. (2014). Efficacy and safety of empagliflozin for type 2 diabetes: a systematic review and meta-analysis. *Diabetes Obes. Metab.* 16, 984–993. doi: 10.1111/dom.12307
- [40] Lonn, E. M., Rambihar, S., Gao, P., Custodis, F. F., Sliwa, K., Teo, K. K., et al. (2014). Heart rate is associated with increased risk of major cardiovascular events, cardiovascular and all-cause death in patients with stable chronic cardiovascular disease: an analysis of ONTARGET/TRANSCEND. *Clin. Res. Cardiol.* 103, 149–159. doi: 10.1007/s00392-013-0644-4
- [41] Low, P. A. (1996). Diabetic autonomic neuropathy. *Semin. Neurol.* 16, 143–151. doi: 10.1055/s-2008-1040970
- [42] Low, P. A., Benrud-Larson, L. M., Sletten, D. M., Opfer-Gehrking, T. L., Weigand, S. D., O'Brien, P. C., et al. (2004). Autonomic symptoms and diabetic neuropathy: a population-based study. *Diabetes Care* 27, 2942–2947. doi: 10.2337/diacare.27.12.2942
- [43] Luque-Casado, A., Zabala, M., Morales, E., Mateo-March, M., and Sanabria, D. (2013). Cognitive performance and heart rate variability: the influence of fitness level. *PLoS One* 8:e56935. doi: 10.1371/journal.pone.0056935
- [44] Lykke, J. A., Tarnow, L., Parving, H. H., and Hilsted, J. (2008). A combined abnormality in heart rate variation and QT corrected interval is a strong predictor of cardiovascular death in type 1 diabetes. *Scand. J. Clin. Lab. Invest.* 68, 654–659. doi: 10.1080/00365510802018330

- [45] Malmberg, K., Norhammar, A., Wedel, H., and Ryden, L. (1999). Glycometabolic state at admission: important risk marker of mortality in conventionally treated patients with diabetes mellitus and acute myocardial infarction: long-term results from the Diabetes and Insulin-Glucose Infusion in Acute Myocardial Infarction (DIGAMI) study. *Circulation* 99, 2626–2632. doi: 10.1161/01.CIR.99.20.2626
- [46] Martin, C. L., Albers, J. W., and Pop-Busui, R. (2014). Neuropathy and related findings in the diabetes control and complications trial/epidemiology of diabetes interventions and complications study. *Diabetes Care* 37, 31–38. doi: 10.2337/dc13-2114
- [47] Maser, R. E., Lenhard, M. J., Irgau, I., and Wynn, G. M. (2007). Impact of surgically induced weight loss on cardiovascular autonomic function: one-year follow-up. *Obesity* 15, 364–369. doi: 10.1038/oby.2007.554
- [48] Maser, R. E., Mitchell, B. D., Vinik, A. I., and Freeman, R. (2003). The association between cardiovascular autonomic neuropathy and mortality in individuals with diabetes: a meta-analysis. *Diabetes Care* 26, 1895–1901. doi: 10.2337/diacare.26.6.1895
- [49] Matsuki, A., Nozawa, T., Igarashi, N., Sobajima, M., Ohori, T., Suzuki, T., et al. (2010). Fluvastatin attenuates diabetes-induced cardiac sympathetic neuropathy in association with a decrease in oxidative stress. *Circ. J.* 74, 468–475. doi: 10.1253/circj.CJ-09-0402
- [50] Morrison, S., Colberg, S. R., Mariano, M., Parson, H. K., and Vinik, A. I. (2010). Balance training reduces falls risk in older individuals with type 2 diabetes. *Diabetes Care* 33, 748–750. doi: 10.2337/dc09-1699
- [51] Motooka, M., Koike, H., Yokoyama, T., and Kennedy, N. L. (2006). Effect of dog-walking on autonomic nervous activity in senior citizens. *Med. J. Aust.* 184, 60–63.
- [52] Nault, I., Nadreau, E., Paquet, C., Brassard, P., Marceau, P., Marceau, S., et al. (2007). Impact of bariatric surgery-induced weight loss on heart rate variability. *Metabolism* 56, 1425–1430. doi: 10.1016/j.metabol.2007.06.006
- [53] Perugini, R. A., Li, Y., Rosenthal, L., Gallagher-Dorval, K., Kelly, J. J., and Czerniach, D. R. (2010). Reduced heart rate variability correlates with insulin resistance but not with measures of obesity in population undergoing laparoscopic Roux-en-Y gastric bypass. *Surg. Obes. Relat. Dis.* 6, 237–241. doi: 10.1016/j.soard.2009.09.012
- [54] Piestrzeniewicz, K., Luczak, K., Lelonek, M., Wrancisz, J. K., and Goch, J. H. (2008). Obesity and heart rate variability in men with myocardial infarction. *Cardiol. J.* 15, 43–49.
- [55] Pop-Busui, R. (2010). Cardiac autonomic neuropathy in diabetes: a clinical perspective. *Diabetes Care* 33, 434–441. doi: 10.2337/dc09-1294
- [56] Pop-Busui, R., Boulton, A. J., Feldman, E. L., Bril, V., Freeman, R., Malik, R. A., et al. (2017). Diabetic neuropathy: a position statement by the American diabetes association. *Diabetes Care* 40, 136–154. doi: 10.2337/dc16-2042
- [57] Pop-Busui, R., Cleary, P. A., Braffett, B. H., Martin, C. L., Herman, W. H., Low, P. A., et al. (2013). Association between cardiovascular autonomic neuropathy and left ventricular dysfunction: DCCT/EDIC study (Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications). *J. Am. Coll. Cardiol.* 61, 447–454. doi: 10.1016/j.jacc.2012.10.028
- [58] Pop-Busui, R., Evans, G. W., Gerstein, H. C., Fonseca, V., Fleg, J. L., Hoogwerf, B. J., et al. (2010). Effects of cardiac autonomic dysfunction on mortality risk in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Diabetes Care* 33, 1578–1584. doi: 10.2337/dc10-0125
- [59] Pop-Busui, R., Low, P. A., Waberski, B. H., Martin, C. L., Albers, J. W., Feldman, E. L., et al. (2009). Effects of prior intensive insulin therapy on cardiac autonomic nervous system function in type 1 diabetes mellitus: the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications study (DCCT/EDIC). *Circulation* 119, 2886–2893. doi: 10.1161/CIRCULATIONAHA.108.837369
- [60] Raskin, P., and Cincotta, A. H. (2016). Bromocriptine-QR therapy for the management of type 2 diabetes mellitus: developmental basis and therapeutic profile summary. *Expert Rev. Endocrinol. Metab.* 11, 113–148. doi: 10.1586/17446651.2016.1131119
- [61] Ravussin, E. (2010). Impact of six month caloric restriction on autonomic nervous system in healthy, overweight individuals. *Obesity* 18, 414–416. doi: 10.1038/oby.2009.408
- [62] Rodrigues, G. D., Gurgel, J. L., Goncalves, T. R., and da Silva Soares, P. P. (2018). Inspiratory muscle training improves physical performance and cardiac autonomic modulation in older women. *Eur. J. Appl. Physiol.* 118, 1143–1152. doi: 10.1007/s00421-018-3844-9
- [63] Roy, B., and Ghatak, S. (2013). Nonlinear methods to assess changes in heart rate variability in type 2 diabetic patients. *Arq. Bras. Cardiol.* 101, 317–327. doi: 10.5935/abc.20130181
- [64] Sassi, R., Cerutti, S., Lombardi, F., Malik, M., Huikuri, H. V., Peng, C. K., et al. (2015). Advances in heart rate variability signal analysis: joint position statement by the e-Cardiology ESC Working Group and the European Heart Rhythm Association co-endorsed by the Asia Pacific Heart Rhythm Society. *Europace* 17, 1341–1353. doi: 10.1093/europace/euv015
- [65] Scirica, B. M., Bhatt, D. L., Braunwald, E., Steg, P. G., Davidson, J., Hirshberg, B., et al. (2013). Saxagliptin and cardiovascular outcomes in patients

- with type 2 diabetes mellitus. *N. Engl. J. Med.* 369, 1317–1326. doi: 10.1056/NEJMoa1307684
- [66] Selvarajah, D., Cash, T., Davies, J., Sankar, A., Rao, G., Grieg, M., et al. (2015). SUDOSCAN: a simple, rapid, and objective method with potential for screening for diabetic peripheral neuropathy. *PLoS One* 10:e0138224. doi: 10.1371/journal.pone.0138224
- [67] Servant, D., Logier, R., Mouster, Y., and Goudemand, M. (2009). [Heart rate variability. Applications in psychiatry]. *Encephale* 35, 423–428. doi: 10.1016/j.encep.2008.06.016
- [68] Sgoifo, A., Carnevali, L., Alfonso, M. L., and Amore, M. (2015). Autonomic dysfunction and heart rate variability in depression. *Stress* 18, 343–352. doi: 10.3109/10253890.2015.1045868
- [69] Silva-E-Oliveira, J., Amelio, P. M., Abranches, I. L. L., Damasceno, D. D., and Furtado, F. (2017). Heart rate variability based on risk stratification for type 2 diabetes mellitus. *Einstein* 15, 141–147. doi: 10.1590/S1679-45082017AO3888
- [70] Smith, A. G., Lessard, M., Reyna, S., Doudova, M., and Singleton, J. R. (2014). The diagnostic utility of Sudoscan for distal symmetric peripheral neuropathy. *J. Diabetes Complications* 28, 511–516. doi: 10.1016/j.jdiacomp.2014.02.013
- [71] Soedamah-Muthu, S. S., Chaturvedi, N., Witte, D. R., Stevens, L. K., Porta, M., and Fuller, J. H. (2008). Relationship between risk factors and mortality in type 1 diabetic patients in Europe: the EURODIAB Prospective Complications Study (PCS). *Diabetes Care* 31, 1360–1366. doi: 10.2337/dc08-0107
- [72] Spallone, V., Ziegler, D., Freeman, R., Bernardi, L., Frontoni, S., Pop-Busui, R., et al. (2011). Cardiovascular autonomic neuropathy in diabetes: clinical impact, assessment, diagnosis, and management. *Diabetes Metab. Res. Rev.* 27, 639–653. doi: 10.1002/dmrr.1239
- [73] Straznicki, N. E., Lambert, G. W., McGrane, M. T., Masuo, K., Dawood, T., Nestel, P. J., et al. (2009). Weight loss may reverse blunted sympathetic neural responsiveness to glucose ingestion in obese subjects with metabolic syndrome. *Diabetes* 58, 1126–1132. doi: 10.2337/db08-1595
- [74] Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology (1996). Heart rate variability: standards of measurement, physiological interpretation and clinical use. *Circulation* 93, 1043–1065. doi: 10.1161/01.CIR.93.5.1043
- [75] Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J. III, and Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neurosci. Biobehav. Rev.* 36, 747–756. doi: 10.1016/j.neubiorev.2011.11.009
- [76] The Advance Collaborative Group (2008). Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N. Engl. J. Med.* 358, 2560–2572. doi: 10.1056/NEJMoa0802987
- [77] The Consensus Committee of the American Autonomic Society and the American Academy of Neurology (1996). Consensus statement on the definition of orthostatic hypotension, pure autonomic failure, and multiple system atrophy. *Neurology* 46:1470. doi: 10.1212/WNL.46.5.1470
- [78] The Diabetes Control and Complications Trial Research Group (1995). The effect of intensive diabetes therapy on the development and progression of neuropathy. *Ann. Intern. Med.* 122, 561–568.
- [79] Tulppo, M. P., Hautala, A. J., Makikallio, T. H., Laukkanen, R. T., Nissilä, S., Hughson, R. L., et al. (2003). Effects of aerobic training on heart rate dynamics in sedentary subjects. *J. Appl. Physiol.* 95, 364–372. doi: 10.1152/jappphysiol.00751.2002
- [80] Valenza, G., Citi, L., Saul, J. P., and Barbieri, R. (2018). Measures of sympathetic and parasympathetic autonomic outflow from heartbeat dynamics. *J. Appl. Physiol.* 125, 19–39. doi: 10.1152/jappphysiol.00842.2017
- [81] Young, L. H., Wackers, F. J., Chyun, D. A., Davey, J. A., Barrett, E. J., Taillefer, R., et al. (2009). Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *JAMA* 301, 1547–1555. doi: 10.1001/jama.2009.476
- [82] Zafir, B., Azencot, M., Dobrecky-Mery, I., Lewis, B. S., Flugelman, M. Y., and Halon, D. A. (2016). Resting heart rate and measures of effort-related cardiac autonomic dysfunction predict cardiovascular events in asymptomatic type 2 diabetes. *Eur. J. Prev. Cardiol.* 23, 1298–1306. doi: 10.1177/2047487315624747
- [83] Ziegler, D., Gries, F., Spuler, M., and Lessmann, F. (1992). The epidemiology of diabetic neuropathy. Diabetic Cardiovascular Autonomic Neuropathy Multicenter Study Group. *J. Diabetes Complications* 6, 49–57. doi: 10.1016/1056-8727(92)90049-Q