

# Bioaccumulation of Heavy Metals in Squid (*Loligo duvaucelii*) from Tuticorin Coastal Waters

K. Venu Babu<sup>1</sup>, B. Veerendra Kumar<sup>2</sup>, K Hari Chandra Prasad<sup>3</sup>

<sup>1</sup>Research Scholar (2019-2023), Department of Biotechnology GITAM School of Science GITAM (Deemed to be University) Visakhapatnam-530045, India

<sup>2</sup>Professor, Head Department of Biotechnology, GITAM School of Science GITAM (Deemed to be University) Visakhapatnam-530045, India

<sup>3</sup>Bureau Veritas (India) Pvt. Ltd. F-2 Thiru Vi Ka Industrial Estate, Phase III, Ekkattuthangal, Guindy, Chennai-600032, India

**Abstract:** This study investigates 9 elements both essential (Cr, Cu, Zn, Fe, Mn and Ni) and non essential (Cd, Hg and Pb) in the tissues and whole of the Squid *loligo duvaucelii* caught off Tuticorin coast (Coromandel Coast of India, Bay of Bengal). The level of elements was determined by Inductively coupled plasma mass spectrometry (ICP-MS). The concentration ranges found for these heavy metals, expressed on a wet weight basis, were as follows: Hg, Cd, Pb, Cu, Zn, Fe, Mn, Cr, and Ni concentrations in Squid muscle samples were 0.01 - 0.07, 0.11-0.67, 0.11-1.14, 0.52-6.08, 4.82-19.32, 0.08-5.84, 0-0.49 and 0-2.11, 0-0.92 ppm respectively. As for other cephalopod species, the liver showed the highest concentrations of many elements highlighting their role in bioaccumulation and detoxification processes. The mean values of highly hazardous metals in the muscle of the *Loligo duvaucelii*, were: Hg = 0.04, Cd = 0.350, Pb = 0.300, Cr = 0.362, all within the international safety limits. The level of contamination in *Loligo duvaucelii* by these heavy metals is compared to those studied in other parts of the world and the legal standards set by international legalizations. The levels of heavy metal in *Loligo duvaucelii* was found to be within the safe limits for human consumption.

**Keywords:** Heavy metal, Bay of Bengal, Accumulation; Squid (*Loligo duvaucelii*)

## 1. Introduction

Cephalopod (Squids and Cuttlefish) represents an increasing component of the world's fisheries (Boyle and Rodhouse 2006; FAO 2007). They are consumed throughout the world, both as food and as feed supplement and have a great commercial value (Navarro and Villanueva 2001). Indeed, they are actually much appreciated as a protein source and because of their gustative quality. Cephalopods are consumed not only fresh, but also manufactured into processed food in huge quantities such as dry, frozen and chilled products (Paredi and Crupkin 1997; Hurtado *et al.* 2001). Trace metals are accumulated by marine invertebrates from both water and food. Trace metal accumulation process vary within the same invertebrate species according to the metal concerned. Some metals, such as Cu and Zn, are essential for proper metabolic functioning, whereas others, like Cd and Hg, are highly toxic to aquatic organisms and have unknown biological activities. Copper is an active centre of both the metal enzymes and the oxygen-transporting proteins, e.g. hemocyanin superoxide dismutase. Zinc is a co factor of several enzymes, such as carbonic anhydrase, alkaline phosphatase and DNA and RNA polymerase. In spite of its biological function, both Zn and Cu could produce toxic problems at very high levels.

Trace metals, particularly cadmium (Cd), mercury (Hg), lead (Pb) etc are persistent pollutants bio-accumulated by marine animals, risking public health through seafood. There has been increasing concern over the safety of food items that may contain harmful chemicals. Many fish-producing, as well as fish-importing countries have therefore instituted regulations and stringent quality requirements and standards for many chemical hazards including toxic metals in fish and fishery products. The EU directive of 1991 (91/493/EEC) and the US regulations of 1997 have set safety limits for such

hazards. It has become mandatory for all fish-exporting countries to monitor the levels of trace metals in their fishery products.

The present paper reports the baseline concentrations of some trace metals in the economically important Squid *loligo duvaucelii* from the Tuticorin south Coromandel Coast of India. These Squids occur in shallow coastal waters up to a depth of 80 m on the continental shelf of both coasts of India. From 1997 to 1998, India exported 75,000 metric tons (mt) of cephalopods and earned US \$163 million in foreign exchange. However, these products were sometimes detained or rejected from the EU markets on the ground that they contain high levels of Pb, Cd or the bacterium *Salmonella*. Heavy metals in Cephalopods have been extensively studied in various parts of the world as in India (Sadig and Alam 1989, Dious and Kasinathan 1992, Prudente *et al.* 1997). Higher levels of cadmium and other metals in squids and Cuttlefish have been reported by Falandysz (1989, 1990, 1991). At present, only a few studies on metal concentrations in Cephalopods from the Indian waters (Lakshmanan 1988, Patel and Chandy 1988, Lakshmanan and Stephen 1993) exist. This paper reports the levels of 9 trace metals in Squid *loligo duvaucelii* from Tuticorin which is major fishing harbor in the Coromandel Coast of India.

## 2. Materials and Methods

Fresh Squid samples were collected directly from fishermen and processing plants landed and processed in the fisheries harbor and processing plants, Tuticorin from August 2021 to February 2022. They were fished off the Coromandel Coast of India. Samples were taken at monthly intervals. The average length and whole body weight of the samples were  $100 \pm 8.00$  mm and weight  $110 \pm 15.0$  g respectively, which

were of commercial grades. Samples were immediately iced and brought to the laboratory in insulated boxes. These were either analyzed fresh or kept frozen (-18°C) until analysis. The whole (after removing the cuttlebone) and whole cleaned soft parts were finely chopped and homogenized and aliquots were taken for wet digestion. Another lot was carefully dissected for various body components, such as muscle, liver, and gills, and subjected to wet digestion using concentrated nitric acid and perchloric acid (AOAC 1990). Metal concentrations were determined using Atomic Absorption Spectrophotometer model (GBC 902) with an oxidizing air-acetylene flame or an Inductively Coupled Plasma Emission Spectroscopy. Samples for mercury analysis were wet digested using concentrated HNO<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub> in the ratio (4:1 v/v) in a Bethge's apparatus. Determination of mercury was carried out following the cold vapor Atomic Absorption technique, using a Mercury Analyzer (model MA 5843) after reducing with stannous chloride and hydrochloric acid. Triplicate analysis was carried out in each case and the data were statistically analyzed.

De-ionized water was used to prepare all aqueous solutions. All mineral acids and oxidants used were of the highest quality (Merck, Germany). All the plastic and glassware were cleaned by soaking overnight in a 10% (w/v) nitric acid solution and then rinsed with deionized water.

An independent *t* test analysis was applied to the observed data to check the level of significance site-wise. Pearson's Correlation Coefficient Statistical functions on Microsoft Excel were used to test the relation between the metal concentrations and the mantle length of the specimens.

### 3. Results and Discussion

Concentrations of 9 heavy metals found in the whole soft tissues of Squid species, *Loligo duvauceli* and in various body components are illustrated in figure 1 and table 2.

Levels of contaminants in marine animals are of particular interest because of the potential risk to humans who consume them. In particular, the tendency of heavy metals to accumulate in various organs of cephalopods, which in turn may enter the human metabolism through consumption causing serious health hazards. Accumulation of metals in marine animals is the function of their respective membrane permeability and enzyme system, which is highly species specific and because this fact different metals accumulated in different orders in the studied Squid samples. Therefore, the present study was undertaken to evaluate the heavy metals accumulation of Hg, Fe, Zn, Cu, Pb, Cr, Ni, Mn and Cd in the mantle tissues of the edible Squid, *Loligo duvauceli*.

#### 3.1 Lead

Lead constitutes a serious health hazard to both children and adults. The adverse toxic effect caused by Pb on human was documented by Subramanian (1988). The high blood lead level can cause kidney dysfunction, brain damage, anaemia and can inhibit the normal functioning of many enzymes (Forstner and Wittmann 1983; USFDA 1993).

In the whole soft parts of Squid (*Loligo duvauceli*), Pb levels varied from 0.46 to 1.48 ppm, in the whole samples analysed 9% showed Pb content above the tolerance limit of 1.5 ppm. However, mean Pb content in the edible muscle was below 1 ppm the range of values being 0.1 to 1.14 ppm at Tuticorin. The lower range of Pb, in the various body components analysed in general was zero. In the liver concentration of Pb ranged from 0 to 4.31 ppm. The mean Pb content was highest in gill samples of Squid, the distribution pattern in general being gills > liver > muscle.

In the present study concentrations of Pb in muscle of Squid (*loligo duvauceli*) were less than the permissible limit (1.5 mg/kg, Table 1.) and were comparable to the values reported in *L. opalescens* (Falandysz, 1991). The levels of Pb observed in the liver of Squid *loligo duvaucelii* were similar to the values reported in *Sepia pharaonis* (Lakshmanan and Stephen, 1993; Prafulla *et al.*, 2001; Saleh Al Farraj *et al.*, 2011). The lower range of Pb, in general was zero indicating more or less a cleaner environment. Another reason may be due to the sparingly soluble property of lead compounds and consequently non-availability to bio-mass in the marine environment. As observed in the present study a greater concentration of Pb was found in the non edible parts of Squid. Also, 9% of whole samples had Pb content above 1.5 ppm, the tolerance limit.

#### 3.2 Cadmium

Concerning Cadmium, the capability of cephalopods to concentrate high quantities of cadmium is well known (Miramand and Guary 1980; Miramand and Bentley 1992; Bustamante *et al.* 1998). The mantle of Squid does not mentioned before as a specific organ for Cd accumulation, whereas, the digestive gland plays a primary role in the accumulation of Cd in Squid. Such a role can be related both to the very efficient detoxification processes of the metal occurring in this organ and to the very high Cd assimilation efficiency (Bustamante *et al.* 2002a; Bustamante *et al.* 2006).

Dorneles *et al.* (2007) reported very high Cd concentrations; reach up 1000 µg g<sup>-1</sup> wet wt, in the digestive gland of the short-lived species *Illex argentinus*. They stated that this value represents the highest cadmium level ever reported for a cephalopod. However, in this study, Concentrations of Cd in whole soft tissues of Squid was 3.247 ± 1.553 ppm. Around 20% of the whole Squid had Cd content > 3 ppm, the tolerance limit. Mean Cd content in the edible muscle was 0.481 ppm, liver was the major site of Cd accumulation in Squid and it contributed significantly to the total body burden of Cd. Liver samples from Tuticorin region had Cd content in the range of 4.2 to 114.0 ppm. However, 25% of the gill samples from Tuticorin region showed Cd content above 2 ppm. The increasing order of abundance of Cd in the body components of Squid was liver > gills > muscle.

Higher levels of Cd in whole *Loligo* spp. had been reported from various parts of the world. Falandysz (1989) found high levels of Cd (2.9 to 10mg/kg wet wt) in the edible parts of canned squid, *Loligo patagonica*. Raw whole squid contained on an average 4.0 mg Cd/kg as reported by Falandysz (1991). Hussain and Mukundan (2005) reported

the cadmium levels in edible parts like meat and tentacles of *Sepia pharaonis* and *Uroteuthis (Photololigo) duvauceli*, the level of cadmium was found to be below 1 ppm.

Lourenço, *et al.* (2009) found that Cd concentrations in the arms and the mantle of *Sepia officinalis* ranged between 0.03 and 1.0 mg kg<sup>-1</sup> basis on wet wt. Saleh Al Farraj *et al.*, (2011) found the average concentration of Cd in the mantle of *Sepia pharaonis* was 0.04 mg/ kg, and higher levels were observed in the liver of Squid. The present results exceeded the maximum levels permitted (2 µg g<sup>-1</sup>) in molluscs that are recommended by FSA (2002) and disagree with Prafulla, *et al.* (2001) who stated that in the squid, mantle tissues accumulated relatively low values for Cd and was far below the tolerance limit. Generally, many studies have suggested that squid mantle could be consumed safely in most countries, considering the Cd concentration is two orders of magnitude lower than those in the digestive gland (Kim *et al.* 2008).

### 3.3 Mercury

Contamination of marine organisms with toxic metals such as mercury (Hg) is of ecological and health concern worldwide (Goldberg, 1995). The presence and behaviour of Hg in aquatic systems are of great interest and importance since it is the only heavy metal which bio accumulates and biomagnifies through all levels of the aquatic food chain (Lindqvist *et al.*, 1991). Mercury has not necessary function in any living organism and is considered as a non-essential metal, is among the most toxic elements to man and many higher animals (Steinnes, 1995).

Mercury was the least abundant toxic metal found in Squid (*Loligo duvauceli*). In whole soft tissues, Hg levels were in the range of 0.035 to 0.098 ppm. The highest value observed for Hg in whole Squid was 0.098 ppm and a reduction of around 30% of Hg content was seen in the edible muscle. Highest mean values of Hg in edible muscle were noted was 0.042 ppm. Among the other body components analysed, only the liver exhibited a value of > 0.10 ppm for Hg and mean concentration was 0.186± 0.0419 ppm. Liver of Squid caught off the Tuticorin region recorded the highest values, *viz* 0.291 ppm (Table 2). Only 12% of the liver samples showed Hg content above 0.2 ppm. Hg content in gills was in the range 0.032 to 0.070. The distribution pattern of Hg in the body components analysed was of the order: liver > gills > muscle.

Comparable levels of Hg have been reported in cephalopods caught from the Arabian Sea (Patel and Chandy 1988m Lakshmanan and Stephen 1993). Only low levels of mercury were reported in *Loligo opalescens* during the past several years (Falandysz 1989, 1990). In this study the levels of mercury in the different body components were also low, and the distribution followed the order liver > gills > muscle.

### Chromium

Chromium is an essential trace element for human body for glucose tolerance (Dung 2001). Despite its high toxicity at high doses (Burger and Gochfeld 1995), mammalian bodies can tolerate 100 to 200 times their total natural body content of Cr without harmful effects (Moore and Ramamoorthy

1984).

The concentrations of Cr are presented in Table 2. The mean Cr content in whole Squid (*Loligo duvauceli*) was 3.616 ppm. Around 11% of whole Squid showed Cr content above 12 ppm. However, Cr content in the muscle of Squid was below the tolerance limit. The lower range of Cr in the edible muscle in general, was zero. The mean value of Cr content in the liver was 2.317 ± 2.312 at Tuticorin. The concentrations of Cr in the different organs of Squid was followed the order: liver > gills > muscle.

Schumacher *et al.* (1992) found that the mean concentrations of chromium in the flesh of cephalopods were ranged from 0.090 to 0.220 µg/g of fresh weight. On the other hand, Prafulla *et al.* (2001) found that the mean muscle value for Cr in the two squids species *Loligo duvauceli* and *Doryteuthis sibogae* was < 1.0 ppm. After their study on some species of the Squid *Sepia*, Ahdy *et al.* (2007) concluded that the average concentration of Cr was 2.2 µg/g. From other point of view, Cr concentrations were higher in juveniles than in adults, in the squid *Sthenoteuthis oualaniensis* (Ichihashi *et al.* 2001).

### Zinc

Zinc is an essential trace element required by all living organisms because of its critical role both as a structural component of proteins and as a cofactor in enzyme catalysis (Lall 1989; Sandstrom 1997; Leigh Ackland and Michalczyk 2006). Zinc is always present in seafood, but concentrations found in mollusks are generally higher (Lall 1995; Celik and Oehlschläger 2004). Zn is involved in numerous protein functions such as the carbonic anhydrase and is efficiently absorbed and strongly retained in *Loligo duvauceli* both from the food and seawater pathways (Villanueva and Bustamante 2006).

The range of values of Zn in the whole soft tissues of Squid was 5.84 to 24.16. In general, the level of Zn was highest in the liver of Squid (Figs. 1). Around 40% of the liver samples from Tuticorin region had Zn content above 50 ppm. However, much lower levels were observed in the muscle with values ranging from 4.82 to 19.32 ppm (Tuticorin). In general, all the Squid gill samples showed Zn content below 50 ppm.

Moreover, it is lower than the values determined by Lourenço *et al.* (2009) in arms and mantles of some common Portuguese Squid (14.0- 22.5 mg kg<sup>-1</sup> wet weight). Craig and Overnell (2003) found that Zn present in the cytosol, as a percentage of total tissue metal, in squid *Loligo forbesi* tissues, in mantle was 11.5, while in the digestive gland was 43.0.

This agrees with the results of previous studies which reported an increase of zinc concentration stored in metal-containing enzymes and metalloproteins in the digestive gland of the cephalopods (e.g. Miramand and Bentley 1992; Gerpe *et al.* 2000; Villanueva and Bustamante 2006; Lacoue-Labarthe *et al.* 2008). Accordingly, although Zn is an essential element for most cephalopods, mantle tissues still accumulate the lower values.

### Copper

Mollusks, particularly cephalopods use the soluble copper containing protein haemocyanin as a respiratory pigment, therefore copper is required in large concentrations (Decleir *et al.* 1978; Viant *et al.* 2002). In addition, Martoja and Marcaillou (1993) stated that Cu accumulates in membrane-limited vesicles (spherules) within the basal cells of the liver the *Sepia officinalis*. They postulated that copper was bound to a “metallothionein-like” protein which has been detected in heat-treated liver cytosols from cephalopods (Bustamante *et al.* 2002a).

Among the various body components of *Loligo duvauceli*, muscle had the lowest Cu content and ranged from 0.18 to 7.45 ppm in Squid. Comparatively higher levels of Cu were found in the liver and gills of Squid (Fig..1). The range of values of Cu in the liver of Squid from Tuticorin, were 14.21 to 128.32, ppm,. 38% of Cu in liver samples from Tuticorin region showed Cu levels above 50 ppm. However, Cu content in the muscle samples reflected low levels of Cu. Hence, the mantle tissue looks as if it is not the target organ for Cu accumulation.

The involvement of this element in several metabolic functions, such as in metal-dependent enzymes (Bustamante *et al.* 2000; Craig and Overnell 2003) may explain its high concentration in the present species. The levels of Cu observed in the muscle of Squid (*loligo duvauceli*) are similar to the values reported in the mantle of *Loligo duvauceli* in Portnovo samples by Dious and Kasinathan (1992). The copper levels in muscle are also quite comparable to the levels in *L duvauceli* reported by Tariq *et al.*, (1991) and Prafulla *et al.*, (2001). Very high levels of Cu in the liver of *L opalescens* (200 to 300 mg/kg wet wt.) were reported by Falandysz (1991). Martin and Flegal (1975) and Oehlschlaeger (1991) also reported high Cu content in the liver of *L opalescens*. Copper content in the liver of Squid are quite comparable to the values reported by Lakshmanan and Stephen (1993) ( $56.16 \pm 45.67$  ppm). Since Squid requires Cu for the synthesis of the respiratory pigment, haemocyanin, a high level in the body may be attributed to the functional necessity (Smith, 1984).

### Nickel

A few years ago, the physiological functions of nickel (Ni) were unclear. Nowadays, it is thought that Ni plays an important role in hormone, lipid and cell membrane metabolism. Several studies on nickel indicated to the role of nickel in various enzymatic system as well as activating enzymes associated with the glucose breakdown and use (Nielsen *et al.* 1984; Lall 1995; Acu-Cell2007).

Nickel (Ni) in excess is toxic to aquatic organisms and persistent in the aquatic environment. In the present study, the distribution of Ni is shown in Fig 1 and Table 1. Nickel content in whole Squid was in the range 0.19 to 5.92 ppm. However, Ni content in the muscle of Squid was below the tolerance limit the mean value being 0.297, slightly elevated levels of Ni were found in the liver of Squid and Ni levels ranged from 0 to 5.32 ppm in gills. The distribution pattern of Ni in the body components were liver > gills > muscle (Table 2).

Ni levels in the edible muscle of Squid *Ioligo duvaucelii* were several folds lower than levels reported by Prafulla *et al.*, (2001). Lourenço, *et al.*, (2009) identified Ni content in the edible parts (arms and mantle) of the Squid *Sepia officinalis* was 0.02-0.09 mg/kg wet wt. Villanueva and Bustamante (2006) observed low concentrations of Ni in the European Squid *Sepia officinalis* as compared to the European squid *Loligo vulgaris* and the common octopus *Octopus vulgaris*. Concerning the relationship between Ni concentration and specimen sizes of squid, Ichihashi *et al.*, (2001) have noticed that the concentration of Ni was greater in adults of the squid *Sthenoteuthis oualaniensis* than in the juveniles. In the present study the Ni content is in permissible limits in all the samples from all the three regions.

### 3.8 Iron

Iron, (Fe) one of the most abundant metals on earth, is essential to most life forms and to normal human physiology. Iron is an integral part of many proteins and enzymes that maintain good health (Institute of Medicine, 2001). In humans, iron is an essential component of proteins involved in oxygen transport (Dallman, 1986). It is also essential for the regulation of cell growth and differentiation (Bothwell, 1979, Andrews, 1986). A deficiency of Fe limits oxygen delivery to cells, resulting in fatigue, poor work performance, and decreased immunity (Institute of Medicine, 2001, Bhaskaram, 2001). On the other hand, excess amounts of iron in human can result in toxicity and even death (Corbett, 1995).

The liver of Squid was rich in Fe content with a mean value of 71.981. The Fe content in liver ranged from 3.24 to 141.56 ppm in Squid *Ioligo duvaucelii*. Distribution of Fe in whole Squid samples from Tuticorin was 5.201. The range of Fe values in muscle of Squid from Tuticorin, were: 0.08 to 5.84 (Table 2).

The level of Fe found in the various body components of Squid (*Ioligo duvauceli*) was quite comparable to the values observed in *L opalescens* (Falandysz 1991) and *L duvauceli* (Prafulla 2001). Liver of *Todarodes pacificus* from Japan showed higher level of Fe in visceral organs than in the edible parts (Ueda *et al.*, 1979). Lozano Soldevilla (1989) found Fe content to be high in whole bodies of *Todarodes sagittatus* when compared to those in mantles and tentacles.

### 3.9 Manganese

Manganese (Mn) is essential in animal body and its deficiency is impaired in the glucose utilization. Mn was found only at low levels in Squid. However, in whole soft parts of Squid the Mn content ranged from 0.11 to 5.46 ppm at Tuticorin. In general, liver contained higher levels of Mn compared to other organs. The mean values being 3.18, ppm at Tuticorin. The average levels of Mn in gills showed a more or less homogenous distribution pattern;

Manganese (Mn) is essential in animal body and its deficiency is impaired in the glucose utilization. Mn was found only at low levels in Squid. The lower range of Mn in the muscle was zero. However, in whole soft parts of Squid

the Mn content ranged from 0.11 to 5.46 ppm. In general, liver contained higher levels of Mn compared to other organs the mean values is 3.18

Miramand *et al.*, (2006) found that the highest significant average Mn concentration approximately 4.0 mg/kg was found in Squid and squid. The level of Mn in the whole of Squid observed in the present study compares well with the values for whole *L. duvauceli* reported by Prafulla *et al.*, (2001). Wide ranges in the concentrations of the elements Fe, Mn, Pb and Zn were observed in the digestive gland of squid, *Nototadarus gouldi* by Smith *et al.*, (1984).

#### 4. Conclusion

The highly toxic metals, viz., Cd, Pb, and Cr often exceeded the tolerance limit in around 20% (Cd), and 11% (Pb and Cr) in whole Squid *Ioligo duvauceli*. However, the mean content of all the metals analysed were significantly lower in the edible parts and far below the tolerance limits. Concentration of Hg was found to be <50 ng/kg in the edible muscle in 90% of the samples and Hg content was far below the limit of 1 mg/kg permitted for seafoods by many fish importing nations and USFDA. Cd content in muscle was < 3 ppm in Squid from all the regions. Squid liver was the major site of accumulation of Cd and other toxic metals. The increasing order of abundance of most of the metals in Squids was: Liver > Gills > Muscle. As liver is the major site of accumulation, in Squids consumption of liver should be avoided at any cost. Gutting stage should be included as a Critical Control Point (CCP) during Squid processing ensuring seafood safety.

#### References

- [1] Acu-Cell, 2007. Available at: <<http://www.acu-cell.com/acn.html>>. Accessed September.
- [2] Boyle P.R., Rodhouse P.G., 2006. Cephalopods: ecology and fisheries. Blackwell Science, Oxford.
- [3] Bustamante P, Caurant F, Fowler S.W, and Miramand P. Cephalopods as a vector for the transfer of cadmium to top marine predators in the north-east Atlantic Ocean. *Science of the Total Environment*, 1998; 220:71-80.
- [4] Bustamante P, Cosson R.P, Gallien I, Caurant F, Miramand P. Cadmium detoxification processes in the digestive gland of cephalopods in relation to accumulated cadmium concentrations, *Marine Environmental Research*, 2002; 53: 227-241.
- [5] CE (Commission Européenne), 2006. Reglement (CE) No. 1881/2006 de la Commission du 19 Décembre 2006.
- [6] Craig S, Overnell J, Metals in squid, *Loligo forbesi*, adults, eggs and hatchlings. No evidence for a role for Cu- or Zn-metallothionein, *Comparative Biochemistry and Physiology*, 2003;134: 311-317.
- [7] Dios S.R.J, Kasinathan R, Concentration of Fe, Mn, Zn and Cu in cephalopod *Loligo duvauceli*, *Indian J. Mar. Sc.*, 1992; 21: 224-225.
- [8] Dorneles, P.R, Lailson-Brito J, Machado Torres J.P. Cephalopods and cetaceans as indicators of offshore bioavailability of cadmium off Central South Brazil Bight, *Environmental Pollution*, 2007; 148: 352-359.
- [9] FAO, 2007. Fisheries commodities production and trade 1976–2005, Food and Agriculture Organization of the United Nations, Fisheries and Aquaculture Information and Statistics Service, available at [www.fao.org/fi/statist/FISOFT/FISHPLUS.asp](http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp)
- [10] Forstner N, Wittmann G, 1983, “Metal Pollution in the Aquatic Environment”, Berlin: Springer-Verlag.
- [11] FSA, 2002, (Food Standards Australia), Issue 53, Standard 1.4.1 and 1.4.2.
- [12] Hurtado J.L, Montero P, Borderial J, Chilled storage of pressurized octopus (*Octopus vulgaris*) muscle, *Journal of Food Science*, 2001; 66: 400-406.
- [13] Ichihashi H, Kohno H, Kannan K, Tsumura A, Yamasaki S.I, Multielemental analysis of purple back flying squid using high resolution inductively coupled plasma-mass spectrometry (HR ICP-MS), *Environmental Science and Technology*, 2001; 35: 3103-3108.
- [14] Lacoue-Labarthe T, Warnau M, Teysse J.L, Bustamante P, Bioaccumulation of inorganic Hg by the juvenile Squid *Sepia officinalis* exposed to <sup>203</sup>Hg radio labeled seawater and food, *Aquatic Biology*, 2008; 91-98.
- [15] Lall, S.P, 1995. Macro and trace elements in fish and shellfish, Fish and fishery products. Composition, nutritive properties and stability, Wallingford: Cab International, pp. 187–213.
- [16] Lourenço H.M, Anacleto P, Afonso C, Ferrara V, Nunes M. L, Elemental composition of cephalopods from Portuguese continental waters, *Food Chemistry*, 2009; 113: 1146-1153.
- [17] Lozano Soldevilla F, Galendo-Martin L, Concentrations of heavy metals in various parts of *Todarodes sagittatus* from the Saharan Bank, *Alimentaria*, 1989; 203: 23-26.
- [18] Martin J.H., Flegal A.R, High copper concentrations in squid livers in association with elevated levels of silver, cadmium and zinc, *Marine Biology*, 1975; 30: 51-55.
- [19] Miramand P, Bentley D, Concentration and distribution of heavy metals in tissues of two cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English Channel, *Marine Biology*, 1992; 114: 407-414.
- [20] Miramand P, Guary J.C, High concentrations of some heavy metals in tissues of the Mediterranean octopus, *Bulletin of Environmental Contamination and Toxicology*, 1980; 24: 783-788.
- [21] Navarro J.C, Villanueva R, Lipid and fatty acid composition of early stages of cephalopods: an approach to their lipid requirements, *Aquaculture*, 2001; 183: 161-177.
- [22] Nielsen F.H, Shuler T.R, McLeod T.G, Zimmerman T.J, Nickel influences iron metabolism through physiologic, pharmacologic and toxicological mechanisms in the rat, *Journal of Nutrition*, 1984; 114: 1280-1288.
- [23] Oehlenschlaeger J, Heavy metal content of deep frozen Californian squid, *Inf. Fischwirtsch*, 1991; 38(2): 62-68.
- [24] Paredi M.E, Crupkin M, Biochemical properties of actomyosin from frozen stored mantles of squid (*Illex argentinus*) at different sexual maturation stages, *Journal of Agricultural and Food Chemistry*, 1997; 45:1629-1632.
- [25] Prafulla V, Francis L, Lakshmanan P.T, Concentrations of Trace Metals in the Squids, *Loligo duvauceli* and

Doryteuthis sibogae Caught from the Southwest Coast of India, Asian Fisheries Science, 2001; 14: 399-410. the Arabian Gulf, Bulletin of Environmental Contamination and Toxicology, 1989; 42: 111-118.

- [26] Smith J.D, Plues L, Heryraud M, Chery R.D, Concentrations of the elements Ag, Al, Ca, Cd, Cu, Fe, Mg, Pb and Zn and the radio nuclides 210Pb and 21cPo in the digestive gland of the squid Notodarus gouldi, Marine Environmental Research, 1984; 13: 55 - 68.
- [27] Subramanian K.S, 1988. "Lead. in Quantitative Trace Analysis of Biological Materials." Mckenzie, H.A. and Smythe, A.(Eds.), Amsterdam: Elsevier.
- [28] USFDA (United States Food and Drug Administration). 1993. Guidance document for lead in shellfish. Washington, DC: Center for Food Safety and Applied Nutrition.
- [29] Villanueva R, Bustamante P, Composition in essential and non-essential elements of early stages of

cephalopods and dietary effects on the elemental profiles of Octopus vulgaris paralarvae, Aquaculture, 2006; 261:225-240.

**Table 1:** Tolerable Daily Intake of Heavy Metals by Human as Prescribed by JECFA (2000)

Heavy metals	Provisional Maximum Tolerable Daily Intake (PMTDI) (mg/kg)
Pb	0.025
Cr	0.1-1.2
Co	0.005-1.8
Ni	0.1-0.3
Hg	0.025
Mg	250-380
Fe	0.8
Zn	0.3-1.0
Cu	0.05-0.5
Mn	0.4-10

**Table 2:** Trace metal concentrations (Mean ± S.D., Range, ppm wet wt) in the whole soft parts and body components of Squid loligo duvaucelii collected off Tuticorin Region

Whole/Body Component	n*	Mercury	Cadmium	Lead	Copper	Zinc	Iron	Manganese	Chromium	Nickel
Whole Squid with Tentacles	30	0.080 ± 0.010 (0.03-0.09)	3.240± 1.550 (0.420-6.10)	0.920 ± -0.270 (0.48-1.40)	12.023 ± 2.960 (4.92-15.83)	14.673± 4.340 (5.84-24.16)	5.201 ± 1.317 (2.32-10.11)	1.868± 1.396 (0.11-5.46)	3.616± 1.823 (0.05-6.43)	3.248± 1.406 (0.19-5.92)
Muscle	30	0.04 ± 0.016 (0.01-0.08)	0.470 ± 0.150 (0.11 - 0.65)	0.525 ± 0.295 (0.11 - 1.14)	2.489 ± 1.451 (0.52-6.08)	9.449± 4.056 (4.82-19.32)	3.182± 1.595 (0.08-5.84)	0.310± 0.092 (0 - 0.49)	0.662 ± 0.479 (0 - 2.11)	0.297± 0.243 (0-0.92)
Liver	30	0.180± 0.038 (0.16-0.30)	52.00 ± 40.025 (4.20-14.00)	0.530± 0.330 (0.1-1.10)	68.530 ± 33.354 (14.21-128.32)	5.0332± 25.207 (28.41-149.87)	71.981± 42.249 (3.24-141.56)	3.181 ± 1.876 (0-9.13)	2.317 ± 2.312 (0-7.11)	1.854± 1.478 (0.21-5.86)

\* Number of samples analyzed.

**Table 3:** Standard Preparation

S.No	Element	Stock solution (A) (ppm)	Preparation of stock solution (B) ppm	Conc. of stock solution (B) ppm	Preparation of working stock soln. (C) ppm	Conc. of working stock solution (C) ppm
1	Pb	1000	2.5 ml of A is diluted to 50 ml	50	1 ml of B is diluted to 50ml	1
2	As					
3	Cd					
4	Hg					
5	Cu					
6	Ca					
7	Cr					
8	Ni					
9	Na					
10	K					
11	Fe					
12	Mn					
13	Mg					
14	Se					
15	Sb					
16	Sn					
17	Si					
18	P					
19	Zn					

Table 4: Analysis sequence

S. No	Sample ID	No. of Replicates	S. No	Sample ID	No. of Replicates	S. No	Sample ID	No. of Replicates
1	Blank(c)	3	11	Acid Blank	3	21	Sample-7	3
2	STD-1	3	12	QA STD	3	22	Sample-8	3
3	STD-2	3	13	Acid Blank	3	23	Sample-9	3
4	STD-3	3	14	Blank-P	3	24	Sample-10	3
5	STD-4	3	15	Sample-1	3	25	Sample-Duplicate	3
6	STD-5	3	16	Sample-2	3	26	Acid Blank	3
7	STD-6	3	17	Sample-3	3	27	QA STD	3
8	STD-7	3	18	Sample-4	3	28	Acid Blank	3
9	STD-8	3	19	Sample-5	3			
10	STD-9	3	20	Sample-6	3			

Table 5 Linearity ranges

S.NO	ELEMENT	RANGE	Unit mg/lit
1	Lead (pb)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
2	Arsenic (As)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
3	Cadmium(Cd)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
4	Copper(Cu)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
5	Chromium(Cr)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
6	Nickel(Ni)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
7	Iron(Fe)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
8	Manganese(Mn)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
9	Zinc(Zn)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
10	Selenium(Se)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
11	Antimony(Sb)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
12	Tin(Sn)	0.01,0.02,0.05,0.1,0.2,0.5,1.0,2.0,5.0,	mg/L
13	Phosphorus(P)	0.10,0.20,0.50,1,2,5	mg/L
14	Sodium(Na)	1.0,2.0,5.0,10.0,20.0,50.0,	mg/L
15	Pottasium(K)	1.0,2.0,5.0,10.0,20.0,50.0,	mg/L
16	Calcium(Ca)	1.0,2.0,5.0,10.0,20.0,50.0,	mg/L
17	Magnesium(Mg)	1.0,2.0,5.0,10.0,20.0,50.0,	mg/L
18	Mercury (Hg)	0.01,0.02,0.05,0.1,0.2	mg/L
19	Silicon(Si)	0.10,0.20,0.50,1,2,5	mg/L

5.7.1 Aspiret

55.7.7 Aspi