

Comparative Study of Heart Rate Variability between Young Premenopausal and Postmenopausal Female Medical Professionals at Resting Supine Posture

Dr. Anuradha Upadhyay¹, Dr. Manila Jain²

¹Ph. D. Research Scholar, Department of Physiology, Index Medical College, Hospital and Research Centre, Indore, Madhya Pradesh, India-452016

Corresponding Author E-Mail: [dranuradha1physio\[at\]gmail.com](mailto:dranuradha1physio[at]gmail.com)

Mobile No: 8949236909

²Professor & Head, Department of Physiology, Index Medical College, Hospital and Research Centre, Indore, Madhya Pradesh, India-452016

Abstract: *Background & Objectives:* Heart rate variability (HRV) measures the degree of autonomic modulation and sympathovagal balance. Menopause causes an imbalance of the cardiac autonomic nervous control that shifts toward sympathetic predominance. Present study compared HRV in frequency domain spectral components, between young premenopausal and postmenopausal medical professional women in the resting supine posture. *Method:* It was a comparative, cross-sectional, observational study. After obtaining ethics committee approval, the study population was divided into two groups as per inclusion and exclusion criteria. One group included 40 young premenopausal women aged 18-24 years and the other group included 40 postmenopausal women aged 45-55 years. The power spectrum density for the HRV was calculated in frequency domain by the traditional Fast Fourier transform (FFT) based method. *Results:* Our study showed that 'Total Power' was significantly lower (p -value < 0.001) in postmenopausal women ($917.89+647.67 \text{ ms}^2$) as compared to young premenopausal women ($4900.82+2876.29 \text{ ms}^2$). 'LF Power' was significantly higher (p -value < 0.001) in postmenopausal women ($693.14+271.21 \text{ ms}^2$) as compared to young premenopausal women ($141.60+107.49 \text{ ms}^2$), while, 'HF Power' was significantly higher (p -value < 0.001) in young premenopausal women ($1186.65+783.73 \text{ ms}^2$) as compared to postmenopausal women ($134.97+109.85 \text{ ms}^2$). In the present study, 'LF/HF ratio' was significantly higher (p -value < 0.001) in postmenopausal women ($8.95+2.47$) as compared to young premenopausal women ($0.22+0.14$). Serum estradiol level was significantly lower in postmenopausal women while 'body fat percentage' was significantly higher in postmenopausal women. *Interpretation & Conclusion:* The 'Total Power', and 'HF Power' is significantly lower, while 'LF Power' and 'LF/HF ratio' is significantly higher in postmenopausal women, which suggests overall reduced parasympathetic drive and poor vasovagal balance. Further, decreased level of serum estradiol and increased body fat percentage indicate sympathetic hyperactivity in postmenopausal women that may lead to higher risk of cardiovascular morbidity and mortality.

Keywords: Heart rate variability, Sympathovagal balance, Total Power, LF/HF ratio

1. Introduction

Heart rate variability (HRV) is defined as the cyclic changes or fluctuations of heart periods (R-R intervals) over time during the respiratory cycle (respiratory sinus arrhythmia) at rest. HRV measures the degree of autonomic modulation. It is a good predictor of cardiovascular morbidity and mortality. HRV depends on the rate of discharge of SA node, which is influenced by autonomic nervous system. SA nodal discharge is mainly controlled by parasympathetic activity. During inspiration and expiration, variations in vagal tone occur, which may lead to respiratory sinus arrhythmia. HRV is largely influenced by parasympathetic activity, though sympathetic activity also influences it.

'Sympathovagal balance' can be well estimated by HRV. There are two components of the power spectrum of HRV. They are high frequency (HF) component and low frequency (LF) component. Parasympathetic activity is indicated by HF component, while sympathetic activity is expressed by LF component. LF/HF ratio reflects sympathovagal balance. Parasympathetic activity also indicated by 'Total Power'. Increased HF, reduced LF, and reduced LF/HF ratio indicate

parasympathetic predominance, whereas reduced HF, increased LF, and increased LF/HF ratio indicate sympathetic predominance. Predominant parasympathetic activity favors good cardiovascular health. Predominant sympathetic activity is an important cardiovascular risk factor.

There are three phases of the menstrual cycle in a young reproductive woman-menstrual phase, luteal phase and follicular phase. Due to cessation of ovarian function, menstrual cycle ceases at an approximate age of 45 to 50 years which is commonly termed as menopause. The termination of menstruation continuously longer than 12 months is a postmenopausal state. Many physiological changes occur during the menstrual cycle and postmenopausal period in women. These changes affect the cardiac functioning of the heart¹. The risk towards the cardiovascular diseases increases after menopause and extensive fluctuation in HRV, are seen during the menopause and even after the menopause. Postmenopausal women had lower total power, lower HF in absolute power, higher relative power of LF, and higher LF/HF ratio². Menopause causes an imbalance of the autonomic nervous

Volume 11 Issue 2, February 2022

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

control of the cardiovascular system that shifts toward sympathetic hyperactivity³.

Epidemiological studies have indicated that postmenopausal women had alteration in their autonomic status with higher sympathetic and lower vagal tone compared to premenopausal women⁴. A significant hypoestrogenism state occurs during postmenopausal period which promotes modifications in the autonomic control of heart rate. These modifications induce changes in sympathetic activity and vagal regulation of cardiovascular system. Akiyoshi et al⁵ evaluated that cardiac parasympathetic function is decreased in postmenopausal women, which is related associated to aging and hormone level. Schillaci G et al⁶ studied that vasomotor symptoms are associated with sympathetic predominance during postmenopausal period. The menopausal women have an imbalance of autonomic control of cardiovascular system. Various techniques and maneuvers have been developed to detect the integrity of autonomic nervous system. Bannister R et al⁷ observed that most of the techniques such as cold pressure test, Valsalva maneuver and tilting table test have focused on the evoked response of autonomic nervous system. After an extensive literature survey, it has been observed that there is no substantial work that delves down to the details of HRV analysis using frequency domain method of HRV, in young and postmenopausal female medical professionals at the resting supine posture. Besides being a non-invasive study procedure, an important advantage of frequency domain analysis of HRV is that it utilizes spontaneous fluctuations in the heart rate to estimate autonomic functions.

The present study was conducted to evaluate the difference in HRV between young premenopausal women and postmenopausal women and to find out the possible role of contributing factors like serum estradiol level, serum triglycerides level and body fat percentage.

2. Material and Method

It was a comparative, cross-sectional, observational study. This study was carried out at Index Medical College Hospital and Research Centre, Indore (Madya Pradesh) after obtaining approval from the institutional ethics committee. All the young premenopausal and postmenopausal subjects were medical professionals working at the same medical institute. In this study total sample size was 80 having power of study 80% and significant p-value<0.05 as per online statistical software for sample size calculation. The study population was divided into two groups as per inclusion and exclusion criteria. One group included 40 young premenopausal women aged 18-24 years and the other group included 40 postmenopausal women aged 45-55 years who had attained menopause naturally at least 1 years ago. Data collection for proposed study was conducted within six months from January 2020 to July 2020. Any known case of acute or chronic illness, subjects taking any medicine that modulates autonomic nervous system; subjects taking any oral contraceptives pills or any hormonal replacement therapy were excluded from the study.

Patient preparation – Experimental protocol was maintained by explaining the study protocol in detail to the participants

of the study and written informed consent was taken from the participants. A thorough history was taken and general physical examinations were done to screen out the subjects for exclusion. After explaining exact experimental procedure, the procedures were performed at resting supine posture in our physiology research laboratory, in the morning time (room temperature was maintained at 20–25°C) in the fasting state. In premenopausal group, it was carried out only in the follicular phase of menstrual cycle. Subjects were refrained from caffeinated beverages for at least 12 hours prior to the experiment. The height and weight of all subjects were screened. After 15 min of rest, basal heart rate, basal blood pressure, and ECG were recorded at supine posture. The basal recording of blood pressure was done using sphygmomanometer by standard Riva-Rocci method. ECG recording was performed at resting supine posture using BIOPAC[®] MP150 system at sampling frequency of 500Hz. Software Acknowledge 4.2 was used for acquisition of HRV signals of all subjects. All the subjects were instructed to breathe quietly during the entire recording period with closed eyes and to avoid talking, moving limbs and body, coughing, and sleeping.

All the subjects underwent the different tests in following order –

- 1) Serum Estradiol Level: 5 ml venous blood sample was withdrawn from antecubital vein and serum was separated. This sample was stored at 2-8^oC. Further analysis and serum estradiol assay was performed by chemiluminescence immunoassay (CLIA) method. In premenopausal group, it was carried out only in the follicular phase of menstrual cycle.
- 2) Body Fat percentage – Includes the measurement of skin fold thickness at triceps, sub scapular and supra-iliac sites by skin fold caliper. The sum of three skin folds was used in age and gender specific equation to obtain an estimate of body fat percentage (Edwards K. D. G. et al.1962).
Body fat percentage = 0.29 SF*+ 3.9 (Weight in Kg.)
(SF*: sum of Skin Fold thickness)
- 3) Heart Rate Variability (HRV) – HRV was recorded by medical analyzer module (based on principle of impedance plethysmography) of NIVOMON, L&T and analysis of signal was done in frequency domain measures. In the frequency domain analysis, traditional Fast Fourier transform (FFT) based method was used to calculate the power spectrum for the HRV. Data were edited manually for artefacts and ectopic beats. Frequency domain measurement including Total Power, HFPower (0.15-0.40 Hz), LF Power (0.04-0.15Hz) and LF/HF ratio was calculated to assess sympathetic/parasympathetic modulation. Total Power and HFn. u. (normalized unites) are measures of parasympathetic activity and LF n. u. and LF/HF ratio are measures of the sympathetic activity.

Statistical analysis: The data was expressed as mean+ SD. Analysis was conducted by using Microsoft excel software, Microsoft Corporation USA, 2003. To find the statistical difference between young premenopausal and postmenopausal groups, we used Student's unpaired t-test in this study. p-value < 0.05 was considered statistically significant.

3. Results

Our study results showed that serum estradiol level was significantly higher (p-value< 0.001) in young premenopausal women (195.53+77.18 Pg/ml) as compared to postmenopausal women (19.04+7.87 Pg/ml) (Table 1).

We observed that serum triglyceride level was significantly higher (p-value< 0.001) in postmenopausal women (127.10+12.61 mg/dl) as compared to young premenopausal women (98.84+7.57mg/dl). Similarly, 'body fat percentage' was significantly higher (p-value< 0.001) in postmenopausal women (18.55+2.13 %) as compared to young premenopausal women (17.04+1.07 %) (Table 1).

In the present study 'Total Power' was significantly lower (p-value< 0.001) in postmenopausal women (917.89+647.67 ms²) as compared to young premenopausal women (4900.82+2876.29 ms²) (Table 2; Fig.1).

We observed that 'LF Power' was significantly higher (p-value< 0.001) in postmenopausal women (693.14+271.21 ms²) as compared to young premenopausal women (141.60+107.49 ms²), while, 'HF Power' was significantly higher (p-value< 0.001) in young premenopausal women (1186.65+783.73 ms²) as compared to postmenopausal women (134.97+109.85 ms²) (Table 2).

In the present study, 'LF/HF ratio' was significantly higher (p-value< 0.001) in postmenopausal women (8.95+2.47) as compared to young premenopausal women (0.22+0.14) (Table 2; Fig.2).

Table 1: Showing the various anthropometric data, general physical examination findings, vitals and related blood investigations

Parameters	Group (n=40)		p-value
	Young Premenopausal (Mean+SD)	Postmenopausal (Mean+SD)	
Age (Years)	20.70+2.69	50.67+3.42	
Height (cm)	158.66+2.57	156.97+2.61	
Weight (Kg)	56.42+3.84	62.34+6.27	
Systolic B. P. (mm of Hg)	121.15+4.79	127.60+7.35	
Diastolic B. P. (mm of Hg)	78.05+1.65	79.70+3.11	
Heart Rate (per minute)	74.45+3.69	76.25+4.21	
Serum Estradiol level (Pg/ml)	195.53+77.18	19.04+7.87	<0.001**
Serum Triglycerides level (mg/dl)	98.84+7.57	127.10+12.61	<0.001**
Body Fat Percentage (%)	17.04+1.07	18.55+2.13	<0.001**

Data expressed as mean+SD; *p<0.05 (Significant); **p<0.001 (Highly Significant).

Table 2: Showing the various frequency domain parameters of HRV between Young Premenopausal and Postmenopausal women group by Unpaired 't' Test

Parameters	Group (n=40)		p-value
	Young Premenopausal (Mean+SD)	Postmenopausal (Mean+SD)	
Total Power (ms ²)	4900.82+2876.29	917.89+647.67	<0.001**
LF Power (ms ²)	141.60+107.49	693.14+271.21	<0.001**
HF Power (ms ²)	1186.65+783.73	134.97+109.85	<0.001**
LF _{nu} (%)	21.24+10.21	71.76+14.57	<0.001**
HF _{nu} (%)	66.97+15.87	19.11+9.75	<0.001**
LF/HF ratio	0.22+0.14	8.95+2.47	<0.001**

Data expressed as mean+SD; *p<0.05 (Significant); **p<0.001 (Highly Significant); LF= Low Frequency, HF= High Frequency, and n. u. = normalized units.




Figure 1: Showing Power Spectral Density (PSD) in terms of 'Total Power' in Young Premenopausal and Postmenopausal women group.




Figure 2: Showing Power Spectral Density (PSD) in terms of 'LF/HF ratio' in Young Premenopausal and Postmenopausal women group.

4. Discussion

Menopause is a part of a women's natural ageing process when her ovaries produce lower level of the estradiol due to depletion of functional ovarian follicles and there is gradual transition from reproductive to non-reproductive phase of life. The median age of menopause is about 51 years. Bhat AN et al⁸ evaluated that the transitional changes are in the women's usual menstrual period and beginning of vasomotor symptoms, suggesting an alteration of cardiovascular reflexes. Sherwood A et al⁹ studied that menopause is associated with decreased 'Heart Rate Variability', which is due to increased sympathetic or reduced parasympathetic out flow to the heart. Massin Met al¹⁰ evaluated that HRV is a non-invasive index of neuronal activity of heart.

In the present study, we used spectral analysis of HRV while subjects were in relaxing and resting supine posture, since the daily physical activity level is considered as one of the potential confounders in the measurement of autonomic activity. Previous studies on comparison of HRV of premenopausal women using time domain indexes had not controlled for phases of menstrual cycle during subject recruitment, despite the possibility of ovarian hormonal influences on ANS function. In the present study, all premenopausal subjects were recruited for the study, during their follicular phase of menstruation cycle to find out the contribution of estrogen for the difference in HRV between the study groups.

Hence, on the basis of observations of our study, here it can be mentioned that the higher relative power of LF, increased ratio of LF/HF, and low relative power of HF components of HRV in postmenopausal women suggest shifting of cardiac autonomic balance towards sympathetic dominance in them. Such physiological changes may be confounded by age, declined serum estrogen level, increased serum triglycerides level, and increased body fat percentage as the status changes from premenopausal to postmenopausal. All these findings of our study are well comparable with the study results of Davy KP et al¹¹, and MoodithayaSetal².

5. Conclusion

It can be concluded from our present study that HRV is a good tool to measure the degree of autonomic modulation. Our results demonstrate that HRV status in postmenopausal women reflects cardiac autonomic disturbances. 'Total Power', and 'HF Power' is significantly lower, while 'LF Power' and 'LF/HF ratio' is significantly higher in postmenopausal women as compared to young premenopausal women, which suggests overall reduced parasympathetic drive and poor vasovagal balance in postmenopausal women. Further, decreased level of serum estradiol, increased serum triglycerides level and increased body fat percentage indicate sympathetic dominance in postmenopausal women that may lead to higher risk of cardiovascular morbidity and mortality. It is important to analyze and detect the variations in HRV between young premenopausal and postmenopausal female in order to have the insight of heart, so that affected women can be instructed for proper precautions and timely interventions.

Acknowledgement: None

Declarations

Source (s) of funding: None

Conflict of interest: None.

Ethical approval: This study was approved by the institutional ethics committee.

References

- [1] Johnson TR: Knowledge and Attitudes Regarding the Menstrual Cycle, Oral Contraceptives, and Sport Performance: The Conceptualization and Development of a Questionnaire for Athletic Coaches. Ph. D. Thesis, Florida State University, Florida (2008).
- [2] Moodithaya SS, Avadhany ST: Comparison of cardiac autonomic activity between pre and postmenopausal women using heart rate variability. *Indian J Physiol Pharmacol* 2009; 53 (3): 227-234.
- [3] Neves VF, Silva de Sa MF, Gallo L Jr, and Catai AM: Autonomic modulation of heart rate of young and postmenopausal women undergoing estrogen therapy. *Braz J Med Biol Res.* 2007; 40 (4): 491-499.
- [4] Liu CC, Kuo TB, Yang CC: Effect of estrogen on gender related autonomic difference in human. *Am. J. physiol heart cir. Physiol.* 2003; 285: 2188-2193.
- [5] Akiyoshi M, Kato K, Owa Y: Relationship between estrogen, vasomotor symptoms, and heart rate variability in climacteric women. *J Med Dent Science.* 2011; 58: 49-59.
- [6] Schillaci G, Verdccchia P, Borgioni C, Ciucci A, Porcellati C: Early cardiac changes after the menopause. *Hypertension* 1998; 32: 764-769.
- [7] Bannister R: Autonomic failure. A text book of clinical disorder of autonomic neuropathy. Oxford University Press. Oxford, New York. 1999.
- [8] Bhat AN, Sadhoo AK, Yograj S, Kaur G: Autonomic functions in postmenopausal women. *Jk Science* 2005; 7 (3): 135-139.
- [9] Sherwood A, Thurston R, Stenffen P, Blumenthal JA, Waugh RA: Blunted night time blood pressure dipping in postmenopausal women. *Am. J. Hypertens* 2001; 286: 749-754.
- [10] Massin M, Bernuth VG: Normal ranges of heart rate variability during infancy and childhood. *Pediatric cardiol.* 1997; 18: 297-302.
- [11] Davy KP, Miniclier N, Taylor JA, Stevenson ET, Seals DR: Elevated heart rate variability in physically active postmenopausal women: a cardioprotective effect. *Am. J. Physio.* 1996: 271: 455-460.