

Fixed Point Results for Parametric Metric Space

Arun Kumar Garg¹, Satyendra Ahirwar²

^{1, 2}Department of Mathematics, Madhyanchal University, Bhopal, M.P., India

¹gargarun1956[at]gmail.com

²Satyendraa07[at]gmail.com

Abstract: There are many generalization of metric space. Parametric metric space is the generalization of metric space too. Which was introduced and studied by Hussian (a new approach to metric space) in 2014. In present paper we prove two fixed point theorems based on injective mapping using contraction conditions.

Mathematics Subject Classification (MSC): 47H10, 54H25

Keywords: unique fixed point, contraction, parametric metric space, injective mapping

1. General Introduction

A metric on a nonempty set X is a mapping $d: X \times X \rightarrow [0, \infty]$ satisfying the following properties:

- (i) $d(x, y) = 0$ If and if $x = y$
- (ii) $d(x, y) = d(y, x)$
- (iii) $d(x, y) \leq d(y, z) + d(z, y)$; then the pair (X, d) is said to be a metric spaces.

The theory of metric spaces is the general theory which covers several branches of mathematical analysis, as real analysis, complex analysis, multidimensional calculus, etc. Due to which, existence and uniqueness of fixed points and common fixed points has become a subject of great concern. In the recent six decades many authors generalized the Banach contraction Principle by moderating the triangular inequality of a metric space as generalized metric space [see 2,5,7-8,14,22 and references therein], cone metric space [see 9 references therein], b metric space [see 2,3,4,6 references therein], cone b metric space [see 9,10,11,14-22 references therein], rectangular metric space [see 17 references therein], cone rectangular metric space [see 12,17,18 references therein], are some of the generalizations of metric space introduced by different authors in past few decades. Analogue Banach contraction principle, Kannan contraction principle, Ćirić contraction principle and lots of the existing fixed point theorems for various generalized contractions were proved in these generalized spaces.

Most of the generalization of metric space are Hausdorff topology but we can also find generalization of metric space which are not necessarily Hausdorff topology (see, ref. [13, 19, 22,]). Tarskian mathematician used non Hausdorff topology for programming language semantics used in computer science.

The purpose of this paper is to prove some fixed point theorems for contraction mapping in parametric metric spaces. An example is also given to distinguish our results.

Preliminaries: proceeding to our main result, let we furnish some definition, proposition, properties & lemmas needed in sequel.

1. Let X be a non empty set and $T_p: X \times X \times (0, \infty) \rightarrow (0, \infty)$ be a map on X such that $\forall x, y, z \in X$ and $t > 0$

(a) $T_p(x, y, t) = 0$ if and only if $x = y$

(b) $T_p(x, y, t) = T_p(y, x, t)$

(c) $T_p(x, y, t) \leq T_p(x, z, t) + T_p(z, y, t)$

Then T_p is called parametric metric and pair (X, d) is called parametric metric space.

(a). If $\log_{n \rightarrow \infty} (x_n, x, t) = 0 \Rightarrow \log_{n \rightarrow \infty} x_n = x$, for all $t > 0$ then sequence $\{x_n\}_{n=1}^{\infty}$ converges $x \in X$

(b). If $\log_{n \rightarrow \infty} (x_n, x_m, t) = 0$ for all $t > 0$ then sequence $\{x_n\}_{n=1}^{\infty}$ is called Cauchy sequence.

(d). If every Cauchy sequence is convergent, then parametric metric space (X, d) is a complete parametric metric space.

(e). Let (X, d) be a parametric metric space and $T: X \rightarrow X$ be a mapping, then We say T is a continuous mapping at p in X , if for any sequence $\{x_n\}_{n=1}^{\infty} \in X$ such that $\log_{n \rightarrow \infty} x_n = x \Rightarrow \log_{n \rightarrow \infty} T x_n = T x$.

2. Main Result

The objective of this paper is to prove some new fixed point theorems in parametric metric space. This paper is divided in two sections. In Section I and II we prove theorems on parametric metric spaces.

Section I:

Theorem 2: Let (X, d) be a complete parametric metric space and $T_p: X \rightarrow X$ be an injective mapping satisfying the condition

$$(2.1) d(T_p x, T_p y, t) \leq$$

$$\alpha d(x, y, t) + \beta \left(\frac{d(x, T_p x, t) \cdot d(y, T_p y, t)}{d\{x, y\} + d(x, T_p x, t)} \right) + \gamma \left(\frac{d(x, T_p x, t) \cdot d(x, T_p y, t)}{d\{x, y\} + d(x, T_p x, t) + 2d(y, T_p y, t)} \right)$$

$\forall t \in [0, 1]; \alpha, \beta, \gamma > 0; x, y \in X$ & $x \neq y$ have a fixed point if $\alpha + \frac{\beta}{2} + \frac{\gamma}{2} < 1$ and moreover a unique fixed point if $\alpha + \gamma < 1$.

Proof: Let $ssx_0 \in X$, Define iterative sequence $\{x_n\}_{n=1}^{\infty}$ follows: $T_p x_n = x_{n+1}$ for $n = 1, 2, 3, \dots$. If for some n , $T_p x_n = x_n$, then x_n is the fixed point. Otherwise $T_p x_n \neq x_n$, using inequality (2.1)

$$d(x_{n+1}, x_{n+2}, t) = d(T_p x_n, T_p x_{n+1}, t)$$

$$\begin{aligned}
&\leq \alpha d(x_n, x_{n+1}, t) + \beta \left(\frac{d(x_n, T_p x_n, t) \cdot d(x_{n+1}, T_p x_{n+1}, t)}{d(x_n, x_{n+1}, t) + d(x_n, T_p x_n, t)} \right) \\
&+ \gamma \left(\frac{d(x_n, T_p x_n, t) \cdot d(x_n, T_p x_{n+1}, t)}{d(x_n, x_{n+1}, t) + d(x_n, x_{n+1}, t) + 2d(x_{n+1}, x_{n+2}, t)} \right) \\
&d(x_{n+1}, x_{n+2}, t) \\
&\leq \alpha d(x_n, x_{n+1}, t) + \beta \left(\frac{d(x_n, x_{n+1}, t) \cdot d(x_{n+1}, x_{n+2}, t)}{d(x_n, x_{n+1}, t) + d(x_n, x_{n+1}, t)} \right) \\
&+ \gamma \left(\frac{d(x_n, x_{n+1}, t) \cdot d(x_n, x_{n+2}, t)}{d(x_n, x_{n+1}, t) + d(x_n, x_{n+1}, t) + 2 \cdot d(x_{n+1}, x_{n+2}, t)} \right) \\
&d(x_{n+1}, x_{n+2}, t) \\
&\leq \alpha d(x_n, x_{n+1}, t) + \beta \left(\frac{d(x_n, x_{n+1}, t) \cdot d(x_{n+1}, x_{n+2}, t)}{2 \cdot d(x_n, x_{n+1}, t)} \right) \\
&+ \gamma \left(\frac{d(x_n, x_{n+1}, t) \cdot d(x_n, x_{n+2}, t)}{2 \cdot d(x_n, x_{n+1}, t) + 2 d(x_{n+1}, x_{n+2}, t)} \right) \\
&(x_{n+1}, x_{n+2}, t) \leq \alpha d(x_n, x_{n+1}, t) + \beta \left(\frac{d(x_{n+1}, x_{n+2}, t)}{2} \right) \\
&+ \gamma \left(\frac{d(x_n, x_{n+1}, t)}{2} \right) \\
&\left(1 - \frac{\beta}{2}\right) d(x_{n+1}, x_{n+2}, t) \leq \left(\alpha + \frac{\gamma}{2}\right) d(x_n, x_{n+1}, t) \\
&d(x_{n+1}, x_{n+2}, t) \leq \frac{\left(\alpha + \frac{\gamma}{2}\right)}{\left(1 - \frac{\beta}{2}\right)} d(x_n, x_{n+1}, t) \\
&d(x_{n+1}, x_{n+2}, t) \leq k d(x_n, x_{n+1}, t) \\
&\forall t \in [0,1] \text{ and } k = \frac{\left(\alpha + \frac{\gamma}{2}\right)}{\left(1 - \frac{\beta}{2}\right)} < 1 \Rightarrow \alpha + \frac{\beta}{2} + \frac{\gamma}{2} < 1.
\end{aligned}$$

Therefore by successive iteration

$$d(x_{n+1}, x_{n+2}, t) \leq k^n d(x_0, x_1, t)$$

As we know if $\{x_n\}_{n \rightarrow \infty}$ be a sequence in parametric space (X, d) such that

$$d(x_{n+1}, x_{n+2}, t) \leq k d(x_n, x_{n+1}, t) \forall t \in [0,1] \& n =$$

$$\begin{aligned}
(3.1) (T_p x, T_p y, t) &\leq \alpha \text{Max}\{d(x, y, t), \frac{d(x, T_p x, t) d(y, T_p y, t)}{d(x, y, t)}, \frac{d(x, T_p y, t) d(y, T_p x, t)}{d(x, y, t)}, \frac{d(x, T_p x, t) d(x, T_p y, t)}{2d(x, y, t)}\} \\
&\forall t \in [0,1]; \alpha > 0; x, y \in X \& x \neq y \text{ and } \alpha \in [0,1], \text{ then } T_p \text{ has a unique fixed point.}
\end{aligned}$$

Proof: Let $x_0 \in X$, Define iterative sequence $\{x_n\}_{n=1}^{\infty}$ follows: $T_p x_n = x_{n+1}$ for $n = 1, 2, 3, \dots$. If for some n $T_p x_n = x_n$, then x_n is the fixed point. Otherwise $T_p x_n \neq x_n$, using inequality (3.1)

$$\begin{aligned}
&\leq \alpha \text{Max}\{d(x_n, x_{n+1}, t), \frac{d(x_n, T_p x_n, t) d(x_{n+1}, T_p x_{n+1}, t)}{d(x_n, x_{n+1}, t)}, \\
&\frac{d(x_n, T_p x_{n+1}, t) d(x_{n+1}, T_p x_n, t)}{d(x_n, x_{n+1}, t)}, \frac{d(x_n, T_p x_n, t) d(x_n, T_p x_{n+1}, t)}{2d(x_n, x_{n+1}, t)}\} \\
&\leq \alpha \text{Max}\{d(x_n, x_{n+1}, t), \frac{d(x_n, x_{n+1}, t) d(x_{n+1}, x_{n+2}, t)}{d(x_n, x_{n+1}, t)}, \\
&\frac{d(x_n, x_{n+2}, t) d(x_{n+1}, x_{n+1}, t)}{d(x_n, x_{n+1}, t)}, \frac{d(x_n, x_{n+1}, t) d(x_n, x_{n+2}, t)}{2d(x_n, x_{n+1}, t)}\} \\
&\leq \alpha \text{Max}\{d(x_n, x_{n+1}, t), \frac{d(x_n, x_{n+1}, t) d(x_{n+1}, x_{n+2}, t)}{d(x_n, x_{n+1}, t)}, 0, \frac{d(x_n, x_{n+1}, t) \{d(x_n, x_{n+1}, t) + d(x_{n+1}, x_{n+2}, t)\}}{2d(x_n, x_{n+1}, t)}\} \\
&d(x_{n+1}, x_{n+2}, t) \leq \alpha \text{Max}\{(d(x_n, x_{n+1}, t), d(x_{n+1}, x_{n+2}, t), 0, \frac{d(x_n, x_{n+1}, t) + d(x_{n+1}, x_{n+2}, t)}{2})\}
\end{aligned}$$

$$\Rightarrow d(x_{n+1}, x_{n+2}, t) \leq \alpha d(x_n, x_{n+1}, t)$$

Therefore by successive iteration

$$d(x_{n+1}, x_{n+2}, t) \leq \alpha^n d(x_0, x_1, t)$$

Volume 11 Issue 7, July 2022

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

1,2,3, ... then $\{x_n\}_{n \rightarrow \infty}$ is a Cauchy sequence in (X, d) . Since (X, d) is a complete parametric space; $\{x_n\}_{n \rightarrow \infty}$ converges. Let $x^* \in X$, then $\lim_{n \rightarrow \infty} x_n \rightarrow x^*$. Again T_p is continuous, therefore

$$T_p x^* = T_p (\lim_{n \rightarrow \infty} x_n) = \lim_{n \rightarrow \infty} T_p x_n = x^* \Rightarrow T_p x^* = x^*$$

Implies T_p has a fixed point $T_p x^* = x^*$ in X .

Now we will show that x^* is unique. for that suppose y^* is another fixed point therefore $T_p y^* = y^*$. Therefore by inequality (2.1) we have

$$\begin{aligned}
&d(T_p x^*, T_p y^*, t) \\
&\leq \alpha d(x^*, y^*, t) + \beta \left(\frac{d(x^* T_p x^*, t) \cdot d(y^*, T_p y^*, t)}{d(x^*, y^*, t) + d(x^*, T_p x^*, t)} \right) \\
&+ \gamma \left(\frac{d(x^*, T_p x^*, t) \cdot d(x^*, T_p y^*, t)}{d(x^*, y^*, t) + d(x^*, T_p x^*, t) + 2d(y^*, T_p y^*, t)} \right) \\
&d(x^*, y^*, t) \\
&\leq \alpha d(x^*, y^*, t) + \beta \left(\frac{d(x^*, x^*, t) \cdot d(y^*, y^*, t)}{d(x^*, y^*, t) + d(x^*, x^*, t)} \right) \\
&+ \gamma \left(\frac{d(x^*, x^*, t) \cdot d(x^*, y^*, t)}{d(x^*, y^*, t) + d(x^*, x^*, t) + 2d(y^*, y^*, t)} \right)
\end{aligned}$$

$$d(x^*, y^*, t) \leq \alpha d(x^*, y^*, t)$$

$$d(x^*, y^*, t) \leq \alpha d(x^*, y^*, t)$$

$$\Rightarrow (1 - \alpha) d(x^*, y^*, t) \leq 0$$

$\Rightarrow d(x^*, y^*, t) = 0$ Since $\alpha < 1 \Rightarrow x^* = y^*$. Hence T_p has a unique point.

Section II:

Theorems: Let (X, T_p) be a complete parametric b- metric space and $T_p: X \rightarrow X$ be a orbitally continuous self-map satisfying the condition

$$d(x_{n+1}, x_{n+2}, t) = d(T_p x_n, T_p x_{n+1}, t)$$

$$d(x_{n+1}, x_{n+2}, t) \leq \alpha^n d(x_0, x_1, t)$$

As we know if $\{x_n\}_{n \rightarrow \infty}$ be a sequence in parametric space (X, d) such that $d(x_{n+1}, x_{n+2}, t) \leq \alpha^n d(x_0, x_1, t)$

$\forall t \in [0, 1]$ & $n = 1, 2, 3, \dots$ then $\{x_n\}_{n \rightarrow \infty}$ is a Cauchy sequence in (X, d) . Since (X, d) is a complete parametric space; $\{x_n\}_{n \rightarrow \infty}$ converges. Let $x^* \in X$, then $\lim_{n \rightarrow \infty} x_n \rightarrow x^*$. Again T_p is continuous, therefore

$$T_p x^* = T_p(\lim_{n \rightarrow \infty} x_n) = \lim_{n \rightarrow \infty} T_p x_n = x^* \Rightarrow T_p x^* = x^*$$

$$d(T_p x^*, T_p y^*, t) \leq \alpha \max\{d(x^*, y^*, t), \frac{d(x^*, T_p x^*, t)d(y^*, T_p y^*, t)}{d(x^*, y^*, t)}, \frac{d(x^*, T_p y^*, t)d(y^*, T_p x^*, t)}{d(x^*, y^*, t)}, \frac{d(x^*, T_p x^*, t)d(x^*, T_p y^*, t)}{2d(x^*, y^*, t)}, \frac{d(x^*, x^*, t)d(y^*, y^*, t)}{d(x^*, y^*, t)}, \frac{d(x^*, y^*, t)d(y^*, x^*, t)}{d(x^*, y^*, t)}, \frac{d(x^*, x^*, t)d(x^*, y^*, t)}{2d(x^*, y^*, t)}\}$$

$$d(T_p x^*, T_p y^*, t) \leq \alpha \max\{d(x^*, y^*, t), 0, d(x^*, y^*, t), 0\}$$

$$d(x^*, y^*, t) \leq \alpha d(x^*, y^*, t)$$

$$\Rightarrow (1 - \alpha)d(x^*, y^*, t) \leq 0$$

$\Rightarrow d(x^*, y^*, t) = 0$ Since $\alpha > 1 \Rightarrow x^* = y^*$. Hence T_p has a unique point.

Implies T_p has a fixed point $T_p x^* = x^*$ in X .

Now we will show that x^* is unique. for that suppose y^* is another fixed point therefore $T_p y^* = y^*$. Therefore by inequality (3.1) we have

References

- [1] Achari J. (1983): "Fixed point theorems for a class of mappings on non-Archimedean probabilistic metric spaces, *Mathematica*, 25, 5-9,
- [2] Alamgir Khan M, Sumitra (2008) "Common fixed point theorem in 2 N. A. Menger PM-space for R-weakly commuting maps of type (P)", *Novi Sad J. Math.* (38)(2), 145-152.
- [3] Alamgir Khan M Sumitra: "A common fixed point theorem in non-Archimedean Menger PM-space", *Novi Sad J. Math.*, Vol. 39(2), 81-87.2009
- [4] Alamgir Khan M and Sumitra: "Common fixed point theorems in non-Archidean Menge PM-space", 5(1), 1-13, 2010
- [5] Bota, M, Molnar, A and Varga, C: "On ekeland's variational principle in b-metric spaces", *Fixed Point Theory* no. 2, 21-28, 12 (2011).
- [6] Chugh, Rkumar, V, and Kadian, T:: Some fixed point theorems for multivalued mappings in generalized b-metric spaces, *International Journal of Mathematical Archive*, no. 3, 1198-1210, 3 (2012).
- [7] Czerwik, S: "Contraction mappings in b-metric spaces", *Acta Mathematica et Informatica Universitatis Ostraviensis*, 5-11. 1 (1993).
- [8] Czerwik S., Nonlinear set-valued contraction mappings in b-metric spaces, *Atti. Sem. Mat. Fis. Univ. Modena* 46 (1998), 263-276.
- [9] J. R. Morales and E. Rojas: "Cone metric spaces and fixed point theorems of T -Kannan contractive mappings" *arXiv: 0907. 3949v2 [math. FA]* 26 Oct. 2009
- [10] Kir, M, Kiziltunc, H: "On some well known fixed point theorems in b-metric spaces", *Turkish Journal of Analysis and Number Theory*, 1, no. 1, 13-16. (2013).
- [11] Khan, M.S: "A fixed point theorem for metric spaces", *Rent. Inst. Mat. Univ. Trieste*, Vol. III, Fasc. 10 1 - 4. (1976).
- [12] L.G. Haug and X. Zhang: "Cone metric spaces and fixed point of contractive mapping", *J. Math. Anal.*
- [13] M. Jleli, B. Samet, *The Kannan's fixed point theorem in a cone rectangular metric space*, *J. Nonlinear Sci. Appl.* 2 (3) (2009), 161-167.2009
- [14] N. Shioji, T. Suzuki and W. Takahashi: "Contraction mappings, Kannan mappings and metric completeness" *Proceedings of the American Mathematical Society*, vol. 126, No. 10, 3117-3124, October 1998
- [15] R. George, Hossam A. Nabwey, K. P. Reshma and R. Rajagopalan: "generalized cone b-metric spaces and contraction principles" *MATEMATIKI VESNIK*, 67, 4 (2015), 246-257, December 2015
- [16] R. George, B. Fisher, "Some generalized results of fixed points in cone b-metric space", *Math. Moravica* 17 (2) (2013), 39-50., 2013
- [17] S. Rezapour and R. Hambarani: "Some notes on the paper Cone metric spaces and fixed point Theorems of contractive mappings" *J. Math. Anal. Appl.*, 345, (2), 719-724, 2008
- [18] Shi, L, Xu, S: "Common fixed point theorems for two weakly compatible self-mappings in cone b-metric spaces", *Fixed Point Theory and Applications*, no. 120. 2013
- [19] S. Moradi: "Kannan fixed point theorem on complete metric spaces and on generalized metric spaces depend on another function" *arXiv: 0903. 1577v1 [math. FA]*
- [20] T. Suzuki, "Generalized metric spaces do not have compatible topology", *Abstract Appl. Anal.* 2014, Art. ID 458098, 5 pages.
- [21] T. Van An, N. Van Dung, Z. Kadelburg, S. Radenović, "Various generalizations of metric spaces and fixed point theorems", *Rev. Real Acad. Cienc. Exac. Fis. Nat., Ser. A, Mat.* 109, 1 (2015), 175-198.
- [22] Z. Kadelburg, S. Radenović, Fixed point results in generalized metric spaces without Hausdorff property, *Math. Sci.* 8:125 (2014).