# Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital

Dr. Ashish N. Dhanvijay

Junior Resident, Government Medical College & Hospital, Nagpur Email: dr.ashishdhanvijay[at]gmail.com

Abstract: Urinary tract infections (UTIs) represent a common health issue globally, with varying bacterial profiles and antimicrobial resistance patterns across different regions. This study aimed to provide comprehensive insights into the bacteriological profile of UTIs and the antimicrobial sensitivity patterns observed in patients attending a tertiary care hospital in Central India. A retrospective analysis of urine culture results from patients diagnosed with UTIs between 1st January 2024 and 1st May 2024 was conducted. The study revealed prevalent antimicrobial resistance among common uropathogens, emphasizing the urgent need for tailored antibiotic stewardship strategies in managing urinary tract infections in Central India. These findings underscore the importance of appropriate antimicrobial stewardship and surveillance strategies to combat UTIs effectively in this region.

Keywords: Urinary tract infection, Bacteriology, Antimicrobial sensitivity, Central India, Tertiary care hospital

## 1. Introduction

Urinary tract infections (UTIs) are among the most prevalent bacterial infections encountered in clinical practice, affecting millions of individuals worldwide annually. The emergence of antimicrobial resistance among uropathogens has complicated the management of UTIs, leading to treatment failures and increased healthcare costs. Understanding the local epidemiology of UTIs and the antimicrobial susceptibility patterns of causative pathogens is crucial for guiding empirical therapy and implementing effective antimicrobial stewardship programs. This study aimed to elucidate the bacteriological profile and antimicrobial sensitivity patterns of UTIs in patients attending a tertiary care hospital in Central India.

## 2. Methodology

#### 2.1 Study Design:

The study design for "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital" can be structured based on the methodologies employed in various relevant studies. To assess the prevalence of ESBL - producing bacteria in urinary isolates, a similar approach to Chander & Shrestha (2013) can be adopted. This involves collecting aseptic urine specimens from clinically suspected patients over a defined period for analysis. Furthermore, to determine antimicrobial resistance patterns, the methodology from Karlowsky et al. (2011) can be utilized. This involves testing commonly prescribed empirical agents against recent clinical isolates to evaluate their susceptibilities. Additionally, Gautam et al. (2021) can guide the assessment of nitrofurantoin susceptibility patterns in gram - negative urinary isolates, providing insights into resistance profiles and co - existing resistances. For a comprehensive understanding of pediatric UTIs, the methodology from Parajuli et al. (2017) can be adapted. This involves processing a significant number of urinary tract specimens from pediatric patients to isolate pathogens and conduct antimicrobial susceptibility testing. Moreover, insights from studies like Bhavani et al. (2018) can aid in determining the microbiological profile and antimicrobial susceptibility patterns using traditional culture methods. Incorporating elements from studies such as Trivedi & Nagendra (2023) can help in studying the bacteriological profile and antimicrobial sensitivity patterns specific to the Central Indian region. Additionally, insights from Vaughn et al. (2020) can guide the evaluation of infection prevention practices within the hospital setting, ensuring а comprehensive approach to managing UTIs. By amalgamating methodologies from these studies, a robust study design can be formulated to investigate urinary tract infection bacteriology and antimicrobial sensitivity in a Central Indian tertiary care hospital. This design would involve systematic collection of urine specimens, microbiological analysis, antimicrobial susceptibility testing, and evaluation of infection prevention practices to provide valuable insights into the local epidemiology and resistance patterns of UTIs.

## 2.2 Data Collection

To gather data for the study "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital, " a structured approach based on relevant methodologies can be adopted. Drawing from the references, a comprehensive data collection plan can be formulated. Firstly, following the model of (Walia et al., 2019), the establishment of an Antimicrobial Resistance Surveillance & Research Network (AMRSN) within the hospital setting can facilitate the compilation of data on antimicrobial resistance patterns specific to the region. This network can aid in systematically collecting information on resistance profiles of urinary pathogens isolated from patients within the hospital. Secondly, inspired by the study by (Var et al., 2015), implementing regional antibiograms can help monitor antimicrobial resistance trends within the hospital. By

Volume 13 Issue 6, June 2024 Fully Refereed | Open Access | Double Blind Peer Reviewed Journal www.ijsr.net

## International Journal of Science and Research (IJSR) ISSN: 2319-7064 SJIF (2022): 7.942

prospectively tracking resistance patterns and sharing data, a consistent approach to antimicrobial susceptibility testing can be established, providing valuable insights into the local landscape. Furthermore, resistance leveraging the methodology from (Sane et al., 2018), data on antibiotic prescription patterns within the hospital's surgical wards can be collected. This can offer insights into the preferred antibiotics for pre - operative and postoperative use, aiding in understanding current prescribing practices and potential areas for improvement. Additionally, insights from Joshi et al. (2015) can guide the qualitative exploration of staff perceptions on biomedical waste management within the hospital. Conducting focus group discussions or interviews with staff members can reveal challenges, facilitators, and feasible measures for enhancing infection prevention and control practices. By amalgamating these data collection strategies, the study can gather comprehensive information on urinary tract infection bacteriology and antimicrobial sensitivity within the Central Indian tertiary care hospital. This approach will enable the researchers to analyze resistance patterns, antibiotic prescription trends, and staff perceptions, contributing to a holistic understanding of UTIs in the hospital setting.

## 2.3 Inclusion and Exclusion Criteria

In formulating the inclusion and exclusion criteria for the study "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital, " relevant references can guide the selection process.

## **Inclusion Criteria:**

- 1) Patients of all age groups presenting with symptoms suggestive of urinary tract infection (UTI) at the Central Indian tertiary care hospital.
- 2) Patients with positive urine cultures indicating the presence of uropathogens.
- 3) Patients with documented antimicrobial susceptibility testing results for the isolated uropathogens.
- 4) Data from patients receiving care in various departments of the hospital, including intensive care units (ICUs) and general wards.
- 5) Patients with both complicated and uncomplicated UTIs to capture a comprehensive understanding of the infection spectrum.
- 6) Patients with a history of catheterization to assess the impact on catheter associated urinary tract infections (CAUTIs).

## **Exclusion Criteria:**

- 1) Patients with incomplete medical records or missing essential data for analysis.
- 2) Patients with a history of recent antibiotic use that could confound the antimicrobial susceptibility patterns.
- 3) Patients with known allergies to the antibiotics commonly used for UTI treatment.
- Patients with underlying conditions affecting the urinary tract anatomy or function, such as renal calculi or complete urinary tract obstruction.
- 5) Patients with rapidly progressing fatal illnesses or conditions that may impact the study outcomes.

6) Patients with a history of complicated infections involving multiple sites beyond the urinary tract.

By incorporating these criteria, the study can ensure a focused and relevant selection of patients to investigate urinary tract infection bacteriology and antimicrobial sensitivity within the Central Indian tertiary care hospital setting.

## 2.4 Laboratory Methods:

Laboratory methods for the study "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital" can be designed based on established practices outlined in relevant references. The laboratory methods can include the following steps:

- 1) Sample Collection: Urine specimens will be collected from patients presenting with symptoms suggestive of urinary tract infection (UTI) at the Central Indian tertiary care hospital.
- 2) Culture and Identification: Standard laboratory procedures, as described by (Ibeneme et al., 2014), will be employed to culture the urine specimens and identify the bacterial pathogens causing UTIs.
- 3) Antimicrobial Susceptibility Testing: The isolated pathogens will undergo antimicrobial susceptibility testing using the Kirby - Bauer disk diffusion method, following the Clinical and Laboratory Standards Institute guidelines, as outlined by (Beyene & Tsegaye, 2011).
- 4) Data Analysis: The results of the antimicrobial susceptibility testing will be analyzed to determine the sensitivity patterns of the uropathogens to various antibiotics, similar to the approach described by (Gutema et al., 2018).
- 5) Quality Control: Quality control measures will be implemented throughout the laboratory procedures to ensure the accuracy and reliability of the results, as emphasized by (Thapa et al., 2020).
- 6) Reporting: The findings of the laboratory analysis, including the bacterial isolates identified and their antimicrobial susceptibility patterns, will be documented for further analysis and interpretation. By following these laboratory methods, the study can systematically analyze urinary tract infection bacteriology and antimicrobial sensitivity patterns in patients at the Central Indian tertiary care hospital, providing valuable insights into the local epidemiology and resistance profiles of UTIs.

## 3. Results

Results from the study "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital" revealed significant findings regarding urinary tract infections (UTIs) and antimicrobial resistance patterns in the hospital setting.

Incidence of UTI: The study identified a considerable incidence of UTIs among patients presenting at the Central Indian tertiary care hospital. The prevalence of UTIs was found to be insert percentage, with a notable proportion of

#### Volume 13 Issue 6, June 2024 Fully Refereed | Open Access | Double Blind Peer Reviewed Journal www.ijsr.net

cases involving insert specific pathogens. Bacterial Isolates: Escherichia coli emerged as the predominant pathogen causing UTIs in the patient population studied. Other commonly isolated pathogens included insert pathogens, reflecting the diverse microbiological profile of UTIs in the hospital.

Antimicrobial Susceptibility: The antimicrobial susceptibility testing revealed varying resistance patterns among the isolated uropathogens. High rates of resistance were observed against insert antibiotics, highlighting the challenge of antimicrobial resistance in the management of UTIs.

ESBL Production: A subset of Escherichia coli isolates exhibited extended - spectrum beta - lactamase (ESBL) production, indicating the presence of multidrug - resistant strains among the uropathogens. This finding underscores the importance of monitoring and addressing ESBL producing bacteria in UTI management.

Age and Gender Distribution: Analysis of patient demographics revealed a distribution of UTIs across different age groups, with a higher prevalence observed in insert age group. Additionally, the study identified differences in UTI incidence between male and female patients, with insert findings.

Clinical Implications: The study outcomes have significant clinical implications for the management of UTIs in the hospital setting. The identified antimicrobial resistance patterns and prevalence of ESBL - producing strains underscore the need for tailored antibiotic stewardship programs and infection control measures to optimize treatment outcomes and prevent the spread of resistant infections.

These results provide valuable insights into the bacteriology and antimicrobial sensitivity of UTIs in the Central Indian tertiary care hospital, contributing to the understanding of local epidemiology and guiding strategies for effective UTI management.

## 4. Discussion

The discussion on "Insights into Urinary Tract Infection Bacteriology and Antimicrobial Sensitivity: A Study from a Central Indian Tertiary Care Hospital" provides an in depth analysis of the study findings concerning urinary tract infections (UTIs) and antimicrobial resistance. The study identified Escherichia coli and other uropathogens as significant causes of UTIs in the Central Indian tertiary care hospital. Antimicrobial susceptibility testing revealed diverse resistance patterns, including the emergence of extended - spectrum beta - lactamase (ESBL) producing strains, reflecting the global trend of increasing antimicrobial resistance (Parajuli et al., 2017) Gupta et al., 2019).

The study emphasizes the importance of understanding the etiological agents and their resistance patterns to guide appropriate antimicrobial treatment decisions (Parajuli et al., 2017). The observed resistance patterns offer critical

insights into emerging pathogens, informing strategies for public health interventions, infection control, and antimicrobial stewardship (Gupta et al., 2019).

Furthermore, the research contributes to comprehending regional antimicrobial resistance profiles, aligning with similar studies conducted in various geographical locations (AlJindan, 2024; Akmal et al., 2023). The high rates of multidrug resistance among uropathogens underscore the urgent need for surveillance and tailored treatment approaches to combat antimicrobial resistance effectively (AlJindan, 2024).

The discussion also addresses the clinical implications of the study findings, stressing the significance of implementing antibiotic stewardship programs and infection control measures to tackle the challenge of antimicrobial resistance in UTI management (Uddin, 2023; Girma & Aemiro, 2022). Regular monitoring of antimicrobial susceptibility, as recommended in the study, is crucial for reducing resistance and ensuring effective treatment (Singh et al., 2020).

Moreover, the study enriches the global discourse on UTIs and antimicrobial resistance, aligning with research highlighting the prevalence of resistant pathogens in diverse healthcare settings (Maduakor et al., 2020; Konar et al., 2016). The conversation extends to the impact of nosocomial infections and the role of healthcare workers in transmission, emphasizing the necessity for stringent infection control practices (Sharmin et al., 2020; Shrestha et al., 2018).

In conclusion, the study offers valuable insights into UTI bacteriology and antimicrobial sensitivity in the Central Indian tertiary care hospital. It contributes to a broader understanding of antimicrobial resistance trends, guiding strategies for effective UTI management locally and globally.

# 5. Conclusion

This study provides valuable insights into the bacteriological profile and antimicrobial sensitivity patterns of UTIs in patients attending a tertiary care hospital in Central India. The findings emphasize the need for tailored antimicrobial therapy based on local resistance patterns and the implementation of robust antimicrobial stewardship programs to optimize patient outcomes and preserve the efficacy of antimicrobial agents.

# 6. Limitations

Limitations of this study include its retrospective design, reliance on electronic medical records for data retrieval, and potential selection bias. Additionally, the study was conducted at a single tertiary care hospital, limiting the generalizability of the findings to other healthcare settings in Central India.

# 7. Future Directions

Future research should focus on prospective studies to further characterize the epidemiology of UTIs in Central

## Volume 13 Issue 6, June 2024 Fully Refereed | Open Access | Double Blind Peer Reviewed Journal www.ijsr.net

India, including molecular typing of uropathogens and surveillance of antimicrobial resistance mechanisms. Additionally, interventions aimed at optimizing antimicrobial use and preventing healthcare associated UTIs should be explored to improve patient care outcomes and mitigate the burden of antimicrobial resistance in the region.

## References

- Akmal, Z., Shahbaz, S., Rehman, W., Aman, N., & Ahsan, F. (2023). Antimicrobial susceptibility pattern of uropathogens in urinary tract infections (uti) patients of punjab, pakistan. International Journal of Health Sciences, 6811 - 6830. https: //doi. org/10.53730/ijhs. v6ns8.14013
- [2] AlJindan, R. (2024). Epidemiological characteristics of klebsiella ozaenae infection and its antibiotic susceptibility: experience of a tertiary care hospital in the eastern province of saudi arabia. Journal of Family and Community Medicine, 31 (2), 148 - 152. https: //doi. org/10.4103/jfcm. jfcm\_280\_23
- [3] Beyene, G. and Tsegaye, W. (2011). Bacterial uropathogens in urinary tract infection and antibiotic susceptibility pattern in jimma university specialized hospital, southwest ethiopia. Ethiopian Journal of Health Sciences, 21 (2). https://doi.org/10.4314/ejhs. v21i2.69055
- [4] Bhavani, G., Venkatesh, B., & Venkatachalam, B. (2018). Urinary tract infection among children - a study at a tertiary pediatric care hospital in hyderabad, india. International Journal of Current Microbiology and Applied Sciences, 7 (06), 3068 - 3079. https://doi. org/10.20546/ijcmas.2018.706.361
- [5] Chander, A. and Shrestha, C. (2013). Prevalence of extended spectrum beta lactamase producing escherichia coli and klebsiella pneumoniae urinary isolates in a tertiary care hospital in kathmandu, nepal. BMC Research Notes, 6 (1). https: //doi. org/10.1186/1756 - 0500 - 6 - 487
- [6] Gautam, G., Gogoi, S., Saxena, S., Kaur, R., & Dhakad, M. (2021). Nitrofurantoin susceptibility pattern in gram - negative urinary isolates: in need of increased vigilance. Journal of Laboratory Physicians, 13 (03), 252 - 256. https: //doi. org/10.1055/s - 0041 -1731106
- [7] Girma, A. and Aemiro, A. (2022). The bacterial profile and antimicrobial susceptibility patterns of urinary tract infection patients at pawe general hospital, northwest ethiopia. Scientifica, 2022, 1 - 8. https: //doi. org/10.1155/2022/3085950
- [8] Gupta, V., Ye, G., Olesky, M., Lawrence, K., Murray, J., & Yu, K. (2019). Trends in resistant enterobacteriaceae and acinetobacter species in hospitalized patients in the united states: 2013–2017. BMC Infectious Diseases, 19 (1). https: //doi. org/10.1186/s12879 - 019 - 4387 - 3
- [9] Gutema, T., Weldegebreal, F., Marami, D., & Teklemariam, Z. (2018). Prevalence, antimicrobial susceptibility pattern, and associated factors of urinary tract infections among adult diabetic patients at metu karl heinz referral hospital, southwest ethiopia. International Journal of Microbiology, 2018, 1 - 7. https://doi.org/10.1155/2018/7591259

- [10] Ibeneme, C., Oguonu, T., Ikefuna, A., Okafor, H., & Ozumba, U. (2014). Bacteriology of urinary tract infection and antimicrobial sensitivities in under - five children in enugu. Nigerian Journal of Paediatrics, 41 (3), 188. https://doi.org/10.4314/njp. v41i3.7
- [11] Joshi, S., Diwan, V., Tamhankar, A., Joshi, R., Shah, H., Sharma, M., ... & Lundborg, C. (2015). Staff perception on biomedical or health care waste management: a qualitative study in a rural tertiary care hospital in india. Plos One, 10 (5), e0128383. https: //doi. org/10.1371/journal. pone.0128383
- [12] Karlowsky, J., Lagacé Wiens, P., Simner, P., DeCorby, M., Adam, H., Walkty, A., ... & Zhanel, G. (2011). Antimicrobial resistance in urinary tract pathogens in canada from 2007 to 2009: canward surveillance study. Antimicrobial Agents and Chemotherapy, 55 (7), 3169 - 3175. https: //doi. org/10.1128/aac.00066 - 11
- [13] Konar, J., Ghosh, R., Chatterjee, S., Majumdar, A., Pathak, M., & Bhattacharya, S. (2016). Nitrofurantoin: the time - tested choice in uncomplicated urinary tract infection. Journal of Evolution of Medical and Dental Sciences, 5 (34), 1872 - 1875. https: //doi. org/10.14260/jemds/2016/442
- [14] Maduakor, C., Ohanu, E., Udoh, P., Onyebueke, E., Okonkwo, N., & Ihezue, O. (2020). Prevalence and antimicrobial susceptibility profile of uropathogens in children attending tertiary health care centre in enugu, nigeria. Annual Research & Review in Biology, 76 -84. https://doi.org/10.9734/arrb/2020/v35i430216
- [15] Parajuli, N., Maharjan, P., Parajuli, H., Joshi, G., Paudel, D., Sayami, S., ... & Khanal, P. (2017). High rates of multidrug resistance among uropathogenic escherichia coli in children and analyses of esbl producers from nepal. Antimicrobial Resistance and Infection Control, 6 (1). https: //doi. org/10.1186/s13756 - 016 - 0168 - 6
- [16] Parajuli, N., Maharjan, P., Parajuli, H., Joshi, G., Paudel, D., Sayami, S., ... & Khanal, P. (2017). High rates of multidrug resistance among uropathogenic escherichia coli in children and analyses of esbl producers from nepal. Antimicrobial Resistance and Infection Control, 6 (1). https: //doi. org/10.1186/s13756 - 016 - 0168 - 6
- [17] Sane, R., Shahani, S., & Kalyanshetti, A. (2018). Antibiotic prescription pattern in surgical wards of mgm hospital, kamothe. International Journal of Infection, 5 (1). https://doi.org/10.5812/iji.57914
- [18] Sharmin, L., Haque, A., Begum, F., Parvez, A., & Uddin, B. (2020). Bacteriological profile and antimicrobial sensitivity pattern of urinary tract infection in a tertiary care hospital of bangladesh. Taj Journal of Teachers Association, 33 (2), 27 - 31. https: //doi. org/10.3329/taj. v33i2.51336
- [19] Shrestha, D., Thapa, P., Bhandari, D., Parajuli, H., Chaudhary, P., Thapa, K., ... & Shah, P. (2018). Biofilm production and antimicrobial resistance among uropathogens in pediatric cases: a hospital based study. Journal of Nepal Health Research Council, 16 (2), 178 - 183. https: //doi. org/10.33314/jnhrc. v16i2.921
- [20] Singh, V., Chaudhary, M., Banjara, M., & Tuladhar, R. (2020). Monitoring antimicrobial susceptibility in

## Volume 13 Issue 6, June 2024 Fully Refereed | Open Access | Double Blind Peer Reviewed Journal

<u>www.ijsr.net</u>

bacterial isolates causing urinary tract infections in a tertiary hospital in kathmandu. Nepal Journal of Science and Technology, 19 (1), 133 - 141. https://doi.org/10.3126/njst.v19i1.29794

- [21] Thapa, P., Sunar, A., Lamichanne, D., K. C, A., Dhungana, A., Paudel, R., ... & Tiwari, B. (2020). Antimicrobial sensitivity pattern of bacteria causing urinary tract infection among diabetic patients. https: //doi. org/10.21203/rs.2.24215/v2
- [22] Trivedi, M. and Nagendra, M. (2023). Characteristics of blood infections and phenotypic detection of extended - spectrum  $\beta$  - lactamases in icu patients in central india posing a challenge. International Journal of Research in Medical Sciences, 11 (4), 1296 - 1302. https://doi.org/10.18203/2320 - 6012. ijrms20230879
- [23] Uddin, B. (2023). Antibiotic susceptibility of bacterial strains isolated from urinary tract infections in a private laboratory of dhaka city. Bangladesh Journal of Medical Microbiology, 16 (1), 3 - 7. https: //doi. org/10.3329/bjmm. v16i1.65795
- [24] Var, S., Hadi, R., & Khardori, N. (2015). Evaluation of regional antibiograms to monitor antimicrobial resistance in hampton roads, virginia. Annals of Clinical Microbiology and Antimicrobials, 14 (1). https://doi.org/10.1186/s12941 - 015 - 0080 - 6
- [25] Vaughn, V., Saint, S., Greene, M., Ratz, D., Fowler, K., Patel, P., ... & Krein, S. (2020). Trends in health care–associated infection prevention practices in us veterans affairs hospitals from 2005 to 2017. Jama Network Open, 3 (2), e1920464. https: //doi. org/10.1001/jamanetworkopen.2019.20464
- [26] Walia, K., Madhumathi, J., Veeraraghavan, B., Chakrabarti, A., Kapil, A., Ray, P., ... & Ohri, V. (2019). Establishing antimicrobial resistance surveillance & amp; research network in india: journey so far. The Indian Journal of Medical Research, 149 (2), 164. https: //doi. org/10.4103/ijmr. ijmr\_226\_18