International Journal of Science and Research (IJSR) ISSN: 2319-7064

Impact Factor 2024: 7.101

Isolated and Combined Effect of Asanas and Pranayama Practice on Pulmonary Function Test among College Level Men

Dr. A. Senthil Kumar

Director of Physical Education, NIFT-TEA College of Knitwear Fashion, Tiruppur, Tamil Nadu, India

Abstract: Human movement is the result of complex interactions between environmental factors and the nervous, muscular and skeletal systems. Brain cell activities within the cerebral cortex are converted by supraspinal centre programming into neural outputs that stimulate the muscular system to produce the requirement. Yoga is a system of attaining perfect physical and mental health. Yoga controls one's sense resulting in an integrated personality. Positive changes in the life style of the people can be brought through by yoga. The study aims to found out the isolated and combined effect of asanas and pranayamas on pulmonary function test among college level men. To achieve these purpose sixty (N = 60) middle aged men were selected randomly from the NIFT-TEA College of Kntwear Fashion, Tiruppur, Tamil Nadu as subjects. Their aged ranged from 18 to 23 (18± 2.8) years. They were divided into three equal groups (n=20). Group I underwent asanas, Group II underwent pranayamand Group III underwent combined (Asanas and Pranayama) alternate sessions practices - 40 to 60 min/session/4days/week for 12 weeks. Data were collected on forced vital capacity (FVC) by PC based USB spirometer before and after the intervention programme. Dependent't' test was used to find out the difference between pre and post fest. After eliminating the influence of pre-test on post-test means of experiment groups the analysis of covariance (ANCOVA) was used to find out the mean gain differences. In addition to this Scheffe's test was employed (p < 0.05). From the results and limitation of this study it was concluded that, twelve weeks of asanas, pranayama and combined practice are effective method to improve forced vital capacity of the middle-aged men. Among the experimental group combined asanas and pranayama group improved FVC much better than the other groups.

Keywords: yoga practice, pulmonary function, asanas, pranayama, forced vital capacity

1. Introduction

The human body and mind are gifts of God: The capital given to us without any interest payable. For anything in life to be fulfilled, the body should be healthy and this needs constant looking after. Just as a car needs constant care, the body and mind need the same. This is the best done by regular exercise. Awareness of this certainly better now than many years ago, but exercise-oriented persons still form a minority in our vast world. In today's fast world, with easy availability of different types of cuisines, both vegetarian and non-vegetarian, junk-food, alcohol and cigarettes and the constant bombarding of our consciousness with advertisement of such foods through different media, we succumb, with resultant health problems The great science of yoga is India's unequalled gift of mankind.

If mankind is to evolve further and if it is to save itself from its own aggressive tendencies, the only path open is through the science of yoga. Though the ultimate goal of this science is the realization of the absolute, in day to day life it is useful and necessary to maintain mental and bodily health. Bodily exercises (asanas), breath control (pranayama) and mind control (dhyana) are all helpful to conquer bodily and mental ills.

The pulmonary and circulatory system are responsible for moving blood from the heart to the lungs and back to the heart and getting rid of waste products in the blood while helping to distribute blood rich in oxygen. Exercise can help strengthen and make this system more efficient. Pulmonary functions are generally determined by respiratory muscle strength, compliance of the lung and thoracic cavity airway resistance and elastic recoil of the lungs It is acknowledged

that low physical activity and fitness level are associated with ill health. Although there is evidence that exercise can be of benefit of health, it must be regular if these benefits are to be attained/maintained. It is important that regular physical activity is encouraged and that appropriate exercise sessions are promoted. However, the mode of exercise must be acceptable to the target population - that is, the participants must find the activity enjoyable and be able to participate regularly, and it may be necessary to show that the exercise prescription is effective in improving fitness levels.

Asana tones up the nervous system function of all vital internal organs, stimulation of the glands and regulation of the blood flow. The muscles in our body are thus formed and strengthened. The exercise of pranayama the correct breathing technique helps to manipulate our energies. Most of us breathe incorrectly, using only half of our lung capacity. Pranayama is an art and has techniques to make the respiratory organs to move and expand intentionally, rhythmically and intensively. It consists of long sustained suitable flow inhalation (puraka), exhalation (recaka) and retention of breath (kumbhaka). Physically, pranayama appears to be a systematic exercise of respiration, which makes the lungs stronger, improves blood circulation, makes the man healthier and bestows upon him the boon of a long life. Hence, the study aim to found out the isolated and combined effect of asanas and pranayamas on vital capacity of middle aged men.

2. Methodology

To achieve these purpose sixty (N = 60) college level men were selected randomly from NIFT-TEA College of

Volume 14 Issue 12, December 2025
Fully Refereed | Open Access | Double Blind Peer Reviewed Journal

Paper ID: SR251209105529

International Journal of Science and Research (IJSR) ISSN: 2319-7064 Impact Factor 2024: 7.101

Kntwear Fashion, Tiruppur, Tamil Nadu as subjects. Their aged ranged from 18 to 23 (18 ± 2.8) years. They were divided into three equal groups (n=20). Group I underwent asanas (Suryanamaskar, Tadasana, Trikonasana and Paschimothanasana), Group II underwent pranayama (Nodisodhana, Samavrithi, Bastrika, Ujjayi, Kapalabhati and Bharamari) and Group III underwent combined (Asanas and Pranayama) alternate sessions practices - 40 to 60 min/session/4days/week for 12 weeks. Data were collected on forced vital capacity (FVC) by PC based USB spirometer

before and after the intervention programme. Dependent't' test was used to find out the difference between pre and post fest. After eliminating the influence of pre-test on post-test means of experiment groups the analysis of covariance (ANCOVA) was used to find out the mean gain differences. In addition to this Scheffe's test was employed (p < 0.05).

3. Results

Table I: ANCOVA and 't' of Experimental Groups on Forced Vital Capacity


Variables	Group	Adjusted post- test mean	't' Value	% improvement	'F' Value	
Forced Vital Capacity (Litter)	Asana	3.12	32.90*	19.89		
	Pranayama	3.18	31.84*	18.02	5.17*	
	Combined	3.20	35.90*	2015		

^{&#}x27;t' table value 2.09 and F table value 3.16 respectively; *p > 0.05

Table II: Scheffe's Test for Means

Variables	Asana	Pranayama	Combined	MG	CI
Yoga	3.12	3.18	-	0.06	
Aerobic	3.12	-	3.20	0.08	0.05
Combined	-	3.18	3.20	0.02	

From the result it was clean that all the experimental (asanas, pranayama and combined) groups improved FVC of the college level men. The improvement of FVC was much better for combined group followed by asana and pranayama groups respectively. The magnitude of improvement also reveals the same.



4. Discussion

Physical inactivity and low cardio-respiratory fitness are recognized as important causes of morbidity and mortality. It is generally accepted that people with higher levels of physical activity tend of have higher levels of fitness and that physical activity tend to have higher levels of fitness and that physical activity can improve cardio-respiratory fitness. In the present study FVC increased significant in the experiment groups after twelve weeks of asanas, pranayama and combination of asana and pranayama practices.

Yoga (asanas), quietitude in the body always stabilizes irritable response. The parasympathetic tone is in excess in asthuma; exercise always help balance the autonomic tone by raising the sympathetic. It is possible to change the very responsiveness of the body. It is a matter of training. Asanas do their part by helping to ease the congestion in the lungs. This promotes relaxation of smooth muscle and lungs. This promotes relaxation of smooth muscle and better

oxygenation. Exercise is the only way to help improve the mechanical efficiency of the lungs. The usual repertoire of exercises is too strenuous for the lungs. Asanas that do not raise the respiratory rate and yet help excretion of the sputum, increases oxygen levels in the blood and elastic recoil of the lungs, prevent recurrent infections, and aerate the whole lungs, invigorating the lungs function at the end of the session. Yoga stands out as the only system eminently meeting all these requirements process. Regular practice of pranayama improves ventilation, better control of smooth muscle as the process of breathing helps in voluntary opening of airways. Cardio respiratory fitness significantly improved and breathlessness decreased over a wide range of work corresponding to activities of daily living. FVC in health people; and thus provides further support for the asanas and pranayama exercise being an important component of pulmonary rehabilitation. Present study also correlates with the above findings and showed that the experimental groups asana, pranayama and combined practice were able to have more powerful and more effective expiration as opposed to what they had been before.

5. Conclusion

From the results and limitation of this study it was concluded that, twelve week of asanas, pranayama and combined practice are effective method to improve forced vital capacity of the college level men. Among the experimental group combined asanas and pranayama group improved FVC much better than the other groups.

References

- [1] Gilbert C. Yoga and bteathing. J bodywork Mov Ther.1999; 3: 44-54.
- [2] Dr Vinayak P Doijad, Dr Anil. D. Surdi, Effect of short term yoga practice on pulmonary function tests. Indian Journal of Basic and Applied Medical Research.2012; 3 (1): 226-30.
- [3] Mandanmohan, Jatiya L, Udapa K, and Bhavanani AB. Effect of yoga training on handgrip, respiratory pressure and pulmonary function. Indian J Physiol Pharmacol. 2003; 47 (2): 387-92
- [4] Chandrasekaran, K. (1999). Sound Health through Yoga. Tamil Nadu: Prem Kalyan Publications.

Volume 14 Issue 12, December 2025
Fully Refereed | Open Access | Double Blind Peer Reviewed Journal
www.ijsr.net

International Journal of Science and Research (IJSR) ISSN: 2319-7064 Impact Factor 2024: 7.101

- [5] Menon N. Narayana (1984), "Yogasana for Health and Longevity", Madras: Sura Books Pvt. Ltd., P.1.
- [6] Iyengar, B. K. S. (1981), "The Light on Pranayama: The Yogic Art of Breathing", Crossroad Publishing Company; New York.
- [7] Nancy (1986), The Art of Breathing, Bantam Books Publishers; New York, PP.56-61.
- [8] Frostell C, Pande JN, Hedenstierna G. Effects of high-frequency breathing on pulmonary ventilation and gas exchange. J Appl Physiol 1983; 55: 1854-61.

Volume 14 Issue 12, December 2025
Fully Refereed | Open Access | Double Blind Peer Reviewed Journal
www.ijsr.net