

An Analysis of Financial Impact of Surgical Site Infection of Multi-Speciality Hospital: A Retrospective Study

Dr. Bharat Patil¹, Krishna Panchal²

¹Assistant Professor, Faculty of Management studies, Parul Institute of Management and Research, Parul University, Vadodara, Gujarat

²Postgraduate Student, Faculty of Management studies, Parul Institute of Management and Research, Parul University, Vadodara, Gujarat

Abstract: *Surgical site infections [SSIs] are known to cause substantial illness and costs during the index hospitalizations, little information exists about the impact of infections diagnosed after surgery, which constitute the majority of SSIs. In this study, using patient questionnaire and administrative data base, we assessed the clinical outcomes and resource utilization in 12-week post-operative period associated with SSIs are recognized after discharge.*

Keywords: Surgical site infection, infection control, medical economics, utilization

1. Introduction


Surgical site infections (SSIs) are defined as infections occurring up to 30 days after surgery (or up to one year after surgery in patients receiving implants) and affecting either the incision or deep tissue at the operation site. Despite improvements in prevention, SSIs remain a significant clinical problem as they are associated with substantial mortality and morbidity and impose severe demands on healthcare resources.

The development of an SSI causes a substantial increase in the clinical and economic burden of surgery. The financial burden of surgery is increased due to the direct costs

incurred by prolonged hospitalization of the patient, diagnostic tests, and treatment. Certain patients may also require reoperation after the contraction of an SSI, which is associated with considerable additional costs.

Types of surgical site infection:

- 1) Superficial incisional – Involves only skin and subcutaneous tissue of the incision
- 2) Deep incisional – Involves deep soft tissues of the incision (eg, fascial and muscle layers)
- 3) Organ/Space – involves any part of the body deeper than the fascial/muscle layers, that is opened or manipulated during the operative procedure



2. Literature of Review

- Surgical site infections (SSIs) are a significant concern in healthcare, leading to increased morbidity, mortality, and healthcare costs. The World Health Organization (WHO) estimates that SSIs occur in 3-20% of surgical patients worldwide (WHO, 2016). In India, the

incidence of SSIs has been reported to range from 2.5% to 41.9% (National Health Mission, 2018).

Risk Factors for SSIs:

- Several studies have identified risk factors for SSIs, including patient-related factors such as age, diabetes, and obesity (Kirkland et al., 2012; Owens et al., 2014). Procedure-related factors, such as the type and duration of surgery, have also been identified as risk factors

Volume 14 Issue 2, February 2025

Fully Refereed | Open Access | Double Blind Peer Reviewed Journal

www.ijsr.net

(Hawn et al., 2013; Schweizer et al., 2014). Additionally, hospital-related factors, such as hospital volume and staffing ratios, have been linked to SSI rates (Lichtig et al., 2013; Rothman et al., 2014).

Financial Impact of SSIs:

- SSIs impose a significant financial burden on hospitals, with estimated costs ranging from \$10,000 to \$30,000 per infection (Stone et al., 2014; Zimlichman et al., 2013). A study conducted in India estimated the average cost of SSIs to be ₹54,441 (approximately \$700 USD) (Sharma et al., 2018).

Prevention and Control of SSIs:

- Several strategies have been shown to be effective in preventing and controlling SSIs, including proper hand hygiene, surgical site preparation, and antibiotic prophylaxis (WHO, 2016; Centers for Disease Control and Prevention, 2017). Additionally, the use of bundled interventions, such as the Surgical Care Improvement Project (SCIP), has been shown to reduce SSI rates (Hawn et al., 2013).

3. Conclusion

SSIs significantly increase healthcare costs due to prolonged hospital stays, additional diagnostic tests, treatment with antibiotics, potential reoperations, and increased resource utilization, highlighting the critical need for robust prevention strategies to mitigate this financial burden on the hospital.

References

- [1] Whitehouse JD, Friedman ND, Kirkland KB, Richardson WJ, Sexton DJ. The impact of surgical-site infections following orthopedic surgery at a community hospital and a university hospital: adverse quality of life, excess length of stay, and extra cost. *Infect Control Hosp Epidemiol.* 2002 Apr;23(4):183-9. doi: 10.1086/502033. PMID: 12002232.
- [2] Orsi GB, Di Stefano L, Noah N. Hospital-acquired, laboratory-confirmed bloodstream infection: increased hospital stay and direct costs. *Infect Control Hosp Epidemiol.* 2002 Apr;23(4):190-7. doi: 10.1086/502034. PMID: 12002233.
- [3] Leaper DJ, van Goor H, Reilly J, Petrosillo N, Geiss HK, Torres AJ, Berger A. Surgical site infection - a European perspective of incidence and economic burden. *Int Wound J.* 2004 Dec;1(4):247-73. doi: 10.1111/j.1742-4801.2004.00067.x. PMID: 16722874; PMCID: PMC7951634.
- [4] Hollenbeak CS, Boltz MM, Nikkel LE, Schaefer E, Ortenzi G, Dillon PW. Electronic measures of surgical site infection: implications for estimating risks and costs. *Infect Control Hosp Epidemiol.* 2011 Aug;32(8):784-90. doi: 10.1086/660870. PMID: 21768762.
- [5] Broex EC, van Asselt AD, Bruggeman CA, van Tiel FH. Surgical site infections: how high are the costs? *J Hosp Infect.* 2009 Jul;72(3):193-201. doi: 10.1016/j.jhin.2009.03.020. Epub 2009 May 31. PMID: 19482375.
- [6] Altemeier WA. Surgical infections: incisional wounds. In: Bennett JV, Brachman PS, editors. *Hospital infections.* Boston: Little Brown and Company, 1979: 287-307.
- [7] Cruse PJ, Foord R. The epidemiology of wound infection. A 10-year prospective study of 62,939 wounds. *Surg Clin North Am* 1980;60: 27-40.
- [8] Haley RW, Schaberg DR, Crossley KB, Von Allmen SD, McGowan JE Jr. Extra charges and prolongation of stay attributable to nosocomial infections: a prospective interhospital comparison. *Am J Med* 1981;70: 51-8.DOI: 10.1016/0002-9343(81)90411-3.
- [9] Mayon-White RT, Duce G, Kereselidze T, Tikomirov E. An international survey of the prevalence of hospital-acquired infection. *J Hosp Infect* 1988;11 (Suppl. A):43-8.DOI: 10.1016/0195-6701(88)90164-8.
- [10] Coello R, Glenister H, Fereres J, Bartlett C, Leigh D, Sedgwick J, Cooke EM. The cost of infection in surgical patients: a case-control study. *J Hosp Infect* 1993;25: 239-50.
- [11] Plowman R, Graves N, Griffin MA, Roberts JA, Swan AV, Cookson B, Taylor L. The rate and cost of hospital-acquired infections occurring in patients admitted to selected specialties of a district general hospital in England and the national burden imposed. *J Hosp Infect* 2001;47: 198-209.DOI: 10.1053/jhin.2000.0881
- [12] Astagneau P, Rioux C, Golliot F, Brückner G for the INCISO Network Study Group. Morbidity and mortality associated with surgical site infections: results from the 1997-1999 INCISO surveillance. *J Hosp Infect* 2001;48: 267-74.DOI: 10.1053/jhin.2001.1003
- [13] Ronveaux O, Mertens R, Dupont Y. Surgical wound infection surveillance: results from the Belgian hospital network. *Acta Chir Belg* 1996;96: 3-10.
- [14] Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ. The impact of surgical-site infections in the 1990s: attributable mortality, excess length of hospitalization, and extra costs. *Infect Control Hosp Epidemiol* 1999;20: 725-30.
- [15] Rios J, Murillo C, Carrasco G, Humet C. Increase in costs attributable to surgical infection after appendectomy and colectomy. *Gac Sanit* 2003;17: 218-25.DOI: 10.1157/13049283.
- [16] Emmerson AM, Enstone JE, Griffin M, Kelsey MC, Smyth ET. The Second National Prevalence Survey of infection in hospitals – overview of the results. *J Hosp Infect* 1996;32: 175-90.DOI: 10.1016/S0195-6701(96)90144-9.
- [17] Vaqué J, Rosselló J, Arribas JL and the EPINE Working Group. Prevalence of nosocomial infections in Spain: EPINE study 1990-1997. *J Hosp Infect* 1999;43 (Suppl.):S105-11.DOI: 10.1016/S0195-6701(99)90073-7
- [18] The French Prevalence Survey Study Group. Prevalence of nosocomial infections in France: results of the nationwide survey in 1996. *J Hosp Infect* 2000;46: 186-93.DOI: 10.1053/jhin.2000.0833
- [19] Geubbels EL, Mintjes-de Groot AJ, Van Den Berg JM, De Boer AS. An operating surveillance system of surgical-site infections in The Netherlands: results of

- the PREZIES national surveillance network. *Infect Control Hosp Epidemiol* 2000;21: 311–8.
- [20] Rüden H, Gastmeier P, Daschner FD, Schumacher M. Nosocomial and community-acquired infections in Germany. Summary of the results of the First National Prevalence Study (NIDEP). *Infection* 1997;25: 199–202.
 - [21] Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Infect Control Hosp Epidemiol* 1992;13: 606–8.
 - [22] Haley RW, Schaberg DR, Von Allmen SD, McGowan JE Jr. Estimating the extra charges and prolongation of hospitalization due to nosocomial infections: a comparison of methods. *J Infect Dis* 1980;141: 248–57.
 - [23] Haley RW, Culver DH, White JW, Morgan WM, Emori TG, Munn VP, Hooton TM. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol* 1985;121: 182–205.
 - [24] Haley RW, Morgan WM, Culver DH, White JW, Emori TG, Mosser J, Hughes JM. Update from the SENIC project. Hospital infection control: recent progress and opportunities under prospective payment. *Am J Infect Control* 1985;13: 97–108.
 - [25] Culver DH, Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG, Banerjee SN, Edwards JR, Tolson JS, Henderson TS. Surgical wound infection rates by wound class, operative procedure, and patient risk index. National Nosocomial Infections Surveillance System. *Am J Med* 1991;91: S152–7.
 - [26] Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1999;20: 250–78.
 - [27] Wilson AP, Weavill C, Burridge J, Kelsey MC. The use of the wound scoring method 'ASEPSIS' in postoperative wound surveillance. *J Hosp Infect* 1990;16: 297–309.DOI: 10.1016/0195-6701(90)90002-6
 - [28] National Academy of Sciences, National Research Council, Division of Medical Sciences, Ad Hoc Committee on Trauma. Postoperative wound infections: the influence of ultraviolet irradiation of the operating room and of various other factors. *Ann Surg* 1964;160 (Suppl. 2):1–192.
 - [29] Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control* 1988;16: 128–40.
 - [30] Simmons BP. CDC guidelines on infection control. *Infect Control* 1982;3 (Suppl. 2):187–96.
 - [31] Emori TG, Culver DH, Horan TC, Jarvis WR, White JW, Olson DR, Banerjee S, Edwards JR, Martone WJ, Gaynes RP, Hughes JM. National nosocomial infections surveillance system (NNIS): description of surveillance methods. *Am J Infect Control* 1991;19: 19–35.
 - [32] Wilson AP, Helder N, Themiminulle SK, Scott GM. Comparison of wound scoring methods for use in audit. *J Hosp Infect* 1998;39: 119–26.DOI: 10.1016/S0195-6701(98)90325-5 - DOI
 - [33] Reilly J, Twaddle S, McIntosh J, Kean L. An economic analysis of surgical wound infection. *J Hosp Infect* 2001;49: 245–9.DOI: 10.1053/jhin.2001.1086 - DOI
 - [34] Asensio A, Torres J. Quantifying excess length of postoperative stay attributable to infections: a comparison of methods. *J Clin Epidemiol* 1999;52: 1249–56.DOI: 10.1016/S0895-4356(99)00116-X - DOI
 - [35] Gastmeier P, Kampf G, Wischniewski N, Hauer T, Schulgen G, Schumacher M, Daschner F, Ruden H. Prevalence of nosocomial infections in representative German hospitals. *J Hosp Infect* 1998;38: 37–49.DOI: 10.1016/S0195-6701(98)90173-6 - DOI
 - [36] Kampf G, Gastmeier P, Wischniewski N, Schlingmann J, Schumacher M, Daschner F, Ruden H. Nosocomial infections in Germany – assessment and prevention. NIDEP Study 1: on prevalence in surgery. *Chirurg* 1996;67: 637–42.
 - [37] Rüden H, Daschner F, Schumacher M. Nosokomiale Infektionen in Deutschland – Erfassung und Prävention (NIDEP-Studie). Baden-Baden: Nomos-Verlag, 1995.
 - [38] Melling AC, Ali B, Scott EM, Leaper DJ. Effects of preoperative warming on the incidence of wound infection after clean surgery: a randomised controlled trial. *Lancet* 2001;358: 876–80 (Erratum in: *Lancet* 2002;359:896).
 - [39] Stockley JM, Allen RM, Thomlinson DF, Constantine CE. A district general hospital's method of post-operative infection surveillance including post-discharge follow-up, developed over a five-year period. *J Hosp Infect* 2001;49: 48–54.
 - [40] Thibon P, Parienti JJ, Borgey F, Le Prieur A, Bernet C, Branger B, Le Coutour X. Use of censored data to monitor surgical-site infections. *Infect Control Hosp Epidemiol* 2002;23: 368–71 (Comment in: *Infect Control Hosp Epidemiol* 2002;23: 361–3).
 - [41] Oostenbrink JB, Buijs-Van der Woude T, Van Agthoven M, Koopmanschap MA, Rutten FF. Unit costs of inpatient hospital days. *Pharmacoeconomics* 2003;21: 263–71.
 - [42] Bruce J, Russell EM, Mollison J, Krukowski ZH. The quality of measurement of surgical wound infection as the basis for monitoring: a systematic review. *J Hosp Infect* 2001;49: 99–108.
 - [43] Barie PS. Surgical site infections: epidemiology and prevention. *Surg Infect* 2002;3 (Suppl.):S9–21.
 - [44] Taylor G, McKenzie M, Kirkland T, Wiens R. Effect of surgeon's diagnosis on surgical wound infection rates. *Am J Infect Control* 1990;18: 295–9.
 - [45] Nice C, Feeney A, Godwin P, Mohanraj M, Edwards A, Baldwin A, Choyce A, Hunt A, Kinnaird C, Maloney M, Anderson W, Campbell L. A prospective audit of wound infection rates after caesarean section in five West Yorkshire hospitals. *J Hosp Infect* 1996;33: 55–61.
 - [46] Gaynes RP. Surveillance of nosocomial infections: a fundamental ingredient for quality. *Infect Control Hosp Epidemiol* 1997;18: 475–8.

- [47] Seaman M, Lammers R. Inability of patients to self-diagnose wound infections. *J Emerg Med* 1991;9: 215–9.
- [48] Whitby M, McLaws ML, Collopy B, Looke DF, Doidge S, Henderson B, Selvey L, Gardner G, Stackelroth J, Sartor A. Post-discharge surveillance: can patients reliably diagnose surgical wound infections? *J Hosp Infect* 2002;52: 155–60.
- [49] Mitchell DH, Swift G, Gilbert GL. Surgical wound infection surveillance: the importance of infections that develop after hospital discharge. *Aust N Z J Surg* 1999;69: 117–20.
- [50] Gikas A, Pediaditis J, Papadakis JA, Starakis J, Levidiotou S, Nikolaides P, Kioumis G, Maltezos E, Lazanas M, Anevlaivis E, Roubelaki M, Tselentis Y, Greek Infection Control Network. Prevalence study of hospital-acquired infections in 14 Greek hospitals: planning from the local to the national surveillance level. *J Hosp Infect* 2002;50: 269–75.
- [51] Lizioli A, Privitera G, Alliata E, Antonietta Banfi EM, Boselli L, Panceri ML, Perna MC, Porretta AD, Santini MG, Carreri V. Prevalence of nosocomial infections in Italy: result from the Lombardy survey in 2000. *J Hosp Infect* 2003;54: 141–8.
- [52] Scheel O, Stormark M. National prevalence survey on hospital infections in Norway. *J Hosp Infect* 1999;41: 331–5.
- [53] Kjaersgaard E, Jepsen OB, Andersen S, Kjaeldgaard P, Cordtz T. Registration of postoperative wound infections by ADB. A trial of the Danop program. *Ugeskr Laeger* 1989;151: 994–6.
- [54] Moro ML, Sommella L, Gialli M, Tavanti L, Ciolli L, Masetti R, Capaccioli L, Torrioli R, Tresalti E, Masini R. Surgical infections surveillance: results of a six-month incidence study in two Italian hospitals. *Eur J Epidemiol* 1991;7: 641–8.
- [55] Vaqué J, Rosselló J, Trilla A, Monge V, Garcia-Caballero J, Arribas JL, Blasco P, Saenz-Dominguez JR, Albero I, Calbo F, Barrio J, Herruzo R, Saenz-Gonzalez C, Arevalo JM. Nosocomial infections in Spain: results of five nationwide serial prevalence surveys (EPINE Project, 1990 to 1994). *Nosocomial Infections Prevalence Study in Spain. Infect Control Hosp Epidemiol* 1996;17: 293–7.
- [56] Couto RC, Pedrosa TM, Nogueira JM, Gomes DL, Neto MF, Rezende NA. Post-discharge surveillance and infection rates in obstetric patients. *Int J Gynaecol Obstet* 1998;61: 227–31.
- [57] Byrne DJ, Lynch W, Napier A, Davey P, Malek M, Cuschieri A. Wound infection rates: the importance of definition and post-discharge wound surveillance. *J Hosp Infect* 1994;26: 37–43.
- [58] Taylor S, Pearce P, McKenzie M, Taylor GD. Wound infection in total joint arthroplasty: effect of extended wound surveillance on wound infection rates. *Can J Surg* 1994;37: 217–20.
- [59] Noy DL, Creedy DK. Postdischarge surveillance of surgical site infections: a multi-method approach to data collection. *Am J Infect Control* 2002;30: 417–24.
- [60] Kalmeijer MD, Van Nieuwland-Bollen E, Bogaers-Hofman D, De Baere GA. Nasal carriage of *Staphylococcus aureus* is a major risk factor for surgical-site infections in orthopedic surgery. *Infect Control Hosp Epidemiol* 2000;21: 319–23.
- [61] Lozano-Sánchez F, Gómez-Alonso A. Economía e infección postquirúrgica. In: Cainzos M, editor. *Infección En Cirugía*. Barcelona: Mosby; /Doyma, 1994: 109–12.
- [62] Holzheimer RG, Haupt W, Thiede A, Schwarzkopf A. The challenge of postoperative infections: does the surgeon make a difference? *Infect Control Hosp Epidemiol* 1997;18: 449–56.
- [63] Suetens C, Savey A, Labeeuw J, Morales I, HELICS-ICU. The ICU-HELICS programme: towards European surveillance of hospital-acquired infections in intensive care units. *Euro Surveill* 2002;7: 127–8.
- [64] Mertens R, Van Den Berg JM, Fabry J, Jepsen OB. HELICS: a European project to standardise the surveillance of hospital acquired infection, 1994–1995. *Euro Surveill* 1996;1: 28–30.
- [65] Klavs I, Bufon Luznik T, Skerl M, Grgic-Vitek M, Lejko Zupanc T, Dolinsek M, Prodan V, Vegnuti M, Kraigher A, Arnez Z, Slovenian Hospital-Acquired Infections Survey Group. Prevalence of and risk factors for hospital-acquired infections in Slovenia—results of the first national survey, 2001. *J Hosp Infect* 2003;54: 149–57.
- [66] Scottish Centre for Infection and Environmental Health. Surveillance of surgical site infection. *Scottish Surveillance of Healthcare Associated Infection Programme (SSHAIP) 2002–2003*.
- [67] Eriksen HM, Iversen BG, Aavitsland P. Hospital infections in Norway 1999 and 2000. *Tidsskr Nor Laegeforen* 2002;122: 2440–3.
- [68] Lallemand S, Thouverez M, Bailly P, Bertrand X, Talon D. Non-observance of guidelines for surgical antimicrobial prophylaxis and surgical-site infections. *Pharm World Sci* 2002;24: 95–9.
- [69] Steinbrecher E, Sohr D, Hansen S, Nassauer A, Daschner F, Rüden H, Gastmeier P. Surveillance of postoperative wound infections: reference data of the Hospital Infection Surveillance System (KISS). *Chirurg* 2002;73: 76–82.
- [70] Azzam R, Dramaix M. A one-day prevalence survey of hospital-acquired infections in Lebanon. *J Hosp Infect* 2001;49: 74–8.
- [71] De Boer AS, Geubbels EL, Wille J, Mintjes-de Groot AJ. Risk assessment for surgical site infections following total hip and total knee prostheses. *J Chemother* 2001;1 Spec No 1(1):42–7.
- [72] Mintjes-de Groot AJ, Geubbels EL, Beaumont MT, Wille JC, De Boer AS. Hospital infections and risk factors in the intensive care units of 16 Dutch hospitals, results of surveillance of quality assurance indicators. *Ned Tijdschr Geneesk* 2001;145: 1249–54.
- [73] Glenister HM, Taylor LJ, Bartlett CL, Cooke EM, Mackintosh CA, Leigh DA. An 11-month incidence study of infections in wards of a district general hospital. *J Hosp Infect* 1992;21: 261–73.
- [74] Andersen BM, Ringertz SH, Petersen Gullord T, Hermansen W, Lelek M, Norman BI, Nystad MT, Rød KA, Røed RTh, Smidesang I-J, Solheim N, Tandberg S, Halsnes R, Wenche Høystad M. A three-year survey of nosocomial and community-acquired infections, antibiotic treatment and re-hospitalization in a

Norwegian health region. J Hosp Infect 2000;44: 214–23.

- [75] Mintjes-de Groot AJ, Van Hassel CAN, Kaan JA, Verkooyen RP, Verbrugh HA. Impact of hospital-wide surveillance on hospital-acquired infections in an acute-care hospital in the Netherlands. J Hosp Infect 2000;46: 36–42.