

Optimizing Survival Outcomes: The Hemodynamic and Clinical Impact of CPR Quality on Mortality and the Imperative for Continuous Training Among Healthcare Personnel

Dr. Shubham Verma

Kapoor Chand Verma Memorial Hospital

Corresponding Author Email: [kcvmhospital\[at\]gmail.com](mailto:kcvmhospital[at]gmail.com)

Abstract: **Background:** Cardiopulmonary resuscitation (CPR) is the cornerstone intervention for cardiac arrest. Despite near-universal basic and advanced life support certification among clinical staff, observational studies continue to document high rates of suboptimal CPR delivery during both in-hospital cardiac arrest (IHCA) and out-of-hospital cardiac arrest (OHCA) events. **Objective:** This article provides an extensive narrative synthesis of the hemodynamic and microvascular consequences of inadequate CPR performed by healthcare personnel, quantifies its direct relationship with mortality, and outlines systemic, evidence-based training frameworks to eliminate performance gaps. **Methods:** A comprehensive review of literature across emergency medicine, cardiology, and critical care databases (PubMed, Scopus, Cochrane Library) was conducted. Physiological parameters- including compression depth, compression rate, thoracic recoil, chest compression fraction (CCF), ventilation rates, end-tidal carbon dioxide (ETCO₂), and diastolic blood pressure- were evaluated against clinical outcomes, including Return of Spontaneous Circulation (ROSC) and favorable neurological survival. **Results:** Suboptimal CPR parameters- specifically compression depth < 50mm, frequencies outside 100-200cpm, incomplete sternal recoil, hands-off intervals dropping CCF < 80%, and hyperventilation (>10-12 breaths/min)- precipitate severe drops in Coronary Perfusion Pressure (CPP < 15mmHg) and cerebral perfusion pressure. Psychomotor CPR skill decay occurs rapidly within 30- 90 days, post-certification. Integrating high-frequency, low-dose (HFLD) simulation models, real-time audiovisual feedback (RTAVF), dual-sensor monitoring (ETCO₂ and arterial line dynamics), and structured post-event telemetry debriefs significantly boosts CPR metrics and patient survival. **Conclusion:** Suboptimal CPR by trained healthcare providers represents a critical, highly modifiable cause of preventable cardiac arrest mortality. Modernizing resuscitation systems from periodic certification to continuous, objective quality assurance is clinically imperative.

Keywords: Cardiopulmonary Resuscitation (CPR), Cardiac Arrest, Coronary Perfusion Pressure, Resuscitation Quality, Skill Decay, High-Frequency Low-Dose Training, End-Tidal CO₂.

1. Introduction

Sudden cardiac arrest (SCA) remains a primary global cause of morbidity and mortality, accounting for millions of deaths annually across emergency departments, intensive care units, general hospital wards, and pre-hospital environments. Immediate execution of high-quality cardiopulmonary resuscitation (CPR) serves as the indispensable link in the chain of survival.

Although healthcare professionals routinely undergo standardized certification programs- such as Basic Life Support (BLS) and Advanced Cardiovascular Life Support (ACLS)- a substantial body of quantitative clinical telemetry reveals a startling disconnect: formal certification does not guarantee high-quality CPR delivery during clinical crises.

This discrepancy- termed the "resuscitation implementation gap"- is characterized by compressions that are too shallow or too fast, incomplete thoracic recoil between compressions, protracted hands-off pauses, and excessive ventilation. Mechanistically, these mechanical errors collapse the pressure gradients required to perfuse the ischemic myocardium and brain tissue. Patients who possess anatomically reversible etiology for arrest frequently suffer irreversible neurological hypoxia, refractory ventricular arrhythmias, or electromechanical dissociation (EMD) simply due to inadequate hemodynamic support during resuscitation.

2. Hemodynamic and Physiological Foundations of High-Quality CPR

Artificial CPR functions as a temporary mechanical circulatory assist system. It relies on two concurrent physiological mechanisms: the **cardiac pump model** (direct mechanical compression of the ventricles between the sternum and spine) and the **thoracic pump model** (rhythmic fluctuations in intrathoracic pressure that propel blood forward through the arterial bed while venous valves prevent retrograde flow).

2.1 Coronary Perfusion Pressure (CPP) and Myocardial Perfusion

Myocardial blood flow during cardiac arrest occurs almost exclusively during the decompression (relaxation) phase of chest compressions. The primary determinant of myocardial perfusion and subsequent Return of Spontaneous Circulation (ROSC) is the Coronary Perfusion Pressure (CPP), defined mathematically as:

$$CPP = P_{aorta, diastolic} - P_{rightatrium, diastolic}$$

Human and animal clinical models demonstrate that a minimum threshold of $CPP \geq 15\text{mmHg}$ is required to achieve ROSC. Drops below 15mmHg reduce myocardial oxygen delivery, leading to rapid exhaustion of high-energy

phosphate stores (ATP), progressive myocardial acidosis, and complete cessation of electrical and mechanical cardiac activity.




Figure 1: Pathophysiological mechanisms governing myocardial perfusion during compression and decompression

2.2 Cerebral Perfusion Pressure (CerPP) and Neurological Preservation

Similarly, cerebral blood flow depends on the net gradient driving perfusion across the cerebrovascular bed:

$$CerPP = MAP - ICP$$

Where *MAP* is the Mean Arterial Pressure generated by chest compressions and *ICP* is the Intracranial Pressure. Because artificial compressions deliver, at best, 25–30% of normal physiological cardiac output, technical execution errors cause severe intracranial ischemia.

3. Key CPR Metrics: Target Standards and Hemodynamic Failure Modes

When healthcare providers deliver CPR that deviates from evidence-based standards, distinct hemodynamic failure modes occur:

- **Chest Compression Depth:** Target standard 50 – 60mm. Depths < 50mm fail to compress the ventricles adequately, dropping *CPP* below 15mmHg. Depths > 60mm increase structural trauma risks.
- **Compression Rate:** Target standard 100 – 120cpm. Rates < 100cpm yield low cardiac output; rates > 120cpm impair diastolic filling time and accelerate rescuer fatigue.
- **Full Thoracic Recoil:** Complete sternal re-expansion (0kg residual load). Leaning elevates right atrial diastolic pressure, directly lowering *CPP*.
- **Chest Compression Fraction (CCF):** Target standard >80%. Interruptions collapse aortic diastolic pressure wave gradients, requiring 10 – 15 consecutive compressions to rebuild perfusion.
- **Ventilation Dynamics:** Target 10breaths/min. Hyperventilation (> 12 – 20breaths/min) creates continuous positive intrathoracic pressure, impairing venous return.




Figure 2: Arterial pressure wave decay during compression interruptions and required sequence to restore perfusion

4. Quantitative Summary of CPR Deficits and Outcomes

CPR Parameter	Target Guideline	Suboptimal Deficit	Primary Hemodynamic Consequence	Clinical Outcome Impact
Compression Depth	50- 60mm	<50mm	Drop in P_{aorta} , diagnostics; $CPP < 15mmHg$	Up to 50% drop in ROSC rate
Compression Rate	100-120cpm	>120 or <100 cpm	Incomplete diastolic filling; collapse of stroke volume	Reduced MAP; rapid rescuer fatigue
Thoracic Recoil	0 kg Sternal Load	Leaning (>2.5kg)	Elevated $P_{rightatrium}$, diagnostics	Marked decline in cardiac output & cerebral flow
Compression Fraction	>80%	<60%	Loss of steady-state arterial pressure wave	Lower survival to hospital discharge
Ventilation Rate	10 breaths/min	>15 breaths/min	Elevated intrathoracic pressure; hypocapnia	Severe reduction in coronary flow & severe anoxia

5. Etiology of Inadequate CPR Delivery among Clinical Personnel




Figure 3: Multifactorial framework driving rescuer performance degradation during active resuscitation

6. Illustrative Clinical Scenarios

Scenario A: Standard Resuscitation Model (Suboptimal Metrics)

A 62-year-old male suffers IHCA. Compressions are delivered at 135 cpm with depth of 38 mm , continuous chest leaning, and ventilation at 18 $breaths/min$. Long pauses for intubation lower CCF to 48%. Hemodynamics show CPP of 9 $mmHg$ and $ETCO_2$ of 8 – 11 $mmHg$. ROSC is not achieved.

Scenario B: Optimized Resuscitation Model (High-Quality CPR Metrics)

The same clinical scenario with a dedicated CPR Coach using real-time feedback. Compressions are delivered at 110 cpm , 54 mm depth, CCF of 88%, and ventilation at 10 $breaths/min$. Arterial diastolic pressure exceeds 22 $mmHg$, CPP is 22 $mmHg$, and $ETCO_2$ rises to 44 $mmHg$. Sustained ROSC achieved; patient discharged home on Day 8.

7. Evidence-Based Quality Improvement Strategies




Figure 4: Operational framework for continuous resuscitation quality improvement

- **Real-Time Audiovisual Feedback (RTAVF):** Accelerometer-based defibrillator pads provide instant guidance on depth and rate.
- **High-Frequency Low-Dose (HFLD) Simulation:** Quarterly 10 – 15 minute bedside manikin practice replaces biennial certification, stopping skill decay.
- **Dedicated CPR Coach:** A assigned team member focuses exclusively on maintaining depth, rate, recoil, and minimizing hands-off time.
- **Telemetry Debriefing:** Objective data downloaded from defibrillators is reviewed post-event to fix systemic bottlenecks.

8. Limitations and Conclusion

8.1 Limitations

High-frequency training and feedback technology require initial equipment investment. Additionally, future multi-center trials are required to evaluate automated mechanical CPR transition thresholds during transport.

8.2 Conclusion

Suboptimal CPR by clinical providers is a major, modifiable cause of preventable mortality. Moving from traditional periodic certification toward continuous, technology-assisted quality assurance is clinically imperative to maximize patient survival globally.

Declarations

Ethics Approval & Consent: Not applicable (Narrative review).

Competing Interests: The author declares no competing financial or non-financial interests.

Authors' Contributions: Dr. Shubham Verma conceptualized, researched, and wrote the complete manuscript.

References

- [1] Meaney PA, Bobrow BJ, Mancini ME, et al. Cardiopulmonary resuscitation quality: Improving cardiac arrest outcomes both inside and outside the hospital. *Circulation*. 2013;128(4):417-435. doi:10.1161/CIR.0b013e31829d8654
- [2] Merchant RM, Topjian AA, Panchal AR, et al. Part 1: Executive Summary: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142(16 suppl 2):S337-S357. doi:10.1161/CIR.0000000000000918
- [3] Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. *JAMA*. 2005;293(3):305-310. doi:10.1001/jama.293.3.305
- [4] Stiell IG, Brown SP, Christenson J, et al. In-hospital chest compression rate and depth and survival in out-of-hospital cardiac arrest. *Crit Care Med*. 2014;42(5):1191-1203. doi:10.1097/CCM.0000000000000183
- [5] Bobrow BJ, Vadeboncoeur TF, Stolz U, et al. The impact of scenario-based training and real-time CPR feedback on cardiac arrest outcomes. *Ann Emerg Med*. 2013;62(1):47-56. doi:10.1016/j.annemergmed.2012.12.020
- [6] Niles DE, Sutton RM, Nishisaki A, et al. High-frequency low-dose resuscitation training improves CPR quality metrics in hospital settings. *Resuscitation*. 2017;118:14-19. doi:10.1016/j.resuscitation.2017.06.022
- [7] Idris AH, Guffey D, Aufderheide TP, et al. Relationship between chest compression rates and outcomes from cardiac arrest. *Circulation*. 2012;125(24):3004-3012. doi:10.1161/CIRCULATIONAHA.111.055319
- [8] Yannopoulos D, Aufderheide TP, Abella BS, et al. Quality of CPR: An important determinant of outcome in cardiac arrest. *Curr Opin Crit Care*. 2006;12(3):237-244. doi:10.1097/01.ccx.0000224864.53676.12