

# Management of Root-Knot Nematode *Abelmoschus esculentus* Using Botanical Amendments

Dr. Resha

Department of Science, Shree Dronacharya P.G College, Dankaur G. B. Nagar, (U. P.), India

Email: [reshareshu\[at\]gmail.com](mailto:reshareshu[at]gmail.com)

**Abstract:** Root-knot nematode (*Meloidogyne incognita*) is one of the most destructive plant-parasitic nematodes causing significant yield losses in okra (*Abelmoschus esculentus*). Excessive dependence on chemical nematicides poses environmental and health risks, highlighting the need for eco-friendly management strategies. The present study was conducted to evaluate the efficacy of selected botanical amendments for the management of *M. incognita* in okra. Botanical materials such as neem cake, neem leaf powder, mustard cake and other locally available plant-based amendments were applied to the soil under controlled field/ pot conditions. Observations were recorded on plant growth parameters, nematode population density, root gall index and yield attributes. Results revealed that botanical amendments significantly reduced nematode infestation and root gall formation compared to the untreated control. Among the treatments, neem-based amendments showed maximum suppression of nematode population along with improved plant growth and yield. The effectiveness of botanicals may be attributed to the release of nematicidal compounds and enhancement of soil health. The study concludes that botanical amendments can serve as an effective, sustainable and environmentally safe alternative for the management of root-knot nematode in okra cultivation.

**Keywords:** *Meloidogyne incognita*, Amendments of *Azadirachta indica*, *Abelmoschus esculentus*, dominant plant-parasitic nematodes

## 1. Introduction

Parasitic nematodes (PPNs) are microscopic tiny and hidden worms that feed on roots of plants, causing damage and disease. PPNs are obligate parasites, meaning they require a plant host to survive and reproduce. *Meloidogyne* spp., is a World -Wide distributed and prevalent plant-parasitic nematode in India, and its population density can vary depending on edaphic factors, the region, soil type, and crop grown. Some *Meloidogyne* population densities in different regions of India. My experimental crop Okra (*Abelmoschus esculentus*), it is a very important food and cash crop by nutrition, health and economic wise. In the present study the management of *Meloidogyne* spp., on okra crop through the organic amendments (choose the nematicidal plant *Azadirachta indica*). The application of various parts of *Azadirachta indica* (oils, cake, seeds, and leaves) altered the physiology of host plant (okra) and developed the strong defensive mechanism of the root against nematodes. so the use of Neem products stimulated and change the physiology of plant cells and tissue to repel the plant nematode parasite. Extracts of fresh greens leaves, showed max. Reduction in egg hatching and cause great mortality of juveniles and followed by fresh green seed and dry seed extracts. In vivo and vitro conditions the powder of fresh seed showed highest plant growth and reduction in root knot formation and followed by fresh leaves and dry seed powder. Overall, *Meloidogyne incognita* is a significant pest in Indian agriculture, and its management is crucial for maintaining crop productivity and soil health.

Nematodes, especially the plant-parasitic are the most abundantly evolved and diversified metazoans on the planet Earth. Nematodes which infest the Kingdom of plantae which known as plant parasitic nematodes, Phyto-nema or Phyto-nematodes which damage more than seventy billion dollars of crop loss in the World each year. PPNs can cause various disease of crop, some symptoms of crop are include: Root-knot or gall formation, Stunted or distorted plant growth,

Yellowing or wilting of leaves and reduced yields or plant death.

These tiny worms can have a significant impact on plant health and agricultural productivity! *Meloidogyne incognita* is a species of plant-parasitic nematode, commonly known as the southern root-knot nematode (SRKN). *M. incognita* is a major pest of many crops, including vegetables, fruits, and ornamentals. It infects plant roots, causing the formation of characteristic swellings or "galls" (also called "root knots"). The nematode's life cycle includes: Egg stage: eggs are laid in the soil, Juvenile stages: hatch from eggs and migrate to plant roots, Adult stage: mature nematodes feed on plant tissues, causing damage. *M. incognita* can lead to significant yield losses, reduced plant growth, and increased susceptibility to other diseases. Management strategies include crop rotation, resistant cultivars, and nematicides.

The highly dangerous Endo-parasitic nematode *Meloidogyne* spp., represented about 300 species all over the World. *Meloidogyne incognita* is a widely distributed and prevalent plant-parasitic nematode in India, and its population density can vary depending on the region, soil type, and crop grown.

However, here are some approximate estimates of *Meloidogyne* population densities in different regions of India: North India: 200-500 eggs per 100 cc of soil (moderate to high infestation). South India: 100-300 eggs per 100 cc of soil (low to moderate infestation), East India: 150-400 eggs per 100 cc of soil (moderate infestation), West India: 250-600 eggs per 100 cc of soil (high infestation), above study show the highest infection is an infected in Western Uttar Pradesh due the favorable environmental conditions like temperature, moisture and pH etc., for growth of *Meloidogyne* spp.

Overall, *Meloidogyne incognita* is a significant pest in Indian agriculture, and its management is crucial for maintaining crop productivity and soil health. Critical conditions for *Meloidogyne incognita* infestation in okra crops in India.

Volume 15 Issue 2, February 2026

Fully Refereed | Open Access | Double Blind Peer Reviewed Journal

[www.ijsr.net](http://www.ijsr.net)

Neem components have attracted global attention for their insecticidal, fungicidal, bactericidal and nematocidal properties. Crude Neem extracts have been used at a local and small-farm level for some time in countries where Neem grows. In the major countries such as USA, Canada and Europe, the commercial Neem insecticides have reached the markets. Integrated Pest Management (IPM) strategies that combine these methods can provide effective overall control of *Meloidogyne incognita* infestation in okra crops.

It's important to note that the most effective method will depend on the severity of infestation, soil type, climate, and other factors, so it's important to monitor the field and adjust the control strategy accordingly.

The effective, inexpensive and environmental friendly control of this major pest is a serious problem. Although the chemical control is effective but their residues remain undegradable in soil for long time. Therefore most of the chemical nematicides are banned now a day. Botanical nematicides are gaining attention because they can be combined with other sustainable practices (crop rotation, resistant cultivars) for better control. A 2024 study showed that neem leaf extract significantly suppressed egg hatchability of *Meloidogyne* spp. (root-knot nematodes) in vitro, with increasing inhibition at higher extract concentrations and longer exposure times. Various workers contributed significantly to the management of root knot nematode. Some of the significant contributors are A. K. Singh et al (2020), J. Singh et al (2019), Graham R. Stirling and John Bridge (2017), Wani (2006); Ram et. al (2009); Sharma et al (2007); Khan (2012); Mohnta et al (2012). To overcome this problem various soil amendments by green and compost Goswami & Neetu Singh (2012). manures, organic, inorganic fertilizers, botanicals, bioagents like nematophagous fungi, soil bacteria are found to be more satisfactory to reduce the nematode population and increased crop yield. R. K. Singh et al (2018) have reported integrated management of root-knot nematode on okra using biological control methods, These Neem extracts can be used as natural nematicides, either alone or in combination with other methods, to manage *Meloidogyne incognita* infestation in okra crops. However, it's important to note that the efficacy of these Neem extracts may vary depending on the severity of infestation, soil type, and other factors.

## 2. Materials and Methods

**In vitro study:** in vitro and vivo study was conducted to test the ovicidal and larvicidal activity of Neem extracts on eggs of dominant plant parasitic nematodes and second stage juveniles. The extracts of fresh green leaves of neem, fresh green and dry seeds were made by chopping and crushing of leaves and seeds [at]20g/70ml in distilled water (w/w) in sox let apparatus at 69±4°C for 43 hours. All the extracts were filtered through what man filter paper and reduced on hot plate and stored in sterilized vials. The standard extracts (10%) were made by {1.5g extracts dissolving in 8.5ml distilled water (w/w) }. Further dilutions were made as 1% (1.5ml of standard extracts dissolved in 8.5ml distilled water).5ml of all the dilutions were poured in glass petri plates and surface sterilized single egg mass was placed in all the plates and replicated thrice. All the plates were incubated in BOD at 27±2°C for 7 days. After 6 days of incubation

hatching of eggs and subsequent mortality of juveniles was recorded under phase contrast microscopy.

**In vivo and vivo study: Pot and field** experiment was conducted on okra plants to root knot nematode, to observe the efficacy of amendments of Neem such as leaves, fresh and dry seed powder, after the establishment of infection by second stage IJ<sub>2</sub>. The study was carried out in following steps:

- 1) **Sowing of seed:** The okra seeds were procured from the NSC, New Delhi. Seeds were raised in the sterilized soil along with the compost manure[at]160g/kg soil, at Nematology Laboratory, Department of Zoology, C. C. S. University, and Meerut. Earthen pots were filled with 1kg sterilized soil and four seeds were sowed in each pot. After 16 days of germination three seedlings was maintained in each pot. And in vivo study, seed of okra was sowing in a field (small field approx. .100meters) of ccs university
- 2) **Culture of (IJ<sub>2</sub>):** The infected okra plant samples were collected from the agricultural fields of Meerut district. The plants were carefully uprooted and after removing the soil, the roots were placed in well labelled polyethylene bags. The collected roots were washed with tap water & eggs masses were isolated in petri plates. The isolated egg masses were incubated for five days at 27±2°C in BOD. Freshly, hatched IJ<sub>2</sub> were collected in water in 250ml beaker. The content was further reduced to small quantity to count the no. of nematodes. Homogeneous aliquot was placed in counting dish and the no. of IJ<sub>2</sub> in 1ml was under the stereoscopic microscope.
- 3) **Experimental Design:** After 3 -4 weeks of seed sowing, okra seedlings were inoculated with freshly hatched *M. incognita* infective[at]1000 IJ<sub>2</sub>. After 7days of inoculation plants were treated with the fresh green leaves, dry and fresh seed powder of Neem[at]8% powder/ soil (w/w). All the experimental pots were replicated thrice and irrigated regulatory as needed.
- 4) **Termination of experiment:** plants were uprooted at 65 days after inoculation. After washing, data were recorded on various plant growth parameters viz. -length of root and shoot, weight of fresh and dry shoot and root and also on disease parameters (root knot index) on the basis of no. root knots/plant. The scale (0-5) given by Taylor & Sasser (1978), was applied to find out root knot index as follows: 1, 0= no galling, 2, 1= light galling (0-25%), 3, 2= moderating galling (25-50%), 4, 3= severe galling (51-75%), 5, 4= very severe galling (76-100%).

## 3. Result and Discussion

An investigation was carried out to study the pathogenicity of root knot nematode, *Meloidogyne incognita* on okra and its management through organic amendments. The present investigation was conducted to okra plants in pot and field conditions. The results revealed determine the efficacy of Neem leaves and seeds (fresh & dry) powder against *M. incognita* infesting okra plants in pot and field conditions. The result revealed that the inoculated plants treated with FSP showed max. Plant growth parameters viz, length (shoot & root) and (fresh & dry) weight (shoot & root) and the plant disease parameters viz, no. Of root galls/root; no. Of egg mass/root; root gall index followed by FGP & DSP. The data

calculated on % reduction in root knot formation in all the treatments showed highest reduction (64.76%) in FSP alone and followed by combined FSP+FLP (64.65%); DSP+FLP (57.73%); FLP (44.26%) & DSP alone (26.29%). The estimation of egg masses reduction was maximum in combined treatment of FSP+FLP (55.16%) and slightly low (53.56%) in FLP alone followed by DSP+FLP (43.25%); FLP (39.77%) and DSP alone (35.27%). The work of Tariq & Siddiqui, 2005; Rather & Siddiqui, 2007; they also found same result in tomato also.

Azadirachtin is the main active and nematicide content of Neem and is reported very effective and target specific to controlling insects and nematodes pests of the various crops. The findings of the present study can be correlated with of R. K. Singh et al (2022) reported that “impact of organic amendments on *Meloidogyne incognita* population and okra yield, Graham R. Stirling and John Bridge (2017), they observed integrated management of root-knot nematodes on okra using crop rotation and organic amendments, A. K. Singh et al (2020) this study evaluated the management of *Meloidogyne incognita* on okra using bioagents and organic amendments, J. Singh et al in 2019 observed some nematicidal activity of some botanicals against *Meloidogyne javanica* on okra, R. K. Singh et al (2018) have reported integrated management of root-knot nematode on okra using biological control methods, Egumjobi & Afolami (1976) where they have reported the successful use of Neem leaves extracts in nematode control. Mishra (1999) states that the Neem formulations are most effective control of root knot nematode *Meloidogyne* sp. as compared to other botanicals.

Frontiers in Plant Science (2025) published research on the effect of botanicals and biocontrol agents (including neem formulations) on *Meloidogyne incognita* infecting tomato, showing enhanced nematode suppression and plant performance under botanical treatments Archana & Prasad (2014) states that organic amendments of soil using dried poultry litter, municipal refuse, oil cakes of ground nut, Neem mustard & neem products (which are commercial available in market) have been found effective in the control of *Meloidogyne incognita*. These all studies provide valuable insights into the management of okra infestation by nematodes using various approaches including biological control, organic amendments and crop rotation. Management of *Meloidogyne incognita* on okra using bioagents and organic amendments, According to A. K. Singh R. K. Sharma A. K. and Singh R. C. (2020): “

In present study thus it may be concluded that changes in protein after infection are related to defense action, because abnormal metabolites are produced in adjacent non-infected tissues. Such metabolites accumulated in infected tissues and are toxic to parasites and inhibit their growth and penetration. The metabolites released from the chemical constituents of Neem (*Azadirachtin*, *salannin*, *limonoids*, *triterpenoids*, *phenolic compounds*, *carotenoids*, *steroids* and *ketones*) stimulated the plant cells to release abnormal metabolites which repel the nematodes from the uninfected cells of plant. However, the green leaves are rich in *azadirachtin*, *salannin*, *meliantrol* and *nimbin* (Jacobson, 1990; National Research Council, 1992). So the use of neem products stimulated and changes the physiology of plant cells and tissue to repel the

nematode parasites. Some **important Facts on *Meloidogyne incognita* in Okra (2023–2025)**

- 1) Root-knot nematode (*Meloidogyne incognita*) continues to be a major pest in okra production worldwide**, causing root galling, stunted growth, reduced nutrient and water uptake, and significant yield losses if unmanaged.
- 2) Eco-friendly management strategies are increasingly emphasized** due to environmental and human health concerns associated with chemical nematicides. Sustainable approaches include soil amendments (e. g., neem cake, organic manures), botanicals, and bio-agents.
- 3) Botanical plant materials show promising control potential:** For example, **moringa leaf powder significantly reduced nematode egg and juvenile populations as well as root galling** in okra under pot conditions, showing comparable effects to some nematicides.
- 4) Organic soil amendments**, such as **neem cake or poultry manure**, improve okra growth parameters and help suppress nematode infestation by improving soil health and releasing bioactive compounds that are nematicidal.
- 5) Biological control agents like fungi (*Trichoderma* spp. and *Purpureocillium lilacinum*) and beneficial microbes (*Pseudomonas fluorescens*) significantly reduce nematode reproduction and enhance plant growth** when used as seed treatments or soil inoculants.
- 6) Recent research (2025) on innovative nano-based approaches shows promise** for managing root-knot nematode in okra. Nanocomposites combining silver (AgNP) and zinc oxide (ZnONP) nanoparticles enhanced plant growth and reduced galling and nematode populations, representing a novel sustainable crop protection strategy.
- 7) Genetic resistance is limited in okra**, and commercial resistant cultivars are still rare; thus, integrated management remains crucial. Screening studies have found only moderate resistance in some cultivars.
- 8) Integrated Pest Management (IPM) frameworks are now widely recommended** to combine cultural, biological, and soil-based strategies for long-term sustainable management of plant-parasitic nematodes including *M. incognita*.

**Table 1:** Month wise population dynamics of *Meloidogyne* spp., infesting okra crop in the field study

Serial No.	Months	Temperature (°C)	No. of Population of Nematodes/ 500gm
1	May	40.93±3.08 (36.22-44.03)	1153.60±223.09 (890-1446)
2	June	45.01±3.67 40.22-45.44)	989.80±90.69 (873-1140)
3	September	34.38±1.73 (32.01-37.33)	2281.60±232.11 (1995-2578)
4	October	24.69±1.28 (25.01-28.22)	1380±139.71 (1008-1470)
5	November	20.03±1.27 (20.33-23.03)	1929±532.13 (1098-2474)
6	December	16.87±2.02 (14.33-19.03)	829.20±238.69 (576-1205)
7	January	10.31±3.22 (8-13.33)	1267±140.28 (1065-1431)
8	February	22.85±1.86 (20.01-25.02)	1760±55.82 (1665-1831)

9	March	31.50±2.15 (26.09-32.06)	784.80±153.83 (585-990)
10	April	32.67±3.57 (29.02-38.45)	1819±542.13 (1008-2274)

Suggestion for farmers and other like a gardening lovers suggest some simple ways to manage *Meloidogyne* infestation in okra crops:

- 1) Crop rotation: Rotate okra with non-host crops like cereals or legumes to break the nematode life cycle.
- 2) Soil solarization: Traps soil moisture and heat under a clear plastic tarp, killing nematodes and eggs.
- 3) Organic amendments: Add organic matter like compost, manure, or neem cake to improve soil health and reduce nematode populations.
- 4) Resistant cultivars: Grow okra varieties with some resistance to *Meloidogyne incognita*.
- 5) Nematode-trapping crops: Plant crops like marigold or sesame that attract and trap nematodes.
- 6) Physical barriers: Use physical barriers like plastic or paper mulch to prevent nematodes from reaching okra roots.
- 7) Biofumigation: Plant crops like mustard or broccoli that release nematicidal compounds when incorporated into soil.
- 8) Soil aeration: Improve soil aeration through tillage or mulching to reduce nematode populations. These simple methods can be used alone or in combination with other management strategies to reduce *Meloidogyne* infestation and improve okra crop productivity.

#### Acknowledgement

One of the authors is grateful to the authorities of D. N. College Meerut, and C. C. S. University, Meerut, for providing me the lab facilities in the department. We are also thankful to Dr. Savita Rani, and Dr. Satyender kumar for the valuable suggestion & guidance to complete the research paper.

#### References

- [1] A. K. Singh R. K. Sharma A. K. and Singh R. C. (2020): "Management of *Meloidogyne incognita* on okra using bioagents and organic amendments" Journal of Agricultural Science and Technology, 20: 234-243.
- [2] Archana U. S., and Prasad D (2014). Management of Plant-parasitic Nematodes by the Use of Botanicals. J Plant Physiol Pathol, 2: 1.
- [3] Babu L. and Rana (2012). Efficacy of Neem products as seed treatment against *Meloidogyne incognita* in Okra. Annals of Plant Protection Sciences Volume: 20: 1; 268-269.
- [4] Egunjobi O. H. and Larinde M. A (1975). Nematodes and maize growth in Nigeria 11. Effects of some amendments on the yield of maize (zea mays) in Nigeria. Nematologica Mediteranean.3: 65-73.
- [5] Ganai A. M., Rehman B., Kavita P., Asi M. and Siddiqui M. A (2014). Phytotherapeutic approach for the management of *Meloidogyne incognita* affecting *Abelmoschus esculentus* (L.) Moench. Archives of Phytopathology and Plant Protection 47: 1797- 1805.
- [6] Goswami B. K. and Neetu S (2012). Combat the Soil Borne Fungal and Root Knot nematode Diseases Infecting Horticultural Crops. Amity CJ Bio pest, 5: 255-258.
- [7] Graham R. Stirling and John Bridge and David J. Hunt (2017): "Integrated management of root-knot nematodes on okra using crop rotation and organic amendments" journal of nematology, 90: 456-465. <https://doi.org/10.1186/s41938-023-00645-7>
- [8] Khan M. R., Solanki R D., Bohra B., Vyas B. N (2012). Evaluation of Achook (Azadirachtin 1500ppm) against root knot nematode (*Meloidogyne incognita*) infecting okra. South Asian Journal of Experimental Biology, 2. Management of plant-parasitic nematodes through botanical amendments under protected cultivation. Journal of Biopesticides, 16 (1), 58–65.
- [9] Mishra S. D (1996). Effects of an aqueous extracts of green leaves of Neem, madar and tulsi on the hatching of eggs of root-knot nematode. *Meloidogyne incognita*. Current Nematology.7 (2): 133-135.
- [10] Mohanta S., Mohanty K. C (2012). Biochemical and physiological alteration in okra infected with root-knot nematode, *Meloidogyne incognita*. Journal of Plant Protection and Environment, 9: 54-56. Nematicidal activity of neem (*Azadirachta indica*) leaf extracts against root-knot nematodes. Egyptian Journal of Biological Pest Control, 33, 112.
- [11] R. K. Singh et al (2018) have reported integrated management of root-knot nematode on okra using biological control methods,
- [12] Ram L., Siddiqui A. U., Parihar A. (2009). Management of Root-Knot Nematode, *Meloidogyne incognita* Infecting Okra Using Oil Cakes. Indian Journal of Nematology, 39: 125-127.
- [13] Satyandra S; Rai, A. B. and Sing, R. K (2011). Bio-management of root-knot disease of Chili (*Capsicum annum*) caused by *Meloidogyne incognita*. Vegetable Science, 38 (1): 63-67.
- [14] Sharma, H. K; Satyendraand Pankaj (2007). Management of *Meloidogyne incognita* with *Paecilomyces lilacinus* and Neem Cake on Okra. Pesticide Research Journal 19: 166-168.
- [15] Singh A. K. Singh S. K. and Singh P. K. (2020); "Assessment of *Meloidogyne incognita* population dynamics on okra (*Abelmoschus esculentus*) under different soil amendments" journal of nematology 52 (2): 123-132.
- [16] Singh J. Singh V. K. and Kumar S. (2019): Efficacy of botanicals against *Meloidogyne javanica* on okra" journal crop protection 116: 105-113.
- [17] Singh R. K., Sharma A. K. and Singh R. C. (2022): "Impact of Organic Amendments on *Meloidogyne incognita* Population and Okra Yield" Journal of Environmental Science and Health, Part B 57 91): 12-20.
- [18] Wani, A. N (2006). *Meloidogyne incognita*, on okra and lentil by soil amendment with oil cakes and leaves of different plants. Nematol. medit, 34: 83-87.