

A Study of Results of Seton Therapy as a Treatment for Complex Anal Fistulas: A Case Series

Suyog Bharambe¹, M. M. Anchalia²

¹PG Student, Department of Gen Surgery, Civil Hospital, Ahmadabad, Gujarat, India
²HOD, Department of Gen Surgery, Civil Hospital, Ahmadabad, Gujarat, India

Abstract: *The challenge of surgery for anal fistula is to eradicate the fistula track while maintaining anal continence. Aim of this study was to analyze the influence of seton technique for high anal fistulas, on results of recurrence and postoperative faecal incontinence and need of permanent stomy. A total of 13 patients of complex anal fistula were operated by seton technique over a period of two and half years, with a follow-up of 6months. Of the 13 patients, 3 developed recurrence (22%), with no reported case incontinence to stool or flatus on repeated follow-ups, thus achieving a healing rate of 78%.*

Keywords: complex fistula, seton, incontinence, recurrence, healing rate.

1. Introduction

Complex anal fistula treatment poses a great challenge to the treating surgeon, as it is often associated with recurrences and anal incontinence.

Treatment must be directed by the:

- Etiology
- Anatomy of the fistula
- Degree of symptoms
- Patient co morbidities
- And the surgeon's experience

Keep in mind the progressive tradeoff between:

- Extent of operative sphincter division
- Postoperative healing rates
- And functional compromise

Seton therapy has been used since a very long time for treatment of anal fistulas. Here we prospectively study a case series of complex anal fistulas treated with the *Fistulectomy with Seton Placement* procedure to reduce recurrences and avoid incontinence.

2. Principle

The principle of seton fistulotomy is that any striated muscle that lies superficial to the fistula track is encircled by a piece of a silk thread. The thread is tied tightly and left in situ so that, for a week or two, the striated muscle is slowly divided by a process of ischemic necrosis due to the seton. However, because the process is slow, the muscle does not spring apart, leaving a defect, but heals behind the process of division, so that fibrous tissue is laid down in the bed of the fistula. This process avoids the development of a gutter deformity, which is usually associated with soiling and impaired continence.

3. Methods

A total of 13 patients of complex anal fistula were operated by seton technique over a period of two and half years, with a follow-up of 6months. All of the patients had

crypto glandular anal fistulas, none of them suffered from Crohn's disease or malignancy. 4 patients had history of previous fistula surgery. The operative steps consisted of identification of internal and external openings of the fistula tract, excision of fistula tract, cutting the perianal skin and skin of anal canal and insertion of a cutting silk seton around both internal and external sphincters. A seton thread impregnated with tetracycline was used in the study. Tetracycline had a local antibiotic as well as its fibrosing action augmented sphincter healing. The seton was tightened twice every week till it cut all the way through the sphincter with subsequent healing of the sphincter. Patient was discharged with seton in situ when the fistulotomy wound showed good healing, and was followed twice every week. The seton thread was changed every two weeks, and care was taken that there was no bridging over the fistulotomy wound during healing.

4. Results

No. of cases	No. of recurrences	Ostomies required	No. of incontinence	Healing rate	Mean Healing time
13	3(22%)	0	0	78%	1.5 to 2 months

The fistula was cured in 10 patients (78 percent) with no reports of altered continence at the time of discharge or in the outpatient review. Recurrence developed in 3 patients (22 percent). Long-term follow-up revealed none of the patients suffered from anal incontinence or fecal soiling or required the need for permanent stomy. The average healing time was 1.5 to 2 months.

5. Conclusions

The research data suggest that medicated cutting setons are effective in treating complex fistula-in-ano with regards to preservation of anal function, acceptable recurrence rates and a good quality of life with an additional advantage of reduced healing time and hospital stay.

6. Future Scope

This study had incorporated a medicated seton impregnated with tetracycline which gave healing results better than those in previous mentioned studies. However, due to the small sample size of patients in this study, this conclusion cannot be standardized. A larger study with larger sample size will appropriately quantify and establish medicated seton therapy as one of the most useful therapies for complex fistula management.

7. Discussion and Literature Survey

A lot of studies have been performed on the most ideal method for anal fistula management. The seton therapy has been studied and used for a long time with varying results in different series and settings. Several modifications of seton have also been used. Below are reports detailing incontinence rates after cutting seton treatment for anal fistula in different studies.

Reference	Country	Type of seton	Incontinence
Charua-Guindic et al [7]	Mexico	Silastic	8/50
Cox et al[8]	USA	Silk no 2	4/8
Culp[4]	USA	Penrose drain	3/20
Decanini-Teran et al[9]	Mexico	Silastic tube	0/42
Deshpande et al[5]	India	No 20 chemically treated linen	0/397
Dziki-Bartos[10]	Poland	Rubber band	12/32
Fasth et al[11]	Sweden	Braided silk	0/7
Flichcarbonel et al[12]	Spain	Non absorbable	3/19
Garcia-Aguilar et al[13]	USA	Rubber band	8/12
Gonzalez-Ruiz et al[14]	USA	Silastic vessel loop	0/31
Gurer et al [3]	Turkey	Self locking cable tie	0/17
Hamalainen and Sainio[15]	Finland	Nonabsorbable braided suture	22/35
Hemel et al[16]	Switzerland	Non absorbable braided suture	0/12
Hasegawa et al[17]	UK	Various	15/28
Held et al[18]	USA	Rubber band	0/9
Ho et al[1]	Singapore	Chemical	3/46
Isbister and alsanea[19]	Saudi Arabia	1-0 silk ligature	17/47
Joy and Williams[20]	UK	Silastic	5/10
McCourtney and Finlay[21]	Scotland	Silk ligature	3/16
Mentes et al[22]	Turkey	Elastic	4/20
Misra and Kapur[23]	India	No1 braided stainless steel	0/53
Mohite et al [6]	India	Medicated seton	0/114
Pescatori et al[24]	Italy	NA	3/17
Qureshi et	Pakistan	Silk thread	2/4

al[25]			
Shukla et al[2]	India	Chemical	8/155
Theera pol et al [26]	Singapore	Prolene 0	0/41
Tocchi et al[27]	Italy	NA	4/28
Vatansev et al[28]	Turkey	Cable tie	5/32
Walfisch et al[29]	Isreal	No 0 heavy silk	0/73
Williams e al[30]	USA	No 1 silk	8/13

7.1 Comparisons of Healing Rate with other Treatment Modalities

Method	Healing rate
Anal fistula plug	40 %(15-85%)
Fibrin glue	15 %(10-70%)
Fistulectomy with primary sphincter reconstruction	90 %(80-95%)
Fistulectomy with secondary sphincter reconstruction	95%
Flap procedure	70 %(50-80%)
Long term Seton therapy(This study)	78%

As compared to other modalities of treatment, seton therapy is cheaper, easier to perform and follow-up, having reasonably higher healing rates, with good quality of life in our set-up.

References

- [1] Ho KS, Tsang C, Seow-Choen F et al. Prospective randomized trial comparing ayurvedic cutting seton and fistulotomy for low fistula-in-ano. Tech Coloproctol2001; 5: 137-41.
- [2] Shukla NK, Narang R, Nair NGK et al. Multicentric randomized controlled clinical trial of Kshaarasootra (Ayurvedic medicated thread) in the management of fistula-in-ano. Indian J Med Res 1991; 94: 177-85.
- [3] Gurer A, Ozlem N, GokakinAKetal. Anovel material in seton treatment of fistula-in-ano. Am J Surg2007; 193: 794-6.
- [4] Culp CE. Use of Penrose drains to treat certain anal fistulas: a primary operative seton. Mayo ClinProc1984; 59: 613-7.
- [5] Deshpande PJ, Sharma KR, Sharma SK et al. Ambulatory treatment of fistulo-in-ano: results in 400 cases. Indian JSurg1975; 37: 85-9.
- [6] Mohite JD, Gawai RS, Rohondia OS et al. Ksharsootra (medicated seton) treatment for fistula-in-ano. Indian JGastroenterol1997; 16: 96-7.
- [7] Charu'a-Guindic L, Me'ndez-Mora'n MA, Avendan'o-EspinosaOet al. Seto'n de corte en el tratamiento de la fi'stula anal compleja. Cir Cir2007; 75: 351-6.
- [8] Cox SW, Senagore AJ, Luchtefeld MA et al. Outcome after incision and drainage with fistulotomy for ischioanalabscess. Am Surg1997; 63: 686-9.

- [9] Decanini-Tera'n CO, Galindo GH-R, Belmonte-Montes C et al. Seton de corte lento ¿la respuestaparalafístulasanales? Cir Cir2002; 70: 428–31.
- [10] Dzik A, Bartos M. Seton treatment of anal fistula: experience with a new modification. Eur J Surg1998; 164: 543–8.
- [11] Fasth SB, Nordgren S, Hulte'n L. Clinical course and management of suprasphincteric and extrasphinctericfistula-in-ano. ActaChirScand1990; 156: 397–402. Incontinence rates after cutting seton treatment for anal fistula R. D. Ritchie et al.570 – 2009 The Authors. Journal Compilation _ 2009 The Association oColoproctology of Great Britain and Ireland. Colorectal Disease, 11, 564–571
- [12] FlichCarbonell J, D'ázFons F, Bolufer Cano JM et al. Fístula perianal. Seccio'n de esfínteres con «seto'n». Rev EspEnfermApar Dig 1987; 72: 339–42.
- [13] Garcí'a-Aguilar J, Belmonte C, Wong DW et al. Cutting setonversus two-stage setonfistulotomy in the surgical management of high anal fistula. Br J Surg1998; 85: 243–5.
- [14] Gonzalez-Ruiz C, Kaiser AM, Vukasin P et al. Intraoperative physical diagnosis in the management of anal fistula. AmSurg2006; 72: 11–5.
- [15] Ha'ma'la'inen K-P, Sainio AP. Cutting seton for anal fistulas. High risk of minor control defects. Dis Colon Rectum 1997; 40: 1443–7.
- [16] Hamel CT, Marti WR, Oertli D. Simplified placement and management of cutting setons in the treatment of transsphinctericanal fistula: technical note. Int J Colorectal Dis2004; 19: 354–6.
- [17] Hasegawa H, Radley S, Keighley MRB. Long term results of cutting setonfistulotomy. ActaChirIugosl2000; 47:19–21.
- [18] Held D, Khubchandani I, Sheets J et al. Management of anorectal horseshoe abscess and fistula. Dis Colon Rectum1986; 29: 793–7.
- [19] Isbister WH, Al Sanea N. The cutting seton. An experience at King Faisal Specialist Hospital. Dis Colon Rectum 2001; 44:722–7.
- [20] Joy HA, Williams JG. The outcome of surgery for complex anal fistula. Colorectal Dis 2002; 4: 254–61.
- [21] McCourtney JS, Finlay IG. Cutting seton without preliminary internal sphincterotomy in management of complex high fistula-in-ano. Dis Colon Rectum 1996; 39: 55–8.
- [22] Menten BB, O' ktemer S, Tezcaner T et al. Elastic one-stage cutting seton for the treatment of high anal fistulas: preliminary results. Tech Coloproctol2004; 8: 159–62.
- [23] Misra MC, Kapur BML. A new non-operative approach to fistula in ano. Br J Surg1988; 75: 1093–4.
- [24] Pescatori M, AyabacaS, Caputo D. Can anal manometrypredict anal incontinence after fistulectomy in males? ColorectalDis 2004; 6: 97–102.
- [25] Qureshi KH, Kamal M, Shah HA et al. Management of fistula-in-ano – a common clinical problem. Pak J Med Res2002; 41: 113–7.
- [26] Theerapol A, So BYJ, Ngoi SS. Routine use of setons for the treatment of anal fistulae. Singapore Med J 2002; 43:305–7.
- [27] Tocchi A, Mazzoni G, Lepre L, Costa G, Liotta G, Maggiolini F. Fistoleanalialterecidive: trattamento consetone. G Chir1997; 18: 375–7.
- [28] Vatansev C, Alabaz O', Tekin A et al. A new seton type for the treatment of anal fistula. Dig Dis Sci2007; 52: 1920–3.
- [29] Walfisch S, Menachem Y, Koretz M. Double seton— a new modified approach to high transsphincteric anal fistula. Dis Colon Rectum 1997; 40: 731–2.
- [30] Williams JG, MacLeod CA, Rothenberger DA et al. Seton treatment of high anal fistulae. Br J Surg1991; 78: 1159–61.

Author Profile



Dr. Suyog Anil Bharambe is a 3rd year Resident in Department of Gen Surgery, Civil Hospital, Ahmadabad. Gujarat, India.



Dr. M. M. Anchalia (M.S. Gen. Surgery, FIAGS). She is HOD of Department of General Surgery in B.J.Medical college and Civil hospital, Ahmadabad. Gujarat, India. She is also P.G. Director in B.J.Medical college and Civil hospital, Ahmadabad. Gujarat, India.