

# On Fixed Point Theorem in Fuzzy 2-Metric Spaces for Integral type Inequality

Rasik M. Patel<sup>1</sup>, Ramakant Bhardwaj<sup>2</sup>

<sup>1</sup>Research Scholar, CMJ University, Shillong, Meghalaya, India

<sup>2</sup>Truba Institute of Engineering & Information Technology, Bhopal, M.P., India

rasik\_maths@yahoo.com  
drkkbhardwaj100@gmail.com

**Abstract:** In the present paper we are proving a common fixed point theorem for fuzzy 2-metric spaces for rational expressions. 2010 Mathematics Subject Classification: Primary 47H10, 54H25.

**Keywords:** Fuzzy metric space, Fuzzy 2-metric space, Common fixed point.

## 1. Introduction and Preliminaries

Impact of fixed point theory in different branches of mathematics and its applications is immense. The first result on fixed points for contractive type mapping was the much celebrated Banach's contraction principle by S. Banach [37] in 1922. In the general setting of complete metric space, this theorem runs as the follows, Theorem 1.1(Banach's contraction principle) Let  $(X, d)$  be a complete metric space,  $c \in (0, 1)$  and  $f: X \rightarrow X$  be a mapping such that for each  $x, y \in X$ ,  $d(fx, fy) \leq c d(x, y)$ . Then  $f$  has a unique fixed point  $a \in X$ , such that for each  $x \in X$ ,  $\lim_{n \rightarrow \infty} f^n x = a$ . After the classical result, R.Kannan [31] gave a subsequently new contractive mapping to prove the fixed point theorem. Since then a number of mathematicians have been worked on fixed point theory dealing with mappings satisfying various type of contractive conditions. In 2002, A. Branciari [1] analyzed the existence of fixed point for mapping  $f$  defined on a complete metric space  $(X, d)$  satisfying a general contractive condition of integral type.

Theorem 1.2(Branciari) Let  $(X, d)$  be a complete metric space,  $c \in (0, 1)$  and let  $f: X \rightarrow X$  be a mapping such that for each  $x, y \in X$ ,  $\int_0^{d(fx, fy)} \varphi(t) dt \leq c \int_0^{d(x, y)} \varphi(t) dt$

Where  $\varphi: [0, +\infty) \rightarrow [0, +\infty)$  is a Lebesgue integrable mapping which is summable on each compact subset of  $[0, +\infty)$ , non negative, and such that for each  $\varepsilon > 0$ ,  $\int_0^\varepsilon \varphi(t) dt > 0$ , then  $f$  has a unique fixed point  $a \in X$  such that for each  $x \in X$ ,  $\lim_{n \rightarrow \infty} f^n x = a$ . After the paper of Branciari, a lot of a research works have been carried out on generalizing contractive condition of integral type for a different contractive mapping satisfying various known properties. A fine work has been done by B.E. Rhoades [2] extending the result of Branciari by replacing the condition [1.2] by the following;

$$\int_0^{d(fx, fy)} \varphi(t) dt \leq \int_0^{\max\{d(x, y), d(x, fx), d(y, fy), \frac{d(x, fy) + d(y, fx)}{2}\}} \varphi(t) dt \quad (1.3)$$

The aim of this paper is to generalize some mixed type of contractive conditions to the mapping and then a pair of mappings, satisfying a general contractive mapping such

as R. Kannan type [31], S.K. Chatterjee type [36], T. Zamfirescu type [38], Turkoglu [39] etc. In 1965, the concept of fuzzy sets was introduced by Zadeh [42]. After that many authors have expansively developed the theory of fuzzy sets and applications. Especially, Deng [10], Erceg [12], Kaleva and Seikkala [25, 26], Kramosil and Michalek [27], have introduced the concept of fuzzy metric spaces in different ways. Recently, many authors [3-5, 13, 19, 22, 23, 24, 29, 30, 34, and 35] have also studied the fixed point theory in the fuzzy metric spaces and [6-9, 21, 28,] have studied for fuzzy mappings which opened an avenue for further development of analysis in such spaces and such mappings. Consequently in due course of time some metric fixed point results were generalized to fuzzy metric spaces by various authors. Gähler in a series of papers [15, 16, and 17] investigated 2-metric spaces. Sharma, Sharma and Iseki [33] studied for the first time contraction type mappings in 2-metric space. We [40, 41] have also worked on 2-Metric spaces and 2-Banach spaces for rational expressions.

**Definition 1.1:** A binary operation  $*$ :  $[0, 1]^3 \rightarrow [0, 1]$  is called a continuous t-norm if  $([0, 1], *)$  is an abelian topological monoid with unit 1 such that  $a_1 * b_1 * c_1 \geq a_2 * b_2 * c_2$  whenever  $a_1 \geq a_2, b_1 \geq b_2, c_1 \geq c_2$  for all  $a_1, b_1, c_1, a_2, b_2, c_2$  are in  $[0, 1]$ .

**Definition 1.2:** the 3-tuple  $(X, M, *)$  is called a fuzzy 2-metric space if  $X$  is an arbitrary set,  $*$  is a continuous t-norm and  $M$  is a fuzzy set in  $X^3 \times [0, \infty)$  satisfying the following conditions: for all  $x, y, z, u \in X$  and  $t_1, t_2, t_3 > 0$ ,

- (1)  $M(x, y, z, 0) = 0$ ,
- (2)  $M(x, y, z, t) = 1$  for all  $t > 0$  (only when the three simplex  $x, y, z$  degenerate)
- (3)  $M(x, y, z, t) = M(x, z, y, t) = M(y, z, x, t)$   
=.....
- (4)  $M(x, y, z, w, t_1, t_2, t_3) \geq *(M(x, y, u, t_1)) * M(x, u, z, t_2) * M(u, y, z, t_3)$
- (5)  $M(x, y, z, \cdot): [0, 1] \times [0, 1]$  is left continuous.

**Definition 1.3:** Let  $(X, M, *)$  be a fuzzy- 2 metric space.

(1) A sequence  $\{x_n\}$  in fuzzy -2 metric space  $X$  is said to be convergent to a point  $x \in X$  (denoted by

$$\lim_{n \rightarrow \infty} x_n = x \text{ or } x_n \rightarrow x$$

if for any  $\lambda \in (0,1)$  and  $t > 0$ , there exists  $n_0 \in \mathbb{N}$  such that for all  $n \geq n_0$  and  $a \in X, M(x_n, x, a, t) > 1 - \lambda$ .

That is  $\lim_{n \rightarrow \infty} M(x_n, x, a, t) = 1$  for all  $a \in X$  and  $t > 0$ .

(2) A sequence  $\{x_n\}$  in fuzzy- 2 metric space  $X$  is called a Cauchy sequence, if for any  $\lambda \in (0,1)$  and  $t > 0$ , there exists

$n_0 \in \mathbb{N}$  such that for all  $m, n \geq n_0$  and  $a \in X, M(x_n, x_m, a, t) > 1 - \lambda$ .

(3) A fuzzy- 2 metric space in which every Cauchy sequence is convergent is said to be complete.

**Definition 1.4:** Self A function  $M$  is continuous in fuzzy 2-metric space if  $x_n \rightarrow x, y_n \rightarrow y$ , then

$$\lim_{n \rightarrow \infty} M(x_n, y_n, a, t) = M(x, y, a, t) \text{ for all } a \in X \text{ and } t > 0.$$

**(3.1a)**

$$\int_0 M(Ax, Ay, a, qt) \zeta(t) dt \geq \int_0 \left\{ \min \left\{ \begin{array}{l} M(Ty, Ay, a, t), M(Sx, Ax, a, t), \\ M(Sx, Ty, a, t), \frac{M(Sx, Ty, a, t)}{M(Ax, Ty, a, t)} \\ \frac{M(Ax, Sx, a, t)}{M(Ay, Sx, a, t)}, M(Ax, Ty, a, t), \end{array} \right. \right\} \zeta(t) dt$$

For all  $x, y, a \in X, t > 0$  and  $q \in (0,1)$

**(3.1b)**

$$\lim_{n \rightarrow \infty} M(x, y, z, t) = 1 \text{ for all } x, y, z \in X.$$

Then  $F, T$  and  $A$  have a unique common fixed point in  $X$ .

Proof:

We define a sequence  $\{x_n\}$  in  $X$  such that  $Ax_{2n} = Sx_{2n-1}$  and  $Ax_{2n-1} = Tx_{2n}$  for  $n = 1, 2, \dots$

We shall prove that  $\{Ax_n\}$  is a Cauchy sequence. For this suppose  $x = x_{2n}$  and  $y = x_{2n+1}$  in (3.1a), we write

**Definition 1.5:** Two mappings  $A$  and  $S$  on fuzzy 2-metric space  $X$  are weakly commuting if

$$M(ASu, SAu, a, t) \geq M(Au, Su, a, t) \text{ for all } a, u \in X \text{ and } t > 0.$$

## 2. Some Basic Results

**Lemma 2.1:** [19] for all  $x, y \in X, M(x, y)$  is non-decreasing.

**Lemma 2.2:** [9] Let  $\{y_n\}$  be a sequence in a fuzzy metric space  $(X, M, *)$  with condition (FM-6) If there exists a number  $q \in (0,1)$  such that  $M(y_{n+2}, y_{n+1}, qt) \geq M(y_{n+1}, y_n, t)$  for  $t > 0$  and  $n = 1, 2, 3, \dots$  then  $\{y_n\}$  is a Cauchy sequence in  $X$ .

**Lemma 2.3:** [30] for all  $x, y \in X$  and for a number  $q \in (0,1)$  such that  $M(x, y, qt) \geq M(x, y, t)$  for  $t > 0$  then  $x = y$ .

**Lemma** (2.1, 2.2, and 2.3) are also true for fuzzy2-metric spaces.

## 3. Main Result

**Theorem 3.1:** Let  $(X, M, *)$  be a complete fuzzy 2-metric space and let  $S$  and  $T$  be continuous mappings of  $X$  in  $X$ . then  $S$  and  $T$  have a common fixed point in  $X$  if there exists continuous mapping  $A$  of  $X$  into  $S(X) \cap T(X)$  which commute weakly with  $S$  and  $T$ ,

$$\int_0^{M(Ax_{2n}, Ax_{2n+1}, a, qt)} \zeta(t) dt \geq \int_0^{\left\{ \min \left\{ \frac{M(Tx_{2n+1}, Ax_{2n+1}, a, t), M(Sx_{2n}, Ax_{2n}, a, t), M(Sx_{2n}, Tx_{2n+1}, a, t)}{M(Sx_{2n}, Tx_{2n+1}, a, t)}, \frac{M(Ax_{2n}, Sx_{2n}, a, t)}{M(Ax_{2n}, Tx_{2n+1}, a, t)}, M(Ax_{2n}, Tx_{2n+1}, a, t) \right\}} \right\}} \zeta(t) dt$$

$$\begin{aligned} \int_0^{M(Ax_{2n}, Ax_{2n+1}, a, qt)} \zeta(t) dt &\geq \int_0^{\left\{ \min \left\{ \frac{M(Ax_{2n}, Ax_{2n+1}, a, t), M(Ax_{2n+1}, Ax_{2n}, a, t), M(Ax_{2n+1}, Ax_{2n}, a, t)}{M(Ax_{2n+1}, Ax_{2n}, a, t)}, \frac{M(Ax_{2n}, Ax_{2n+1}, a, t)}{M(Ax_{2n}, Ax_{2n}, a, t)}, M(Ax_{2n}, Ax_{2n}, a, t) \right\}} \right\}} \zeta(t) dt \\ &= \int_0^{\left\{ \min \left\{ \frac{M(Ax_{2n}, Ax_{2n+1}, a, t), M(Ax_{2n+1}, Ax_{2n}, a, t), M(Ax_{2n+1}, Ax_{2n}, a, t)}{M(Ax_{2n+1}, Ax_{2n}, a, t)}, M(Ax_{2n}, Ax_{2n+1}, a, t), 1 \right\}} \right\}} \zeta(t) dt \\ &\geq \int_0^{\left\{ \min \left\{ M(Ax_{2n-1}, Ax_{2n}, a, \frac{t}{q}), M(Ax_{2n}, Ax_{2n-1}, a, \frac{t}{q}) \right\}} \right\}} \zeta(t) dt \end{aligned}$$

Therefore

$$\int_0^{M(Ax_{2n}, Ax_{2n+1}, a, qt)} \zeta(t) dt \geq \int_0^{\left\{ \min \left\{ M(Ax_{2n-1}, Ax_{2n}, a, \frac{t}{q}) \right\}} \right\}} \zeta(t) dt$$

By induction

$$\int_0^{M(Ax_{2k}, Ax_{2m+1}, a, qt)} \zeta(t) dt \geq \int_0^{\left\{ \min \left\{ M(Ax_{2m}, Ax_{2k-1}, a, \frac{t}{q}) \right\}} \right\}} \zeta(t) dt$$

for every k and m in N, further if 2m+1 > 2k, then

$$\begin{aligned} \int_0^{M(Ax_{2k}, Ax_{2m+1}, a, qt)} \zeta(t) dt &\geq \int_0^{\left\{ M(Ax_{2k}, Ax_{2m}, a, \frac{t}{q}) \right\}} \zeta(t) dt \\ &\dots\dots\dots \\ &\geq \int_0^{\left\{ M(Ax_0, Ax_{2m+1}, a, \frac{t}{2^k}) \right\}} \zeta(t) dt \dots\dots(3.1c) \end{aligned}$$

If 2k > 2m+1, then

$$\int_0^M(Ax_{2k}, Ax_{2m+1}, a, qt) \zeta(t) dt \geq \int_0^{\left\{ M(Ax_{2k-1}, Ax_{2m}, a, \frac{t}{q}) \right\}} \zeta(t) dt$$

.....

$$\geq \int_0^{\left\{ M(Ax_{2k-(2m+1)}, Ax_0, a, \frac{t}{q^{2m+1}}) \right\}} \zeta(t) dt \dots\dots(3.1d)$$

By simple induction with (3.1c) and (3.1d) we have

$$\int_0^M(Ax_n, Ax_{n+p}, a, qt) \zeta(t) dt \geq \int_0^{\left\{ M(Ax_0, Ax_p, a, \frac{t}{q^n}) \right\}} \zeta(t) dt$$

For  $n = 2k, p = 2m+1$  or  $n = 2k+1, p = 2m+1$

$$\int_0^M(Ax_n, Ax_{n+p}, a, qt) \zeta(t) dt \geq \int_0^{\left\{ M(Ax_0, Ax_1, a, \frac{t}{2q^n}) \right\} * \left\{ M(Ax_1, Ax_p, a, \frac{t}{q^n}) \right\}} \zeta(t) dt \dots\dots(3.1e)$$

If  $n = 2k, p = 2m$  or  $n = 2k+1, p = 2m$ ,

For every positive integer  $p$  and  $n$  in  $\mathbb{N}$ , by noting that

$$\int_0^{\left\{ M(Ax_0, Ax_p, a, \frac{t}{q^n}) \right\}} \zeta(t) dt \rightarrow 1 \text{ as } n \rightarrow \infty$$

Thus  $\{Ax_n\}$  is a Cauchy sequence. Since the space  $X$  is complete there exists  $z \in X$ , such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_{2n-1} = \lim_{n \rightarrow \infty} Tx_{2n} = z.$$

It follows that  $Az = Sz = Tz$  and Therefore

$$\int_0^M(Az, AAz, a, qt) \zeta(t) dt \geq \int_0^{\left\{ \min \left\{ \begin{array}{l} M(TAz, AAz, a, t), M(Sz, Az, a, t), \\ M(Sz, TAz, a, t), \frac{M(Sz, TAz, a, t)}{M(Az, TAz, a, t)}, \\ \frac{M(Az, Sz, a, t)}{M(AAz, Sz, a, t)}, M(Az, TAz, a, t), \end{array} \right\} \right\}} \zeta(t) dt$$

$$\begin{aligned} \int_0^t M(Az, A^2z, a, qt) \zeta(t) dt &\geq \int_0^t \{M(Sz, TAz, a, t)\} \zeta(t) dt \\ &\geq \int_0^t \{M(Sz, ATz, a, t)\} \zeta(t) dt \\ &\geq \int_0^t \{M(Az, A^2z, a, t)\} \zeta(t) dt \\ &\geq \int_0^t \left\{M\left(Az, A^2z, a, \frac{t}{q^n}\right)\right\} \zeta(t) dt. \end{aligned}$$

Since,

$$\lim_{n \rightarrow \infty} M\left(Az, A^2z, a, \frac{t}{q^n}\right) = 1 \rightarrow Az = A^2z$$

Thus z is common fixed point of A, S and T.

For uniqueness, let v (v ≠ z) be another common fixed point of S, T and A, By (3.1a) we write

$$\int_0^t M(Az, Av, a, qt) \zeta(t) dt \geq \int_0^t \left\{ \min \left\{ \begin{array}{l} M(Tv, Av, a, t), M(Sz, Az, a, t), \\ M(Sz, Tv, a, t), \frac{M(Sz, Tv, a, t)}{M(Az, Tv, a, t)}, \\ \frac{M(Az, Sz, a, t)}{M(Av, Sz, a, t)}, M(Az, Tv, a, t), \end{array} \right\} \right\} \zeta(t) dt$$

$$\int_0^t M(Az, Av, a, qt) \zeta(t) dt \geq \int_0^t \{M(z, v, a, t)\} \zeta(t) dt$$

This implies that

$$\int_0^t M(z, v, a, qt) \zeta(t) dt \geq \int_0^t \{M(z, v, a, t)\} \zeta(t) dt$$

#### 4. Acknowledgement

Dr. Ramakant Bhardwaj, Associate Professor and PI of MPCOST Bhopal for the project (Application of Fixed Point Theory), Department of Mathematics, Truba Institute of Engineering & Information Technology, Bhopal, M.P, India

#### References

- [4] A.Branciari, A fixed point theorem for mappings satisfying a general contractive condition of integral type, Int.J.Math.Sci. 29(2002), no.9, 531 - 536.
- [5] B.E. Rhoades, Two fixed point theorems for mappings satisfying a general contractive condition of integral type. International Journal of Mathematics and Mathematical Sciences, 63, (2003), 4007 - 4013
- [6] Badard, R. "Fixed point theorems for fuzzy numbers" Fuzzy sets and systems 13 (1984) 291-302
- [7] Bose, B.K. and Sahani, D. "Fuzzy mappings and fixed point theorems" Fuzzy sets and Systems 21 (1984) 53-58.
- [8] Butnariu, D. "Fixed point for fuzzy mappings" Fuzzy sets and Systems 7(1982) 191-207
- [9] Chang, S.S. "Fixed point theorems for fuzzy mappings" Fuzzy Sets and Systems 17(1985) 181-187.
- [10] Chang, S.S., Cho, Y.J., Lee, B.S. and Lee, G.M. "Fixed degree and fixed point theorems for fuzzy mappings" Fuzzy Sets and Systems 17(1985) 181-187.

- mappings" *Fuzzy Sets and Systems* 87 (1997) 325-334
- [11] Change, S.S., Cho, Y.J., Lee B.S., June, J.S. and Kang, S.M. "Coincidence point and minimization theorems in fuzzy metric spaces" *Fuzzy Sets and Systems*, 88 (1) (1997) 119-128
- [12] Cho, Y.J. "Fixed points and fuzzy metric spaces" *J. Fuzzy Math.*5 (1997) no. 4, 949-962.
- [13] Deng, Z. "Fuzzy pseudo-metric space" *J. Math, Anal, Appl.*86 (1982) 74-95.
- [14] Eklund, I. and Gähler, S. "Basic notions for fuzzy topology" *Fuzzy Sets and System* 26 (1988) 336-356.
- [15] Erceg, M.A "Metric space in fuzzy set theory" *J. Math, Anal, Appl.*69 (1979) 205-230.
- [16] Fang, J.X. "On fixed point theorems in fuzzy metric spaces" *Fuzzy Sets and Systems* 46 (1979) 107-113
- [17] Fisher, B. "Mappings with a common fixed point" *Math. Seminar notes Kobe university* 7 (1979) 81-84.
- [18] Gähler, S. *Linear 2-normierte Raume*" *Math. Nachr.* 28 (1964) 1-43.
- [19] Gähler, S. *Über 2-Banach Raume*, *Math. Nachr.* 42 (1969) 335-347
- [20] Gähler, S. *2- metrics Raume and ihre topologische structure*, *Math. Nachr.*26 (1983) 115-148.
- [21] George, A. and Veramani, P."On some results in fuzzy metric spaces" *Fuzzy Sets and Systems* 64(1994)395-399.
- [22] Grabiec, M. "Fixed points in fuzzy metric space" *Fuzzy Sets and Systems* 27 (1988) 385-389.
- [23] Hu, C. "Fuzzy topological space" *J. Math. Anal. Appl.* 110(1985)141-178.
- [24] Helpert, S. "Fuzzy mappings and fixed point theorems" *J. Math. Anal. Appl.*83 (1981) 566-569.
- [25] Hadzic, O. "Fixed point theorems, for multi-valued mappings in some classes of fuzzy metric space" *Fuzzy Sets and Systems* 29(1989) 115-125.
- [26] Jung, J.S., Cho. Y.J. and Kim J.K.: *Minimization theorems for fixed point theorems in fuzzy metric spaces and applications*, *Fuzzy Sets and Systems* 61 (1994) 199-207.
- [27] Jung, J.S., Cho, Y.J., Chang, S.S. and Kang, S.M. "Coincidence theorems for set valued mappings and Eklund's variational principle in fuzzy metric spaces" *Fuzzy Sets and Systems* 79 (1996) 239-250.
- [28] Kaleva, O. and Seikkala, S. "On Fuzzy metric spaces" *Fuzzy Sets and Systems* 12 (1984) 215-229.
- [29] Kaleva, O. "The Completion of fuzzy metric spaces" *J. Math. Anal. Appl.* 109(1985) 194-198.
- [30] Kramosil, I. and Michalek, J. "Fuzzy metric and statistical metric spaces" *Kybernetika* 11 (1975) 326-334.
- [31] Lee, B.S., Cho, Y.J. and Jung, J.S. "Fixed point theorems for fuzzy mappings and applications" *Comm. Korean Math. Sci.* 11 (1966) 89-108.
- [32] Kutukcu, S., Sharma, S. and Tokgoz, H. "A Fixed point theorem in fuzzy metric spaces" *Int. Journal of Math. Analysis* 1 (2007) no.18, 861-872.
- [33] Mishra, S.N., Sharma, N. and Singh, S.L. "Common fixed points of maps on fuzzy metric spaces" *Internet. J. Math. & Math Sci.*17 (1966) 89-108.
- [34] R. Kannan, *Some results on fixed points*, *Bull. Calcutta Math. Soc.*, 60(1968), 71-76.
- [35] Schweitzer, B. and Sklar, A. "Statistical metric spaces" *Pacific Journal Math.*10 (1960) 313-334.
- [36] Sharma. P.L., Sharma. B.K. and Iseki, K. "Contractive type mapping on 2- metric spaces" *Math. Japonica* 21 (1976) 67-70.
- [37] Sharma, S. "On fuzzy Metric space" *Southeast Asian Bulletin of Mathematics* 26 (2002) 133-145.
- [38] Som, T. "Some results on common fixed point in fuzzy Metric spaces" *Soochow Journal of Mathematics* 4 (2007) 553-561.
- [40] S.K.Chatterjea, *Fixed point theorems*, *C.R.Acad.Bulgare Sci.* 25(1972), 727-730.
- [41] S.Banach, *Sur les oprations dans les ensembles abstraits et leur application aux quations intgrales*, *Fund. Math.*3,(1922)133-181 (French).
- [42] T.Zamfrescu, *Fixed point theorems in metric spaces*, *Arch. Math. (Basel)* 23(1972), 292-298
- [43] Turkoglu, D and Rhodes, B.E. "A fixed fuzzy point for fuzzy mapping in complete metric spaces" *Mathematical Communications*10 (2005)115-121.
- [44] Yadava, R.N., Rajput, S.S. Choudhary, S. and Bhardwaj, R.K "Some fixed point theorems for rational inequality in 2- metric spaces" *Acta Ciencia Indica* 33 (2007) 709-714.
- [45] Yadava, R.N., Rajput, S.S. Choudhary, S. and Bhardwaj, R.K. "Some fixed point theorems for non contraction type mapping on 2- Banach spaces" *Acta Ciencia Indica*, 33: 737-744 (2007).
- [46] Zadeh, L.A. "Fuzzy Sets" *Information and control* 8 (1965) 338-353.