

Figure 2: DNA Polymorphism Observed In Maize Hybrid (DHM-117) And Parental Lines Using SSR Markers Phi053 And Umc1600




Figure 3: Testing the genetic purity of 20 randomly selected plants from BML 6, representing the off types




Figure 4: Testing the genetic purity of 20 randomly selected plants from BML 7, representing the off types




Figure 5: Testing the genetic purity of 20 randomly selected plants from DHM 117 showing 100 per cent genetic purity




Figure 6: Testing the genetic purity 20 randomly selected plants from in DHM 117, representing the off types

## References

- [1] Cooke, R.J. (1995). Gel electrophoresis for the identification of plant varieties. *J. of Chrom.* 698(1): 281-299.
- [2] Ichii, M. Hong, D.L. Ohara, Y. Zhao, C .M. and Taketa, S. (2003). Characterization of CMS and maintainer lines in indica rice (*Oryza sativa* L.) based on RAPD marker analysis. *Euphy.* 129: 249-252.
- [3] Kumar, L.S. (1999). DNA markers in plant improvement: an over view. *Biotechnol. Adv.* 17: 143-182.
- [4] Lubberstedt, A.E. Melchinger, C. Dussle, M. Vuylsteke and Kuiper, M. (2000). Relations among early European maize inbreds: IV. Genetic diversity revealed with AFLP markers and comparison with RFLP, RAPD and pedigree data. *Crop Sci.*40: 783-791.
- [5] Maccaferri, M. Stefanelli, S. Rotondo, F. Tuberosa, R. and Sanguineti, M.C. (2007). Relationships among durum wheat accessions. I. Comparative analysis of SSR, AFLP, and phenotypic data. *Gen.* 50(4): 373-384.
- [6] Murray, M.G. and Thompson, W.F. (1980). Rapid isolation of high molecular weight. *DNA. Nucleic acids Res.* 8: 4321-4325.
- [7] Nandakumar, N. Singh, A. Sharma, K. Mohapara, R.K. Prabhu, T.K.V and Zaman, F.U. (2004). Molecular fingerprinting of Hybrids and assessment of genetic purity of hybrid seeds in rice using microsatellite markers. *Euphy.* 136: 257-264.
- [8] Paniego, N. Echaide, M. Munoz, M. Fernandez, L. Torales, S. Faccio, P. Fuxan, I. Crrera, M. Zandomeni, R. Syarez, E.Y. and Hopp, E. (2002). Microsatellite isolation and characterization in sunflower (*Helianthus annuus* L.). *Genome* 45: 34-43.

- [9] Scanlon, M.G. Sapirstein, H.D. and Bhushuk, W. (1989). Computerized wheat varietal identification by high-performance liquid chromatography. *Cereal Chem.* 66: 439-4443.
- [10] Sharopova, N. McMullen, M.D. Schultz, L. Schreoder, S. Sanchez- Villeda, H. Gardiner, J. Bergstrom, D. Houchins, K. Melia- Hancock, S. and Musket, T. (2002). Development and mapping of SSR markers for maize. *Plant Mol.Biol.* 48(5-6): 463-48.
- [11] Smith, J.S.C. Chin, E.C.L. Shu, H. Smith, O.S. Wall, S.J. Senior, M.L. Mitchell, S.E. Kresorich, S. and Ziegler, J. (1997). An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L.). Comparison with data from RFLP and Pedigree. *Theor.Appl. Genet.* 95(1):163-173.
- [12] Sundaram, R.M. Naveenkumar, B. Biradar, S.K. Balachandran, S.M. Mishra, B. Ilyasahmed, M. Viraktamath, B.C. Ramesha, M.S. and Sarma, N.P. (2008). Identification of informative SSR markers capable of distinguishing hybrid rice parental lines and their utilization in seed purity assessment. *Euphy.* 163: 215-224
- [13] Tkachuk, R. and Mellish, V.J. (1980). Wheat cultivars identification by high voltage electrophoresis. *Ann.Technol. Agric.* 29: 207-212.
- [14] Wang, J. Zhong, G.Y. Chin, E.C.L. Register, III J.C. Riley, R.D. Nicbar, W.S. Smith, J.S.C. (2002). Identification of parents of F<sub>1</sub> hybrids through SSR profiling of maternal and hybrid tissue. *Euphy.* 124:29-34.
- [15] Williams, J.G.K. Kubelik, A.R. Leavk, K.J. Rafaski, J.A. and Tingey, S.V. (1990). DNA polymorphism amplification by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18: 6531-6535.
- [16] Yamagishi, M. (1995). Detection of section specific random amplified polymorphic DNA (RAPD) markers in *Lilium*. *Theor. Appl. Genet.* 91: 830-835.
- [17] Yashitola, J. Thirumurgan, T. Sundaram, R.M. Naseerullah, M.K. Ramesha, M.S. Sarma, N.P. and Santi, R.V. (2002). Assessment of purity of rice hybrid using microsatellite and STS markers. *Crop Sci.* 42: 1369-1373.
- [18] Yu, Y. Yuan, D. Liang, S. Li, X. Wang, X. Lin, Z. and Zhang, X. (2011). Genome structure of cotton revealed by a genome- wide SSR genetic map constructed from a BC<sub>1</sub> population between *Gossypium hirsutum* and *G. barbadense*. *BMC Genomics.* 12:15.
- [19] Zilman, R.R. and Bushuk, W. (1979). Wheat cultivars identification by glidinelectrophore-grams.III. Catalogue of electrophoregram formulas of Canadian wheat cultivars. *Can. J. Plant Sci.* 59: 287-298.



**Sambasiva Rao Palaparthy** is presently working as Professor, Department of Crop Physiology, Water Technology Centre, College of Agriculture, Acharya NG Ranga Agricultural University, Rajendranagr, Hyderabad, India. His research areas include Plant water relations of Aerobic rice varieties and water stress management.

**LV Subba Rao** is working as Principal Scientist (Plant Breeding), Directorate of Rice Research, Rajendranagar, Hyderabad, India.

## Author Profile



**Sudharani Madugula** is Senior Scientist (Plant Breeding), Seed Research & Technology Centre, ANGRAU, Rajendranagar, Hyderabad, India. She has 15 years research experience and involved in developing 2 rice varieties for salinity and gall midge tolerance. Currently working in Seed Physiology, Storage and Testing. She developed 15 Seed technologies and disseminated to farmers and she has more than 30 research publications and 20 popular articles to her credit.