# Location and Stability of Equilibrium Points in a PRTBP under Perturbations, Bigger Primary is an **Oblate Spheroid**

Dr. M. N. Haque<sup>1</sup>, Vikash Kumar<sup>2</sup>

<sup>1, 2</sup>M S College, B.R.A. Bihar University, Muzaffarpur, India

Abstract: We have examined the location and linear stability of triangular points in photogravitational restricted three body problems, bigger primary is an oblate spheroid. We observe that the triangular points  $L_4$ ,  $L_5$  form equilateral triangle with the primaries and are linearly stable if the mass ratio  $\mu$  is less than the critical mass value  $\mu_c = 0.0385209 \dots \dots$ 

Keywords: PRTBP, Equilateral Triangle, Triangular Points, Critical mass

#### 1. Introduction

The present work "Location and stability of equilibrium points in a Photogravitational restricted three body problem, bigger primary is an oblate spheroid" deals with the equation of motion and the triangular stability. Here, both the primaries are taken as sources of radiation and the bigger primary as an oblate spheroid. The perturbations in coriolis and centrifugal forces are taken into consideration.

#### **(A)** Equations of motion and Location of triangular points:-

The equations of motion of a perturbed photogravitational restricted three body problem in bigger primary is an oblate spheroid is obtained as:

$$\ddot{\mathbf{x}} - 2\alpha \dot{\mathbf{y}} = \frac{\partial \Omega}{\partial \mathbf{x}}$$
$$\ddot{\mathbf{y}} + 2\alpha \dot{\mathbf{x}} = \frac{\partial \Omega}{\partial \mathbf{y}}$$
where  $(1)$ 

$$\Omega = \frac{1}{2} (x^{2} + y^{2})\beta + \frac{1}{1 + \frac{3A}{2}} \left\{ \frac{(1 - \mu)q_{1}}{r_{1}} + \frac{\mu q_{2}}{r_{2}} + \frac{(1 - \mu)q_{1}A}{2r_{1}^{3}} \right\} \dots \dots$$
(2)  
and

$$\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\end{array}\\
\end{array} \\ q_{1} = 1 - \delta_{1}; \ 0 < \delta_{1} < 1 \\ \\
\end{array} \\ q_{2} = 1 - \delta_{2}; \ 0 < \delta_{2} < 1 \\ \\
\end{array} \\ \vdots \\ \alpha = 1 + \epsilon_{1}; |\epsilon_{1}| \ll 1 \\ \beta = 1 + \epsilon_{2}; |\epsilon_{2}| \ll 1 \\ \\
\end{array}$$
(3)
$$\begin{array}{c}
\end{array} \\ \beta = 1 + \epsilon_{2}; |\epsilon_{2}| \ll 1 \\ \\
\end{array}$$
and

$$r_1^2 = (x + \mu)^2 + y^2 r_2^2 = (x - 1 + \mu)^2 + y^2$$
 ...... (4)

The coordinates of the triangular equilibrium points, upto the first order terms in the parameters  $\epsilon_2$ ,  $q_1$ ,  $q_2$  and A are:

$$\begin{aligned} x &= \frac{1}{2} - \mu - \frac{1}{3}\delta_1 + \frac{1}{3}\delta_2 + \frac{1}{2}A + \frac{1}{3}A\delta_1 - \frac{7}{6}A\delta_2 \\ y &= \pm \frac{\sqrt{3}}{2}(1 - \frac{2}{9}\delta_1 - \frac{2}{9}\delta_2 - \frac{4}{9}\epsilon_2 - \frac{1}{3}A + \frac{4}{9}A\delta_1 + \frac{5}{9}A\delta_2 \\ \dots \dots \dots \end{aligned}$$
(5)

Stability of triangular points :-**(B)** Differentiating (4), we have

$$\frac{\partial r_1}{\partial x} = \frac{x + \mu}{r_1}; \qquad \qquad \frac{\partial r_1}{\partial y} = \frac{y}{r_1}$$
$$\frac{\partial r_2}{\partial x} = \frac{x - 1 + \mu}{r_2}; \qquad \qquad \frac{\partial r_2}{\partial y} = \frac{y}{r_2}$$

Now differentiating (2) partially with respect to x, we get

### Volume 3 Issue 10, October 2014 www.ijsr.net Licensed Under Creative Commons Attribution CC BY

1139

$$= \beta - \frac{1}{1 + \frac{3A}{2}} \left\{ \frac{(1 - \mu)q_1}{r_1^3} + \frac{\mu q_2}{r_2^3} + \frac{3(1 - \mu)q_1A}{2r_1^5} \right\} + \frac{1}{1 + \frac{3A}{2}} \left\{ \frac{3(1 - \mu)q_1}{r_1^5} + \frac{3\mu q_2}{r_2^5} + \frac{15(1 - \mu)q_1A}{2r_1^7} \right\} y^2 \qquad \dots \dots \dots (9)$$

Substituting these values in (9), (10) and (11), we get  $\Omega_{xx}^{0} = \frac{3}{4} - \frac{1}{2}\delta_{1} + \delta_{2} + \frac{5}{4}\epsilon_{2} + \frac{9}{4}A + \mu\left(\frac{3}{2}\delta_{1} - \frac{3}{2}\delta_{2} - 3A\right)$   $\Omega_{yy}^{0} = \frac{9}{4} + \frac{1}{2}\delta_{1} - \delta_{2} + \frac{7}{4}\epsilon_{2} + \frac{3}{4}A - \mu\left(\frac{3}{2}\delta_{1} - \frac{3}{2}\delta_{2}\right)$   $\Omega_{xy}^{0} = \frac{3\sqrt{3}}{4} - \frac{1}{2\sqrt{3}}\delta_{1} + \frac{1}{\sqrt{3}}\delta_{2} + \frac{11}{4\sqrt{3}}\epsilon_{2} + \frac{5\sqrt{3}}{4}A - \mu\left(\frac{3\sqrt{3}}{2} + \frac{1}{2\sqrt{3}}\delta_{1} + \frac{1}{2\sqrt{3}}\delta_{2} + \frac{11}{2\sqrt{3}}\epsilon_{2} + \frac{11}{2\sqrt{3}}\epsilon_{2} + \frac{11}{2\sqrt{3}}\epsilon_{2} + \frac{11}{2\sqrt{3}}\epsilon_{3} + \frac{1}{2\sqrt{3}}\epsilon_{3} + \frac{1}{2\sqrt{3}}\epsilon_{$ 

Using these values, we obtain

 $\cap$ 

$$\Omega_{xx}^{0} + \Omega_{yy}^{0} - 4\alpha^{2} = -1 - 8\epsilon_{1} + 3\epsilon_{2} + 3\mu A$$
  
and  $\Omega_{xx}^{0} \cdot \Omega_{yy}^{0} - (\Omega_{yy}^{0})^{2}$   
 $= \frac{27}{4}\mu(1-\mu)\left(1 + \frac{2}{9}\delta_{1} + \frac{2}{9}\delta_{2} + \frac{4}{3}A + \frac{22}{9}\epsilon_{2}\right)$ 

The characteristic equation for the triangular equilibrium points, therefore, takes the form

 $\lambda^{4} + (1 + 8\epsilon_{1} - 3\epsilon_{2} - 3\mu A)\lambda^{2} + \frac{27}{4}\mu(1 - \mu)\left(1 + \frac{2}{9}\delta_{1} + \frac{2}{9}\delta_{2} + \frac{4}{3}A + \frac{22}{9}\epsilon_{2}\right) = 0 \qquad \dots \dots (10)$ 

This is a quadratic equation in  $\lambda^2$ . The discriminant D of the quadratic equation (10) is given by D = b<sup>2</sup> - 4ac

where 
$$b = 1 + 8\epsilon_1 - 3\epsilon_2 - 3\mu A$$
  
 $a = 1$   
 $c = \frac{27}{4}\mu(1-\mu)\left(1 + \frac{2}{9}\delta_1 + \frac{2}{9}\delta_2 + \frac{4}{3}A + \frac{22}{9}\epsilon_2\right)$ 

In order that  $\lambda$  should be pure imaginary in conjugate pair,  $\lambda^2$  must be negative quantity.

Here,  $\lambda^2 = \frac{-b \pm \sqrt{D}}{2a} = -\frac{b}{2} \pm \frac{1}{2}\sqrt{D}$ =  $-\frac{1}{2}(1 + 8\epsilon_1 - 3\epsilon_2 - 3\mu A) \pm \frac{1}{2}\sqrt{D}$ The roots are pure imaginary if  $D \ge 0$ . This inequality may be written as  $1 + 16\epsilon_1 - 6\epsilon_2 - 6\mu A$ 

$$-27\mu(1-\mu)\left(1+\frac{2}{9}\delta_{1}+\frac{2}{9}\delta_{2}+\frac{4}{3}A+\frac{22}{9}\epsilon_{2}\right)-\epsilon \ge 0 \qquad \dots (11)$$

where  $\epsilon$  is a positive quantity whose limit is zero. The equation (11) may be written as

$$\mu^{2} - \left(1 + \frac{2}{9}A\right)\mu + \frac{1}{27}\left(1 - \frac{2}{9}\delta_{1} - \frac{2}{9}\delta_{2} - \frac{4}{3}A + 16\epsilon_{1} - \frac{76}{9}\epsilon_{2} - \epsilon\right)$$
  
= 0

Since  $\mu$  represents the mass which is less than one half, the negative sign must be taken. At the limit  $\epsilon = 0$   $\mu$ 

$$=\frac{1+\frac{2}{9}A-\sqrt{1+\frac{4}{9}A-\frac{4}{27}\left(1-\frac{2}{9}\delta_{1}-\frac{2}{9}\delta_{2}-\frac{4}{3}A+16\epsilon_{1}-\frac{76}{9}\epsilon_{2}\right)}}{2}$$
$$=\frac{1}{2}\left\{1+\frac{2}{9}A-\sqrt{\frac{23}{27}\left(1+\frac{4}{207}\delta_{1}+\frac{4}{207}\delta_{2}+\frac{26}{69}A-\frac{32}{23}\epsilon_{1}+\frac{152}{207}\epsilon_{2}\right)}\right\}$$

The critical value of 
$$\mu$$
 is easily obtained as  

$$\mu_c = 0.1371742 - 0.0070111\delta_1 - 0.0070111\delta_2 - 0.02560585A - 0.500480105\varepsilon_1 - 0.2664228\varepsilon_2$$

Thus, when  $0 \le \mu < \mu_c$ ,  $D \ge 0$ , the values of  $\lambda^2$  given by the characteristic equation are negative and all the four roots of the characteristic equation are purely imaginary. Thus the triangular equilibrium points are stable. Moreover the range of stability of triangular equilibrium points decreases on account of oblateness and photo gravitational effect of the primaries. When oblateness and radiating effects are ignored, the critical value reduces to  $\mu_c = 0.0385209.....$ 

#### References

- Bhatnagar, K. B; Chawla, J. M., "The effect of oblateness of the bigger primary on collinear libration points in the restricted problem of three bodies", Celestial Mechanics, 16, pp. 129 – 136, 1977.
- [2] Bhatnagar, K. B.; Hallan, P. P, "Effect of Perturbation in coriolis and centrifugal forces on the stability of libration points in the restricted problem", Celestial Mechanics, 18, pp. 105, 1978.
- [3] Chaudhary, P. K.; Manaziruddin and Singh, R. B., "First order normalization in the restricted problem with one radiative and the other oblate primary", Proc. Math. Soc., B.H.U., 3, pp. 57-65, 1987.
- [4] Duboshin, G.N., "Libration points in general restricted three body problem", Astronomical journal, Vol. 73, No.2, pp. 104-109, 1970.
- [5] Haque, M.; Chaudhary, P.K. and Manaziruddin, "Non-Linear stability of the triangular points of equilibrium in the restricted three body problem when the bigger primary is an oblate spheroid", Proc. Nat. Acad. Sci. India 59 (A) – III, pp. 457 – 464, 1989.
- [6] Haque, M. N., "Effect of Perturbations on the Stability of Equilibrium Points in the Photogravitational Restricted Problem of Three Bodies", Ph.D. Thesis Submitted to the Bihar University, Oct., 1992.
- [7] Ishwar, S. K. "Effect of Perturbation on the Stability of Libration Points in the Generalised Photogravitational Restricted Three Body Problem" Ph. D. Thesis submitted to B. R. Ambedkar Bihar University, Muzaffarpur, Feb., 2004.
- [8] Danby, J. M., "Fundamental of Celestial Mechanics", Macmillan Co., New York, 1962.

## Volume 3 Issue 10, October 2014

Paper ID: OCT14415

www.ijsr.net Licensed Under Creative Commons Attribution CC BY