

Study of the Trends in Sunshine Duration Hours in Nigeria During 1961 – 2012 using the Linear Regression Model

S.O. Amadi¹, S.O. Udo², D.B. Pekene³, T.O. Chigbu⁴

¹Department of Physics, Geology and Geophysics, Federal University Ndufu-Alike Ikwo, Ebonyi State.

² Department of Physics, University of Calabar, Calabar, Cross River State

³ Department of Physics, Cross River University of Technology, Calabar, Cross River State.

⁴ Department of Physics and Electronics, Abia State Polytechnic, Aba, Abia State.

Abstract: *The trends in sunshine duration hours in Nigeria during 1961 – 2012 are presented in the context of renewable energy. The study is based on measurement and observation data from 20 meteorological stations across different climatic belts in Nigeria. The basic data which comprises monthly mean daily sunshine duration hours were obtained from the Nigeria Meteorological Agency, Oshodi, Lagos, Nigeria. The trend analysis was based on the linear regression model using the least squares method. It was found that 7 stations indicate downward trends with only 4 stations indicating significant downward trends at the 1% and 5% levels. 13 stations show upward trends with 7 stations indicating significant upward trends at 1% and 5% levels. The results indicate that solar energy resource is a strong candidate in Nigeria as alternative energy solution. The results equally appear to be hinting at the possibility of an impact of anthropogenic aerosols emissions on the dynamics of atmospheric circulations at synoptic scales.*

Keywords: Trends, sunshine duration, linear regression model, alternative energy, Nigeria.

1. Introduction

Within the last few decades, there have been detected considerable changes in the physical climate system at global and regional scales. The changes of major climate variables such as cloud cover, sunshine duration hours and the solar flux at the earth's surface have made it imperative, nay essential, to understand the potential of solar energy resource in different parts of Nigeria. This study examines the trends in sunshine duration hours in different regions of Nigeria using the least squares method of the linear regression model.

Some studies have examined the relationship between sunshine duration hours and climate change. Stanhill and Cohen (2005) observed upward and downward trends in the United States region during the 20th Century. Sanchez – Lorenzo *et al*, (2009) showed significant positive and negative trends at different time scales over the Iberian Peninsula during 1961 – 2004. In Turkey, Askoy (1999) found significant downward trends for annual sunshine duration hours at over 70% of the selected synoptic stations. Durlo (2006) found an insignificant increasing trend of effective sunshine duration in the BeskidSadecki Mountains in Poland during 1971 – 2005 based on measurement and observation data from six meteorological stations situated in the area. Sanchez – Lorenzo *et al*, (2008) found a direct correlation between the dimming and brightening phenomena of the sun with a negative trend for sunshine duration for Western Europe during 1938 – 2004. In Iran, Rahimzadehet *al*, (2014) found positive trends in seasonal and annual variations of sunshine hours for the vast majority of the stations during the period 1981 – 2007. Studies in China have indicated that the annual number of sunshine

hours has decreased during 1961 – 1998 and 1965 – 1999 (Chen *et al*, 2006; Kaiser and Qian, 2002; Yang *et al*, 2009).

Quite a number of studies related to sunshine duration in Nigeria have been done. Ewona and Udo (2011a) reported a positive correlation in sunshine duration between 1989 and 1996 in Niger Delta region of Nigeria. Ewona and Udo (2011b) reported mean daily sunshine hours of 4.69 hours per day for Calabar between 1985 and 2003. Yakubu&Medugu (2012) studied the relationship between the global solar radiation and sunshine duration hours in Abuja, Nigeria during 1991 – 2005 for the purpose of predicting global solar radiation. They observed a good agreement between the measured and predicted global solar radiation. Abdusalamet *al*, (2012) assessed the solar radiation patterns for sustainable implementation of solar home systems in Nigeria and concluded that solar energy resource is a favourable alternative by reason of availability of sunshine in the country. Ogolo (2014) investigated the trends of some common and related atmospheric variables including sunshine hours in the context of climate change for the period 1975 – 2006 using 13 meteorological stations. He observed that 98.8% of the stations indicated upward trend in sunshine hours of which 16% were significant at the 1% level. Ewonaet *al*, (2014) assessed decadal variation of sunshine duration in 23 locations in Nigeria from 1978 – 2007. They observed that sunshine duration displayed positive trends and showed marked latitudinal dependence. By virtue of the geographical location of Nigeria (tropical location), the long-term mean daily sunshine duration is supposedly high. Consequently, a rather high potential for solar energy production exists. Apart from the obvious role played by the direct solar radiation in plants and human lives, there are possibilities for its utilization for energy

production, as a clean alternative to the exploitation fossil fuel resources. Inferences drawn from climate change impacts on sunshine duration hours have provoked this study to examine the trends in sunshine duration hours as an excellent proxy measure of global solar radiation on interannual and decadal scales (e.g. Stanhill, 2003; Stanhill and Cohen, 2008; Liang and Xia, 2005). Sanchez – Lorenzo *et al* (2009) defined sunshine duration as the amount of time, usually expressed in hours, that direct solar radiation exceeds a certain threshold (usually taken at 120 Wm^{-2}). Rahimzadeh *et al*, (2014) reported that the desired minimum amount of sunshine needed to generate effective energy from solar panels is roughly 4 – 6 hours per day. It is important to understand the potential of solar energy resource in different parts of Nigeria due to prognosticated changes of major climate variable such as sunshine duration, atmospheric transmissivity and cloud-cover.

The aim of this study is to investigate the trends in the effective sunshine duration in Nigeria on the basis of the long-term measurements and observations conducted at 20 meteorological stations in Nigeria during the period 1961 – 2012.

2. Study Area and Data

Nigeria sits between longitude 2°E and 15°E and between latitude 4°N and 14°N . The monthly mean daily sunshine duration data in hours from 20 synoptic stations across Nigeria were obtained from the Nigerian Meteorological Agency, Oshodi, Lagos for the period 1961 – 2012. Fig 1 shows the map of Nigeria indicating the stations used for the study.



Figure 1: Map of Nigeria, showing the areas covered in the study.

3. Methodology

The trends, representing a general direction of changes in sunshine duration during the period 1961 – 2012 were determined. The monthly mean daily sunshine hours were used. The trend testing was based on the linear regression model, using the method of least squares. The estimation of the quality of the model fitting was carried out on the basis of the coefficient of determination and residual standard deviation.

3.1 Co-Efficient of Determination R^2

$$R^2 = \frac{\sum_{i=1}^n (\hat{y}_i - \bar{y})^2}{\sum_{i=1}^n (y_i - \bar{y})^2} \quad \text{.....(1)}$$

where R^2 = coefficient of determination; n = sample size; y_i = value of next observation i of y variable; \hat{y} = value of

regression function for x_i ; \bar{y} is the arithmetic mean of y variable.

3.2 Residual Standard Deviation

$$Se = \sqrt{\frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{y}_i)^2} \quad \text{.....(2)}$$

where Se is residual standard deviation.

To estimate the significance of regression coefficients, the values of standard errors of parameter estimators were used. The linear regression model is represented thus:

$$\bar{y} = \beta_1 \bar{x} + \beta_0 \quad \text{.....(3)}$$

The null hypothesis is that the slope coefficient, $\beta_1 = 0$. The t – statistic on β_1 is tested to determine if it is significantly different from zero. If β_1 is significantly non zero, the null

hypothesis is rejected and it can be concluded that there is a linear trend in y over time with the rate $= \beta_1$. Missing values are allowed. Values of β_1 (the slope coefficient) and β_0 (the intercept), and their errors were determined on the basis of the following formulae:

$$\beta_1 = \frac{\sum_{i=1}^n (x_i - \bar{x}) \cdot (y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \dots\dots\dots(4)$$

$$\beta_0 = \bar{y} - \beta_1 \cdot \bar{x} \dots\dots\dots(5)$$

where x is the independent variable (time), and y is the dependent variable (Sunshine duration).

$$D(\beta_1) = \frac{Se}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2}} \dots\dots\dots(6)$$

$$D(\beta_0) = Se \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2}} \dots\dots\dots(7)$$

where $D(\beta_1)$ and $D(\beta_0)$ are standard errors of the model estimators.

The significance of model parameters were tested on the basis of the t – statistic determined by the formulae:

$$t_{stat} = \frac{\beta_1 - 0}{D(\beta_1)} \dots\dots\dots(8)$$

$$t_{stat} = \frac{\beta_0 - 0}{D(\beta_0)} \dots\dots\dots(9)$$

The statistics has student's t – distribution. Testing of the significance of the model parameters was done on the basis of the following hypothesis:

Null hypothesis, $H_0 : \beta_1 = 0$ (lack of linear dependence):

Alternative hypothesis, $H_1 : \beta_1 \neq 0$ (Linear dependence exists).

The calculated values using eqns (8) and (9) were compared with the critical values determined from student's t – distribution at the assumed levels of confidence ($\alpha = 0.01$ and $\alpha = 0.05$). The null hypothesis was rejected if $t_{stat} > t_{\alpha}$ (ie, $p < \alpha$), thus accepting the alternative hypothesis, H_1 .

4. Results and Discussion

Table 1 is the result of linear trend estimation of the model estimators for sunshine duration hours in the stations studied. Also indicated in the table are the standard errors, the t – statistics and the p values of the model estimators.

Table 1: Results of linear trend estimation, standard errors of model estimators, t – statistics and the p -values.

	Parameter	Estimate	Std. Error	t	p -value	Slope Estimate	
						Hrs per yr	Hrs per decade
Yelwa	Slope	0.0020*	0.0010	2.0710	0.0390	0.024	0.24
	Intercept	6.0840	0.4840	12.5630	0.0000		
Sokoto	Slope	-0.0029**	0.0006	-4.9986	0.0000	-0.0348	-0.348
	Intercept	8.5090	0.2623	32.4385	0.0000		
Kaduna	Slope	-0.0007	0.0007	-0.9909	0.3226	-0.0084	-0.084
	Intercept	7.9503	0.3153	25.2135	0.0000		
Kano	Slope	0.0014*	0.0006	2.5265	0.0121	0.0168	0.168
	Intercept	7.1234	0.2531	28.1478	0.0000		
Bauchi	Slope	-0.0020**	0.0007	-2.6642	0.0082	-0.024	-0.24
	Intercept	7.3911	0.3381	21.8630	0.0000		
Maiduguri	Slope	-0.0004	0.0005	-0.7745	0.4393	-0.0048	-0.048
	Intercept	8.2004	0.2448	33.4952	0.0000		
Ilorin	Slope	0.0017**	0.0006	2.6208	0.0092	0.0204	0.204
	Intercept	5.3532	0.2911	18.3871	0.0000		
Yola	Slope	-0.0003	0.0006	-0.4241	0.6718	-0.0036	-0.036
	Intercept	7.4532	0.2678	27.8264	0.0000		
Ikeja	Slope	0.0024**	0.0007	3.6969	0.0003	0.0288	0.288
	Intercept	4.1784	0.2970	14.0672	0.0000		
Ibadan	Slope	0.0003	0.0007	0.4135	0.6796	0.0036	0.036
	Intercept	4.6746	0.3227	14.4879	0.0000		
Oshogbo	Slope	0.0003	0.0007	0.4436	0.6577	0.0036	0.036
	Intercept	5.1253	0.3180	16.1156	0.0000		
Benin	Slope	0.0011	0.0006	1.7829	0.0757	0.0132	0.132
	Intercept	4.3187	0.2762	15.6370	0.0000		
Warri	Slope	0.0040**	0.0007	6.0323	0.0000	0.048	0.48
	Intercept	2.6781	0.3007	8.9053	0.0000		
Lokoja	Slope	-0.0020**	0.0006	-3.2755	0.0012	-0.024	-0.24
	Intercept	6.7939	0.2817	24.1160	0.0000		
Port Harcourt	Slope	0.0025**	0.0007	3.5695	0.0004	0.03	0.3
	Intercept	3.1164	0.3131	9.9544	0.0000		
Owerri	Slope	0.0007	0.0006	1.1642	0.2453	0.0084	0.084
	Intercept	4.1542	0.2889	14.3791	0.0000		
Enugu	Slope	0.0004	0.0006	0.5910	0.5550	0.0048	0.048
	Intercept	5.2446	0.2758	19.0147	0.0000		
Calabar	Slope	0.0018**	0.0007	2.7044	0.0073	0.0216	0.216
	Intercept	2.8064	0.2999	9.3565	0.0000		

Makurdi	Slope	0.0001	0.0006	0.1517	0.8795	0.0012	0.012
	Intercept	6.0249	0.2802	21.5016	0.0000		
Ogoja	Slope	-0.0016*	0.0007	-2.2111	0.0278	-0.0192	-0.192
	Intercept	6.1740	0.3219	19.1811	0.0000		

** Trend is significant at the 1% level (2-tailed) * Trend is significant at the 5% level (2-tailed).

7 stations show downward trends with 4 stations having significant trends. Apart from Ogoja and Lokoja, all the stations that show downward trends fall within the Sudan and Sahelian Savanna regions of the country. 13 stations show upward trends with 7 stations showing significant positive trends.

Several explanations could account for the observed trends in sunshine duration in Nigeria. Clouds exert a dominant influence on the global energy balance and can be attributed to be the leading cause of the trend variations across Nigeria. Cloudiness can contribute to dimming, i.e. low-level clouds types linked to their high albedo, and also brightening, ie, high clouds types emit less radiation out to space than do low clouds, or the clear atmosphere (Mace *et al*, 2006). Changes in atmospheric transmissivity (aerosol optical thickness) due to changes in the concentrations and optical properties of aerosols can also cause dimming or brightening depending on the type of aerosol causing the local pollution.

Uncertainties of the indirect effects of aerosols on clouds and precipitation could also result to either induced changes in the cloud properties such as albedo and lifetime, or the modifications of precipitation forming processes. These aerosol indirect effects on clouds and precipitation are in the form of cloud-albedo effect, cloud lifetime effect, semi direct effect, glaciation effect and the thermodynamic effect. The cloud- albedo and cloud lifetime effects can be brought about by sulphate aerosols that have the tendency to produce small cloud droplets that reflect solar radiation more efficiently, decrease precipitation formation and prolong cloud lifetime. These two effects presumably contribute to the observed downward trends in sunshine duration in some of the locations. Soot aerosols (particulate black carbon) absorb solar radiation and re-emit it as thermal radiation which consequently heats the air – mass, causing evaporation of cloud droplets. This aerosol heating within cloud layers reduce cloud fractions, and cause change in cloud amounts (Ramanathan, *et al*, 2001). This semi-direct effect, as it is known, can contribute to the increasing trends in sunshine durations in locations with large fossil fuel burning and several other manufacturing concerns such as Port Harcourt, Ikeja and Warri etc. Large aerosols have the consequences of precipitation with the formation of fewer and larger droplets. These large aerosols are efficiently scavenged by precipitation which is their main atmospheric sink, resulting in atmospheric lifetime of few days or few weeks. All things being equal, this effect will consequently increase effective sunshine hours and can also account for the upward trends observed in some stations.

Given the observed trends at the stations in Nigeria over the 52 year period, and realising the high sunshine duration experienced across the regions, Nigeria has favourable conditions for solar energy resource. Descriptive statistical analysis (not shown) shows that the monthly mean daily sunshine hours for the 52 year period (ranging from 444 –

624 months) range from over 4 hours per day in the coastal locations to over 8 hours per day in the high latitude locations. The effective sunshine duration in Nigeria is characterised by high variation in space and time. Seasonal variation plots (not shown) indicate that maximum daily sunshine hours are observed from November to March in all the stations, exceeding 6.5hours in the southern low latitudes and having up to 9hours in the high latitude locations in the north. Minimum daily sunshine hours are observed from June to September across the stations recording about 6hours in the high latitude stations and 3 hours in the low latitude stations.

The results of this study are in agreement with earlier studies particularly Ogolo (2014), Ewona *et al*, (2014), Ewona and Udo (2011a), and Ewona and Udo (2011b).

5. Conclusion

The trends of sunshine duration hours in Nigeria for the period 1961 – 2012 have been investigated using the least square method of the linear regression model. The results indicate, overall regions, that the pattern of sunshine duration exhibited sinusoidal increases and decreases in seasonal variations. The trends indicate that the sign of the trends for the vast majority of the station is positive. Given the increasing trends observed at many stations and some decreasing trends that are not significant in some stations over the 52 years period, and realising the relatively high monthly mean daily sunshine duration hours, solar energy resource is a good candidate in Nigeria as alternative energy solution to offset carbon emissions. The effective sunshine duration across Nigeria is characterised by high spatial and temporal variations which could be linked to some factors such as the effects due to cloud cover, direct and indirect aerosol effects, latitudinal locations, and orographic conditions. Nevertheless, the results in some cities, intriguing as they appear, may be hinting at the possibility of an impact of anthropogenic aerosols emissions on the dynamics of the atmospheric circulation at synoptic scales, and needs to be explored further in subsequent times.

As a result of large geographical coverage of Nigeria, the 20 stations used in this study do not have the capacity to generate complete meteorological data for the Country even though they are representatives of different climatic belts and agro-ecological zones in Nigeria. Further research could improve on this by examining more stations to get finer grids especially in the north. Furthermore, the non-parametric Mann-Kendall's rank correlation test, which is more robust to missing data and outliers, could be used for analysis in further works in this regard to create room for comparison of the parametric and non-parametric tests.

References

- [1] Stanhill, G and Cohen, S. (2005). Solar radiation changes in the United States during the 20th Century: Evidence from sunshine duration measurements. *J.Clim.* 18(10), 1503-1512, doi:10.1175/jcli 3354.1.
- [2] Sanchez-Lorenzo, A., Calbo, J., Brunetti, M., and Deser, C. (2009). Dimming/brightening over the Iberian Peninsula: Trends in sunshine duration and cloud cover and their relations with atmospheric circulation. *J.Geophys.Res.* 114, D00D09, doi:10.1029/2008JD011394.
- [3] Askoy, B. (1999). Analysis of changes in sunshine duration data for Ankara, Turkey. *Theor. Appl. Climatol.*, 64, 229 – 237.
- [4] Durlo, G., (2006). Multi annual variation of the effective sunshine duration in the Beskid Sadecki Mountains. *Electronic Journal of Polish Agricultural Universities*, 9(4), 12 pages.
- [5] Sanchez- Lorenzo, A., Calbo, J., Martin-Vade, J. (2008). Spatial and temporal trends in sunshine duration over Western Europe (1938 - 2004). *J.Clim.* 21, 6089 – 6098, doi: 10.1175/2008jcli 2442.1.
- [6] Rahimzadeh, F., Pedram, M., and Kruk, M.C. (2014). An examination of trend in sunshine hours over Iran. *Meteorological Applications*, 21(2), 309-315 doi: 10.1002/met.1334.
- [7] Chen, R., Kang, E., Yang, J., Ji, X., and Zhan, Z (2006). Trends of the global solar radiation and duration of sunshine. *Q.J.R Meteorol. Soc.*, 80, 231 – 235.
- [8] Kaiser, D.P. and Qian, Y. (2002). Decreasing trends of sunshine duration over China for 1954 – 1998: Indications for increased haze pollution? *Geophys.Res. Lett.* 21, 2042 – 2046.
- [9] Yang, Y., Zhao, N., Hao, X. and Li, C. (2009). Decreasing trend of sunshine hours and related driving forces in North China. *Theor.Appl.Climatol.*, 97, 91 – 98.
- [10] Ewona, I.O. and Udo, S.O. (2011a). Changes in some meteorological parameters in the Nigeria Delta Region of Nigeria between 1989 – 1996. *Global Journal of Pure and Applied Sciences*, 17(1), 61 – 69.
- [11] Ewona, I.O. and Udo, S.O. (2011b). Climatic condition of Calabar as typified by some meteorological parameters. *Global Journal of Pure and Applied Sciences*, 17(1), 81 – 86.
- [12] Yakubu, D. and Medugu, D.W. (2012). Relationship between the global solar radiation and the sunshine duration in Abuja, Nigeria. *Ozean Journal of Applied Sciences*, 5(3), 221 – 228.
- [13] Abdusalam, D., Mbamali, I., Mamman, M. and Saleh, Y.M. (2012). An assessment of solar radiation patterns for sustainable implementation of solar home system in Nigeria. *American International Journal of Contemporary Research*, 2(6), 238 – 243.
- [14] Ogolo, E. O. (2014). Estimation of global solar radiation in Nigeria using a modified Angstrom model and trend analysis of the allied meteorological components. *Indian Journal of Radio and Space physics*, 43, 213 – 224.
- [15] Ewona, I.O., Udo, S.O., and Osang, J.E. (2014). An assessment of decadal variation of sunshine duration in Nigeria. *Journal of Current Research in Science*, 2(1), 59 – 62.
- [16] Stanhill, G. (2003). Through a glass brightly: some new light on the Campbell-Stokes sunshine recorder. *Weather*, 58, 3-11, doi: 10.1256/wea.278.01.
- [17] Stanhill, G and Cohen, S. (2008). Solar radiation changes in Japan during the 20th century: Evidence from sunshine duration measurements. *J. Meteorol. Soc.Jpn*, 86, 57 – 76, doi:10.2151/jmsj.86.57.
- [18] Liang, F. and Xia, X.A. (2005). Long-term trends in solar radiation and the associated climatic factors over China for 1961 – 2000. *Ann.Geophys.*, 23, 2425 – 2432.
- [19] Mace, G.G., Benson, S. and Kato, S. (2006). Cloud radiative forcing at the Atmospheric Radiation Measurement Program Climate Research Facility: Vertical redistribution of radiant energy by clouds. *J.Geophys.Res.*, 111, D11591, doi:10.1029/2005 JD005922.
- [20] Ramanathan, V., Crutzen, P.J., Kiehl, J.T., and Rosenfeld, D. (2001). Aerosols, climate and the hydrological cycle, *Science*, 294, 2119 – 2124, doi:10.1126/science. 1064034.