





Figure 1: Procedure of beta-glucan extraction from barley

## 2.4 Analysis on isolated $\beta$ -glucan pellets

### 2.4.1 Estimation of $\beta$ -glucan

The isolated pellets were estimated for  $\beta$ -glucan content by using megazyme assay kit, Ireland. This was carried out in order to identify the recovery percentage of  $\beta$ -glucan content from the barley flour by hot water extraction method.

### 2.4.2 Proximate composition

Crude fat and ash content were determined according to the approved methods of the American Association of Cereal Chemists Method 08-01 and method 30-10, respectively [9]. Nitrogen content was determined by the kjeldahl method [10] and converted to protein content by a factor of 6.25. Total, Soluble and Insoluble dietary fibre content was determined by following the method of IS 11062-1984.

## 2.5 Functional Property Analysis

### 2.5.1 Foaming property

The foaming capacity and foam stability (FS) were studied by following the method of Temelli [11].

### 2.5.2 Water binding capacity

The water binding capacity (WBC) was determined by the modified method described by Wong and Cheung [12].

### 2.5.3 Viscosity

Viscosity was measured using Brookfield viscometer.

## 2.6 Biological Properties (In-vitro)

### 2.6.1 Anti-oxidant activity

In the present part of the study, antioxidant activity of the isolated  $\beta$ -glucan pellets were identified by measuring Free radical scavenging activity and Metal chelating activity according to the method of Braca [13] and Benzie [14]. The

sample taken for each analysis is 5 $\mu$ l and the assay was performed in three triplicates.

### 2.6.2 Anti-bacterial activity

Well-diffusion assay was used to evaluate the anti-bacterial activity of isolated  $\beta$ -glucan pellets against *S.aureus* (gram positive) and *E.coli* (gram negative) bacteria on MHA medium. Sample of different volume (10 to 40 $\mu$ l) at various concentrations was added into the wells. The plates were incubated for 24 hours at 37°C and the zone of inhibition was measured using zone inhibitory scale. Kanamycin (30mg) acted as control. The assays were performed in triplicates.

## 3. Results and Discussion

### 3.1 Estimation of Beta-glucan

Beta glucan is a polysaccharide that are composed of glucose units linked together by beta glycosidic linkage having (1,3) and (1,4) bonds. The most documented nutritional benefit of beta-glucan is flattening postprandial glucose, insulin and cholesterol rise. FDA has acknowledged and recommended 3gm of beta glucan per day for achieving the health benefits. As beta glucan has shown to have positive influence on human health, in the present study it was estimated in both selected oats and barley.

Table 1: Estimation of beta-glucan in Oats and Barley

Samples	Beta-glucan (%) (w/w)
<i>Oats</i>	
I	2.91
II	2.80
III	2.55
<i>Barley</i>	
Hulled barley	4.83
Broken barley	3.96
Pearl barley	2.95

From the estimated results, it was found barley had the maximum content of beta glucan compared to oats. This was proved by many scientists that of the common cereals like wheat, rye oats and barley, the largest amounts of  $\beta$ -glucan are found in barley (3-11%) and in oats (3-7%) [15]. Differences in beta glucan content among the barley were also identified. This may be due to the sample selected although barley is generally considered as the higher beta-glucan level material [16].

### 3.2 Extraction of Beta-glucan

The general extraction procedure of beta glucan was discussed in the methodology. The aim of this part of study is to produce beta-glucan with maximum yield and recovery. In-order to achieve this best and cheaper method was carried out and the yielded pellets were analyzed for the chemical and biological parameters to determine its efficacy.

### 3.3 Analysis on isolated $\beta$ -glucan pellets

#### 3.3.1 Estimation of $\beta$ -glucan

**Table 2:** Yield and percentage recovery of beta-glucan from barley

Weight of flour (gm)	Weight of isolated Pellets (gm)	Percentage of Extract (%)	
100	5.4	5.4	
Percentage of Recovery of beta-glucan content (%)			
Beta-glucan (%) (w/w)	Name of the Sample	Beta-glucan content (%)	Percentage of Recovery (%)
	Flour	4.83	83.8
	Extract	4.05	

From the extraction, 5.4 gm of pellets was yielded from 100gm of whole grain flour which represent 5.4% of extract. As the yielded pellets did not represent the whole quality of beta-glucan isolated the percentage of recovery was identified and it was found to be of 83.8%. This maximum recovery was achieved in the present study due to the best extraction method – hot water extraction. Study conducted by Asif Ahmad et al on different extraction procedures showed hot water extraction process had significant yield and recovery.

Previous studies by various authors [17] reported 57.8% - 88.4% by using different temperature and pH in the extraction methods. Lower efficiency by hot water extraction method was observed by Symnons and Brennan[18] that might be due to the beta-glucan cleavage by  $\beta$ -glucanase enzyme whereas in current study hot water method of extraction showed maximum recovery (83.8%). This may be due to the preliminary treatment involved in the extraction procedure like refluxing with 80% ethanol inactivates the  $\beta$ -glucanase enzyme which results in higher recovery and another reason might be due to maximum starch gelatinisation and protein solubilisation. Moreover indigenous enzyme system was inactivated.

#### 3.3.2 Proximate composition

The chemical composition of the isolated pellets is present in the below table

**Table 3:** Chemical composition of beta-glucan pellet

Chemical components (%)	Beta-glucan Pellet
Protein	7.02
Fat	0.79
Ash	1.20
Total dietary fiber	93.81
Soluble dietary fiber	87.52
Insoluble dietary fiber	6.01

From the obtained result it was found that protein content of 7.02% in the pellets. The protein content of isolated gum may affect the various functional characteristics of the product in which they are added as a functional additive. With respect to fat content, 0.79% was obtained, this appreciable removal of fatty portion is essential, since the oxidative degradation of fat is important during processing and storage. This may produce bitter taste in the foods in which beta-glucan is added as a functional ingredient, thus the overall quality of the product can be deteriorated [19].

As concerned to ash (minerals) content of beta-glucan pellets, 1.20% was found. Detailed analysis of beta-glucan gum by Ahmad et al. showed that an appreciable amount of phosphorus (3010mg/kg) and potassium (3318mg/kg) is present in the isolated samples.

Regarding dietary fiber, major compositions of the beta-glucan consisted of soluble dietary fiber (87.52%) which is the sole component in the reduction of sugars and are responsible for reduced glycaemic index [20], [21] with relatively low amount of insoluble fiber (6.01%).

#### 3.4 Functional Property Analysis

The functional properties are the intrinsic physico-chemical characteristics which affect the behavior of a food ingredient in food systems during processing, manufacturing, storage and preparation. Such functional properties are important in determining the quality of the final product [22].

**Table 4:** Functional analysis of beta-glucan pellet

Functional parameters	Beta-glucan pellet
Water-binding capacity (g/g)	3.62
Foaming capacity (%)	170
Foam stability (%)	63.10
Viscosity (cP)	50.19

The water binding capacity measures the amount water retaining capacity of the isolated pellets when it is subjected to centrifugation. This hydration property is found to be of 3.62g/g in the present study, which is higher than the water binding capacity of some dietary fibers such as wheat bran 2.6g/g, corn bran 2.5g/g and soybean bran 2.4g/g [23]. The extract with appreciable binding capacity obtained in our study could be utilized as a functional ingredient in the preparation of various food products like jam, jellies, cheese etc to overcome syneresis problem.

Foaming capacity and stability are the two important factors when the beta-glucan is used as a functional ingredient. The above table reveals 170% of foam capacity and 63.10% of foam stability which represents high foaming capacity and stability that shows more desirable characteristics in the preparation of cakes and batter. Regarding viscosity, 50.19 cP was determined at 20°C. Different extraction procedures of varying pH and temperature may result low or high viscosity this may be due to the sensitivity of (1,3) b-D bond. Similarly the high viscosity of the beta glucan content depends on the soluble fibre than the insoluble fiber which tends to lower the viscosity.

#### 3.5 Biological Properties (In-vitro)

Biological property describes the beneficial or adverse effects of a compound on living matter. Among the various properties of chemical compounds, biological property plays a crucial role since it suggests uses of compounds in the medical applications.

##### 3.5.1 Anti-oxidant activity

Antioxidant compounds play an important role as a health protecting factor. Scientific evidence suggests that antioxidants reduce the risk of chronic diseases including

cancer and heart disease by its ability to trap free radicals in the biological system [24].

**Table 5:** Antioxidant activity of beta-glucan pellet

Method	Percent of Activity
DPPH	40.07
Metal chelating	39.22

The reactive oxygen species removal was studied in-vitro by applying free radical scavenging activity of Beta Glucan. From the result it was found that the activity on the inhibition or removal of free radical by Beta Glucan, thus exhibiting a potential scavenging role by DPPH scavenging activity of 40.07% and Metal chelating activity of 39.22%. Overall the removal of free radical by scavenging activity and by metal binding ability of Beta-glucan makes it a super molecule-bioactive substance. The present result coincides with the study conducted by Kyoko kofuji et al [25], in which beta-glucan derived from barley extracted by various acidic to alkaline conditions proved to have high scavenging activity than various polymers that are used as food additives.

### 3.5.2 Anti-bacterial activity

**Table 6:** Antibacterial activity of beta-glucan pellet

Inhibition zone Measurement (mm)						
Bacteria	Strain - NCIM	Control - Kanamycin	Volume ( $\mu$ l)			
			10	20	30	40
			Concentration ( $\mu$ g)			
			400	800	1200	1600
S.aureus	2079	16	4	12	19	23
E.coli	2065	21	0	6	13	19

From the results, it was found that beta-glucan extract showed inhibition at the minimum concentration of 30 $\mu$ l (1.2mg concentration) and at 40 $\mu$ l (1.6mg concentration) it demonstrated a significant inhibiting property towards gram positive bacteria (S.aureus) than the control. Whereas there was no significant inhibition towards gram negative bacteria (E.coli), minimum inhibition zone was observed at the 30 $\mu$ l of beta glucan extract having 1.2mg concentration. The antibacterial activity of cereal beta glucan was also proved by yun et al [26].

## 4. Conclusion

It was evident from the study that the  $\beta$ -glucan extracted by hot water extraction method has both scientific and commercial value and it is a cheap and promising additive, having a great potential with respect to its properties. Hence beta-glucan from barley could find the ability to act as a functional and nutraceutical ingredient in food products.

## 5. Recommendations for Future Work

The use of barley beta-glucan as a functional food ingredient definitely warrants future research. This study has generated a number of areas for further research. Particularly, beta-glucan interactions with other food components are necessary that might be able to increase its use in the development of food and other new novel products.

## References

- [1] Newman, R. K., Newman, C. W. and Graham, H. 1989. The hypocholesterolemic function of barley  $\beta$ -glucan. *Cereal Foods World* 36: 800 – 805.
- [2] McIntosh, G. H., Whyte, J., McArthur, R. and Nestel, P.J. 1991. Barley and wheat foods: Influence on plasma cholesterol concentrations in hypercholesterolemic men. *Am. J. Clin. Nutr.* 53: 1205-1209.
- [3] Kloprenstein, C.F. and Hosney, R.C. 1987. Cholesterol lowering effect of beta-glucan-enriched bread. *Nutr. Rep. Int.* 36:1091.
- [4] McIntosh, G.H., Jorgensen, L. and Royle, P. 1993. The potential of an insoluble dietary-fiber rich source from barley to protect from DMH-induced intestinal tumors in rats. *Nutrition and Cancer – An International Journal* 19:213-221.
- [5] Jenkins, D. J., Wolever, T.M., Nineham, R., Taylor, R., Metz, G.L., Bacon, S. and Hockaday, T.D. 1978. Guar crispbread in the diabetic diet. *British Medical Journal* 2:1744-1746.
- [6] Bhatti, R. S. 1993. Extraction and enrichment of (1,3) (1,4)  $\beta$ -glucan from barley and oat brans. *Cereal Chemistry* 70: 73-77.
- [7] McCleary, B. V. and Codd, R. 1991. Measurement of (1,3) (1,4)- $\beta$ -glucan in barley and oats: a streamlined enzymic procedure. *Journal of the Science of Food and Agriculture* 55: 302 – 312.
- [8] Asif Ahmad., Faqir Muhammad Anjum., TahirZahoor., Haq Nawaz. and Ahmad Din; 2009. Physicochemical and functional properties of barley  $\beta$ -glucan affected by different extraction procedures. *International Journal of Food Science and Technology*.44:181 – 187.
- [9] AACC., 2003. Approved Methods of American Association of Cereal Chemists. St. Paul: The American Association of Cereal Chemists, Inc.
- [10] AACC. 1990. Approved Methods of American Association of Cereal Chemists. St Paul: The American Association of Cereal Chemists, Inc.
- [11] Temelli, F. 1997. Extraction and functional properties of barley  $\beta$ -glucan as affected by temperature and pH. *Journal of Food science* 69: 1192 – 1201.
- [12] Wong, K. H. and Cheung, P. C. K. 2005. Dietary fiber from mushrooms sclerotia: preparation and physicochemical and functional properties. *Journal of Agriculture and Food Chemistry* 53: 9395 – 9400.
- [13] Braca A, Sortino C, Politi M et al. Anti-oxidant activity of flavonoids from Licanialicaniaeflora. *Journal of Ethnopharmacol* 79: 379-381, 2002.
- [14] Benzie, I. F. F. and Strain, J. J. 1999. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods in Enzymology* 299: 15–27.
- [15] Wood, P. J. 2007. Cereal  $\beta$ -glucan in diet and health. *Journal of Cereal Science* 46: 230 – 238.
- [16] Poppitt, S.D. 2007. Soluble oat and barley  $\beta$ -glucan enriched products: Can we predict cholesterol-lowering effects? *Br. J. Nutr* 97:1049-1050.
- [17] Brukus, Z. and Temelli, F. 1998. Effect of extraction conditions on yield, composition and viscosity stability of barley  $\beta$ -glucan gum. *Cereal Chemistry* 75: 805-809.

- [18] Symons, L. J. and Brennan, C. S. 2004. The effect of barley  $\beta$ -glucan fiber fractions on starch gelatinization and pasting characteristics. *Journal of Food Science* 69: 257-261.
- [19] Molteberg, E. L., Gjermund, V., Astrid, N and Wenche, F. 1995. Effect of storage and heat processing on the content and composition of free fatty acids in oats. *Cereal chemistry* 72: 88-93.
- [20] Cavallero, A., Empilli, S., Brighenti, F. and Stanca, A. M. 2002. High (1,3) (1,4)  $\beta$ -glucanbarle fractions in bread making and their effects on human glycemic response. *Journal of Cereal Science* 36: 59-66.
- [21] Symons, L. J. and Brennan, C. S. 2004. The influence of (1,3) (1,4)  $\beta$ -glucan rich fraction from barley on the physicochemical properties and in-vitro reducing sugar release of white wheat breads. *Journal of Food Science* 69: 463-467.
- [22] Lakshmi Menon., SwarnaliDuttaMajumdar. AndUsha Ravi. 2014. Mango kernel flour as a potential ingredient in the development of composite flour bread. *Indian Journal of Natural Products and Resources* 5(1): 75-82.
- [23] Dreher, M. L. 1987. *Handbook of Dietary Fiber: An Applied Approach*. New York: Marcel Dekker.
- [24] Balalyeswarya, M., Manigundan, K., Usha, J. and Shanmugapriya, A. 2014. Antioxidant and Anti-proliferative activity of methanolic leaf extract of *Eupatorium glandulosum*. L. *Internation journal of Advanced Research* 2(7): 717-723.
- [25] Kyoko kofuji., Ayumi Aoki., KazufumiTsubaki., Masanori Konishi., Takashi Isobe and Yoshifumi Murata. 2012. Antioxidant Activity of  $\beta$ -glucan. *ISRN Pharm*: 125864.
- [26] Yun, C. H., Estrada, A., Van Kessel, A., Byung-Chul P.I. and Laarveld, B. 2003. Beta-glucan, extracted from oat, enhances disease resistance against bacterial and parasitic infection. *FEMS Immun. Med. Microbiol.* 35: 67-75.

### Author Profile

**Kayal vizhi V.**, Ph. D, Research scholar at Bharathidasan Government College for women, has been awarded Gold Medal in PG level. Presented several papers and won two best awards on National and International conference.

**Dr. Josephine Nirmala Many**, Associated Professor, BGCW (Autonomous), Department of Home Science has about two decade of teaching experience and guided 35 PG and 10 Research Projects.