

HPLC, Antioxidant and Antibacterial Activities of Methanolic Extract of *Berberis aristata* Stem

Ovas Dar¹, Reena Lawrence², Sikander Dar³

^{1,3}Research scholar, Chemistry Department,
Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P, India

²Assistant Professor, Chemistry Department, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, U.P, India

Abstract: The components in methanolic extract of *Berberis aristata* stem were identified by HPLC. The three components which were identified are Berberine (95.52%), Palmitine (2.64%) and Jatrorrhizine (1.83%). The antioxidant activity of the methanolic extract were determined by DPPH (1, 1-Diphenyl-2-picryl hydrazyl) assay and Nitric oxide scavenging method. The methanolic extract was able to reduce stable purple colored DPPH reaching 50% of reduction with IC₅₀ values as – IC₅₀ for methanolic extract=0.90 mg/ml, IC₅₀ for Gallic acid = 0.61 mg/ml IC₅₀ for BHT = 0.41 mg/ml. The methanolic extract reduced the concentration of DPPH free radical with an efficacy near to that standard Gallic acid. In case of Nitric oxide scavenging assay, IC₅₀ values have been found to be 0.80 mg/ml, 0.65 mg/ml and 0.60mg/ml for the methanolic extract, gallic acid and BHT respectively. The antibacterial activity of methanolic extract was evaluated on bacteria strains like *Escherichia coli*, *Staphylococcus aureus* and *Bacillus subtilis* by using agar well diffusion method. The results were compared with ampicillin, a commercial antibacterial agent. Methanolic extract showed maximum (19mm) zone of inhibition against *Bacillus subtilis*. The MIC values of methanolic extract for different pathogenic bacteria ranged from 37 to 111µg/ml.

Keywords: Methanolic extract, HPLC, IC₅₀, Zone of inhibition, MIC

1. Introduction

Berberis aristata commonly known as “Daru haldhi and chitra” is spinous shrub native to northern Himalayan region. The plant is widely distributed from Himalayas to Srilanka, Bhutan, and hilly areas of Nepal in Himalaya region (1). *Berberis aristata* is used in ayurvedic medicines from very long time. The plant is used traditionally in inflammation, wound healing, skin disease, menorrhagia, diarrhoea, jaundice and diabetes (2).

Berberis aristata contains protoberberine and bis isoquinoline type of alkaloid. Root of plant *B. aristata* contains alkaloids which are berbamine, berberine, oxycanthine, epiberberine, palmatine, dehydrocaroline, jatrorrhizine and columbamine (3), karachine (4), dihydrokarachine, alkaloids, pakistanine, 1-Omethylpakistanine, pseudopalmatine chloride and pseudoberberine chloride were also isolated from *Berberis aristata* (5). *Berberis aristata* roots have been used in treatment of jaundice in Ayurveda. Hypoglycemic effect of *Berberis aristata* root was evaluated. Dried and powdered root extracted with water and methanol was administrated to normal and alloxan induced diabetic albino rabbit. The results show that *B. aristata* roots contain potent and orally effective antidiabetic component which either triggers the formation of insulin or shows insulin like effect (6). Ethanolic root extract of *B. aristata* shows antifungal activity was found against *Candida* species and *Aspergillus* species (7). Thus, the aim of this study is to investigate the chemical composition, antioxidant and antibacterial activities of methanolic extract of *Berberis aristata* stem.

2. Material and Methods

2.1 Sample Preparation

The dried stem of *Berberis aristata* was available in pieces of variable length and thickness. The dried stem was pulverized using sterile laboratory mortar and pestle to obtain the powdered form. These were stored in airtight glass containers protected from sunlight until required for analysis.

2.2 Preparation of Extract

The dried and powdered plant material was extracted with methanol using soxhlet extractor for 5 h at a temperature not exceeding the boiling point of the Solvent (8). The extract was filtered and then concentrated to dryness.

2.3 HPLC

HPLC fingerprinting and content determination methods were applied to evaluate methanol extract of *Berberis aristata*. The sample was analyzed in model Shimadzu CLASS –VP V6.14 SPI, AYOTO Japan/LC/20 ATPC. Two pumps were used STP-20 and SPD. Detector used was AVP-UV. Reverse phase HPLC column (C-18) was used with length 260X4.6mm, thickness being 5µ. Column Temperature was 36°C. Injected volume was 10 µl. Mobile phase was methanol-water (water containing 4% acetic acid). The ratio of methanol: water being 66:34 v/v. Flow rate 1ml/min. Two detectors were of 290 and 250nm.

2.4 Antioxidant activities assay

2.4.1 DPPH Free Radical Scavenging Assay

The DPPH radical scavenging activity assay elucidated by [9] was followed. The hydrogen atom or electron donating ability of the corresponding methanolic extract was

measured from the bleaching of the purple colored methanol solution of DPPH. This spectrophotometric assay uses the stable radical, 2, 2- diphenyl-1-picrylhydrazyl (DPPH), as a reagent. The working solution of the test extract was prepared in methanol. BHT and Gallic acid were used as standard in 1-100µg/ml solution. DPPH (0.002%) was prepared in methanol and 1ml of sample solution separately. The solution mixture was kept in dark for 30min and absorbance was measured at 517nm. DPPH solution (1ml) was used as blank. The absorbance was recorded and % inhibition was calculated using the formula given below.

$$\% \text{ scavenging activity} = \frac{A - B}{A} \times 100$$

Where A= absorbance of the control and

B= absorbance of the sample

IC₅₀= value, which represented, as the concentration of methanolic extract that caused 50% scavenging as calculated from the plot inhibition percentage against concentration.

2.4.2 Nitric Oxide scavenging method

The Nitric Oxide scavenging method elucidated by (10) was followed. Nitric oxide (NO) was generated from Sodium Nitroprusside (SNP) and was measured by Griess reagent. SNP in aqueous solution at physiological pH spontaneously generate NO, which interacts with oxygen to produce nitrite ions that can be estimated by the use of Griess Reagent. Scavenging of NO compete with oxygen leading to reduced production of NO. SNP (10mM) in phosphate buffer saline (PBS) was mixed with different concentration of extract (100-1000µg/ml) of the drug dissolved in methanol and incubated at 25°C for 180 min. The samples from the above were reacted with Griess reagent (1% sulphanilic acid, 0.1% naphthylamine and 3% phosphoric acid). The absorbance of the chromophores formed during the diazotization of nitrite with sulphanilic acid and subsequent coupling with naphthylamine was recorded at 546 nm and referred to the absorbance of Gallic acid, used as a positive control treated in the same way with Griess reagent.

2.4.3 Antibacterial activity

Antibiogram was done by agar well diffusion method (11, 12) using plant extracts and commonly used antibiotics. The test quantity of specific extracts were dissolved in dimethylsulphoxide (DMSO), depending upon the solubility of the extracts. The dissolution of the organic extracts was aided by 1% (v/v) DMSO, which did not affect the growth of microorganisms, in accordance with our control experiments. The surface of media was inoculated with bacteria from a broth culture. Wells or cups of 5 mm size were made with sterile borer into agar plates containing the bacterial inoculums. 20µl volume of the sample extracts of

concentration (2.0 mg/ ml) was poured into a well of inoculated plates. Ampicillin (10mg/ml) was used as a positive control which was introduced into the well instead of plant extract. After 18 h of incubation at a specific temperature the plates were examined and the diameters of the inhibition zones were measured to the nearest millimeter. Antibacterial activity was recorded if the radius of zone of inhibition was greater than 4 mm (13). The antibacterial activity results was considered as inactive if < 4.5 mm; 4.5-6 mm as partially active; while 6.5-9 mm as active and greater than 9mm as very active (14).

2.5 Minimum Inhibitory Concentration assay

Minimum Inhibitory Concentration of fruit peel extracts was also determined (15, 16). Dilution of the fruit peel extracts of Citrus Karna was prepared in sterile nutrient broth to achieve a decreasing concentration ranging from 1000µg/ml to 12µg/ml in sterile tubes labeled 1 to 5. Each dilution was seeded with 10µl of the standardized bacterial inoculums (108-109) CFU/ml. The inoculated culture tubes were incubated at 37°C for 24hrs. A set of tubes containing only seed broth (i.e. without plant extract) was kept as control. The lowest dilution of the plant extract that retained its inhibitory effect resulting in no growth (absence of turbidity) of a microorganism was recorded as the MIC value of the extract. After incubation, 10µl of content of each test tube was transferred with a loop on to nutrient agar media. Agar plates were incubated for 24 hours at 37°C. The lower concentration that did not permit any visible growth when compared with the control was considered as the MIC.

3. Results and Discussion

3.1 HPLC of the methanol extract of *Berberis aristata*

HPLC fingerprinting was established with good repeatability. Following three peaks have been shown in spectra. The compounds have been identified by comparing with the standards and also by comparing with the data reported for the *Berberis aristata* in literature. The identification has been done on the basis of Retention time and percentage height. *Berberis aristata* contains alkaloids which are berbamine, berberine, oxycanthine, epiberberine, palmatine, dehydrocaroline, jatrorrhizine and columbamine (17). The major alkaloid found in *B. aristata* is Berberine followed by Palamatine. While analyzing the components present in methanol extract of *Berberis aristata* through HPLC the three components which were identified are Berberine of 95.52%, Palmitine of 2.64% and Jatrorrhizine of 1.83% (Figure 1).




Figure 1: Spectrum for the HPLC of methanolic extract of *Berberis aristata*.

3.2 Antioxidant Activities

The DPPH free radical scavenging activity of methanolic extract of *Berberis aristata* at various concentrations was determined and compared with that of the standard BHT and Gallic acid (Table 2 and Figure 2). The methanolic extract reduced the concentration of DPPH free radical with efficiency near to that of Gallic acid but less than BHT. The IC₅₀ value for methanolic extract was 0.90mg/ml.

Nitric oxide generated from sodium nitroprusside in aqueous solution at physiological pH interacts with oxygen to produce nitrite ions which were measured by Griess Reaction. Nitric oxide radical generated from nitroprusside at physiological pH was found to be inhibited by the methanolic extract as shown in the (Table 3 and Figure 3).

The IC₅₀ values have been found to be 0.80 mg/ml, 0.65 mg/ml and 0.60mg/ml for the methanolic extract, gallic acid and BHT respectively.

Table 2: DPPH free radical scavenging assay (%) of methanolic extract of *Berberis aristata*, Gallic acid and BHT.

Conc.(mg/ml)	Extract	%	Gallic acid	%	BHT	%
1.0	0.321±0.006	58.31	0.383±0.011	58.45	0.234±0.018	66.42
0.8	0.447±0.017	41.94	0.410±0.006	55.53	0.265±0.006	61.97
0.6	0.535±0.028	30.51	0.470±0.007	49.02	0.332±0.011	52.36
0.4	0.640±0.046	16.88	0.532±0.028	42.29	0.365±0.007	47.63
0.2	0.660±0.054	14.28	0.554±0.037	39.91	0.445±0.042	36.15




Figure 2: DPPH scavenging activity for BHT, Gallic acid and methanol extract of *Berberis aristata*.

Table 3: Nitric Oxide Scavenging method (%) of methanolic extract of *Berberis aristata*, Gallic acid and BHT.

Conc.	Extract	%	Gallic acid	%	BHT	%
1.0	0.517±0.016	54.32	0.539±0.010	53.67	0.698±0.060	56.51
0.8	0.569±0.045	49.73	0.552±0.010	51.74	0.758±0.036	52.77
0.6	0.606±0.072	46.46	0.579±0.005	49.38	0.806±0.016	49.78
0.4	0.734±0.052	35.15	0.633±0.008	44.66	0.837±0.023	47.85
0.2	0.813±0.060	28.18	0.684±0.010	40.20	0.868±0.013	45.91

Values were expressed as MEAN ± S.D. (n=3)





Figure 3: Nitric Oxide scavenging activity for BHT, Gallic acid and methanolic extract of *Berberis aristata*.

3.3 Antibacterial activity

The antibacterial activity reveals that the methanolic extract of *Berberis aristata* stem is highly active against both Gram positive and Gram negative bacteria. The methanolic extract shows highest zone of inhibition (19mm) against *B.subtilis* followed by (18mm) zone of inhibition against *S.aureus* and lowest (16mm) zone of inhibition against *E.coli*. Graph 1 summarizes the microbial growth of methanolic extract and ampicillin used as a positive control.



Graph 1: Antibacterial activity (in mm) of methanolic extract and ampicillin (standard)

*0 to 25 zone of inhibition in mm; borer size=5mm




Figure 4: Antibacterial activity of *Berberis aristata* extract by agar well diffusion.




Figure 5: Antibacterial activity of Ampicillin

The MIC which is lowest concentration of a plant extract that still retained an inhibitory effect against the growth of a microorganism was assessed by using broth dilution method. The MIC values of methanol extract of *Berberis aristata* for different pathogenic bacteria ranged from (37 to 111) $\mu\text{g/ml}$.

Table 5: Intial MIC of *Berberis aristata* methanolic extract against bacteria

Conc.($\mu\text{g/ml}$)	<i>Berberis aristata</i> methanol extract ($\mu\text{g/ml}$) against bacteria		
	<i>S.aureus</i>	<i>B.subtilis</i>	<i>E.coli</i>
1000	-	-	-
333	-	-	-
111	-	*	*
37	*	+	+
12	+	+	+
MIC	37	111	111

(+)Growth, (*)MIC and (-)No growth

4. Conclusion

On the basis of results obtained it can be concluded that *Berberis aristata* stem possess alkaloids and have potent antioxidant and antibacterial activities. Further the potential of this plant can be explored more and more, in order to develop an alternative therapy for treatment of various diseases. The present study also suggests that the use of this medicinal plant may be exploited for health supplements. Thus justifying its traditional use.

References

- [1] Sharma B. D., Balakrishnan N. P., Rao R. R. and Hajra P. K. (1993). Botanical Survey of India, Calcutta. *Ranunculaceae- Barclayaceae*; (1).
- [2] Jha P. K., Choudhary R. S. and Choudhary S. K. (1997). Studies of medicinal plants of Plamau(Bihar)-(2nd part) *Biojournal.*; 9: 21-38.
- [3] Chatterjee R. P. (1951). *J Indian Chem. Soc.* 28: 225.
- [4] Blasko G. (1982). Karachine, an unusual protoberberine alkaloid. *J of American chemical society*; 104(7): 2039-2041.
- [5] Bhakuni D. S., Shoheb A. and Popali S. P. (1968). Chemical constituents of *Berberis aristata*. *Indian journal of Chemistry*; 6(2): 123.
- [6] Akhtar M. S., Sajid M. S., Ahmad M. (2008). Hypoglycemic effect of *Berberis aristata*, its aqueous and methanolic extract in normal and alloxan induced diabetic rabbits. *Pharmacology online* (Italy); 2: 845-856.
- [7] Shahid M., Rahim T., Shahzad A, Tajuddi, Latif, A., Fatma T, Rashid M, Raza Adil and Mustafa S. (2008). Ethnobotanical studies on *Berberis aristata* DC root extracts. *African Journal of Biotechnology*; (8): 556-563.
- [8] Linn J., Opoku A. R., Geheeb-Keller M., Hutchings A. D., Terblanche S. E., Jager A. K. and Staden J. V. (1999). Preliminary screening of some traditional Zulu materials for anti-inflammatory and antimicrobial activities. *Journal of Ethnopharmacology*; 68: 267-274.
- [9] Nooman A., Khalaf A., Shakya, Atif Al-Othman, Zaha El-Agbar and Farah H. (2008). Antioxidant activity of some common plants. *Turk J.Biol.*; 32: 51-55.
- [10] Balak K. N., Ponda A. B., Raj N. R, Shrivastava A. and Prathani R (2009). The evaluation of Nitric Oxide Scavenging Activity of *Acalypha indica* Linn Root. *Asian J Research chemistry*; 2: 69-74.
- [11] Ahmad I. and Beg A. J. (2001). Atimicrobial and phytochemical studies on 45 Indian medicinal plants against multidrug resistant human pathogens. *J. Ethnopharmacology*; 74: 111-123.
- [12] Srinivasan D., Nathan S., Suresh T. and Perumalsamy P.L. (2001). Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *J. Ethnopharma*; 74: 217-220.
- [13] Hammer K. A., Carson C. F. and Riley T.V. (1999). Antimicrobial activity of essential oils and other plant extracts. *J. Appl. Microbiol.*; 86: 885-990.
- [14] Junior A. and Zani C. (2000). Biological screening of Brazilian medicinal plants. *J. Braz.Sci.*; 95: 367-373.
- [15] Andrews H.J.M. (2001). Determination of minimum inhibitory concentration. *J. Antimicrob.*; 48: 5-16.
- [16] Rasooli I. and Abyanek M.R.(2004). Inhibitory effect of thyme oils on growth and aflotoxin production by *Aspergillus parasiticus*. *J. Food Cont.*; 15: 479-483.
- [17] Saied S, Batool S and Naz S (2007). Phytochemical studies of *Berberis aristata*. *J. of Basic and applied sciences*; 3 (1): 1-4.