



technique. A 1 ml volume of V-8 juice broth medium containing spores of each fungus ( $50\mu\text{l}$ ,  $10^5$  spores  $\text{ml}^{-1}$ ), was mixed with different product, separately on a sterile glass slide. After 24h of incubation at  $25\text{C}^0$  and 100% RH, spore germination was examined microscopically under an Olympus microscope (BX50+BX-FLA, Olympus Optical, Germany). The percentage of germinated spores within a count of 100 spores was determined on triplicate slides for each sample.

### 2.2.2 Evaluation of the ability of elicitors to induce the wheat defense response against pathogens under controlled conditions

Seeds of wheat of highly susceptible cultivar to diseases (cv. Giza 160) were grown in 30 cm- diameter plastic pots (2 plants per pots) under a 16 h photoperiod and 15 to  $30^0\text{C}$ . Seeds were sown in 30 cm diameter plastic pots (5 seeds per pot) and grown under a 16 h photoperiod at 22 to  $25^0\text{C}$  (Boheria governorate). The seeds were placed into insert in the sterile glass beakers containing a suspension of bioagents cells ( $1.0 \times 10^6$  /ml) for 3 h after that sown in soil fortified with biocopmost. Plants were watered every second day with water supplemented with 20N:20P:20K greenhouse fertilizer. The greenhouse experiment was designed as a randomized complete block. Soil and plant were inoculated with pathogens inoculation as follows:

Plants were sprayed with spore suspension  $10^5$  spores, according to infection stages. Uredospores of *Puccinia recondita* f. sp. *tritici* inoculations were performed on seedlings at 2-3 leaf stage. Uredospores were suspended in a light mineral oil and inoculated using concentration of  $10^5$  spores /ml with 0.01% surfactant, using a hand held atomizer (about 0.5 ml per leave). Blotch pathogen i.e. *Pyrenophora teres*, was grown separately, on V-8 juice agar at  $20^0\text{C}$ <sup>18</sup>. After seven days in culture, conidia were washed from the agar surface, filtered through a  $330\mu\text{m}$  strainer and made up into an aqueous suspension containing 10,000 conidia  $\text{mL}^{-1}$  and 0.01% Tween-80. After inoculation, seedlings were incubated overnight in complete darkness and at a relative humidity of 100%. Seedlings were then transferred to a growth chamber at  $18-22^0\text{C}$  and white fluorescent light (12 h light/12 h dark). Inoculated seedlings and plants 18-21h at greenhouse temperatures in a dark room. Each treatment was replicated three times. Seedling and plants were moved to natural light greenhouse chambers at  $22-24^0\text{C}$ , and disease responses were assessed 15 days after inoculation. The severity of rust disease was assessed as the percentage area of leaves infected during growth periods.

Disease index was scored on whole plants with ordinal rating scales. This scale is qualitative and based on lesion size and morphology. Ratings from 1-5 scale were classified as immune (0) to highly susceptible (5) according to Rosielle (Rosielle, 1972). Disease severity (R) was calculated according to the formula, having added the per cent of the affected leaf area of each leaf and having divided the sum by the number of assessed leaves:

$$\frac{R}{N} = \frac{\sum(n \cdot b)}{N}$$

- Where  $\sum(n \cdot b)$  – sum of product of the number of leaves with the same percent of severity and value of severity, N – number of assessed leaves.

### 2.3 Chemical Analysis

Ten days after inoculation, three leaves per plant were separately collected, frozen for 36 h, dried and powdered. Generally, 100 mg dried sample were used for analysis.

- **Determination of phenol content:** Free and conjugated phenols were determined in treated leaves, 15 days after plant spraying with chemical elicitors according to A.O.A.C. (1975) using the Folin–Danis reagent. Phenols were identified by HPLC using a reverse phase C8 column and compared with a catecol standard (Sigma chemicals).
- **Peroxidase and chitinase assays:** Peroxidase activity was measured according to the methods described by Allam (1972). The chitinase activity was determined by the colorimetric method of Boller and Mauch (1988). Soluble protein extraction was carried out according to Bollag and Edelstein (1992).
- **Statistical analysis:** Disease assessment results were analyzed using an ANOVA of square- root-transformed data. Data were transformed to acquire the normal distribution necessary for statistical analysis to be carried out. Significant differences were assessed by comparison of sample mean differences with the LSD value.

## 3. Results and Discussion

### 3.1 Bioassay

#### 3.1.1 Effects of products on spores germination.

*Acremonium*, *Streptomyces*, *Bacillus subtilis*, *B. megaterium*, *Pseudomonas putida*, *P. fluorescens*, *Rhodotorula glutinis*, methyl jasmonate (MJ), chitosan (CHI), an antioxidant in the form of ascorbic acid and a polyamine (PA) in the form of putrescein) were effective against pathogens germination isolated from wheat plants as showed in Table (1). The most treatment against spot diseases and rust was *Pseudomonas putida*, *P. fluorescens* in both crops. *Bacillus subtilis* was effective against powdery mildew. Methyl jasmonate and Chitosan resulted in a significantly greater decrease in the germination of all pathogens in wheat.

#### 3.1.2 Evaluation of the ability of elicitors to induce the wheat defense response against biotic stress under controlled conditions.

The effects of different compounds on control of diseases of wheat was evaluated and presented in Tables (2-3). Leaf blotches, Powdery mildew, leaf rust are the most important diseases that cause severe losses, Tables (2 & 3). Results showed that all compound have potentiality to reduce the diseases incidence in wheat. Significant differences were obtained among treatments and untreated control. Analysis of data indicated that all compounds treatments significantly reduced disease severity under artificial conditions of wheat.

Biocontrol agents were more effective in controlling all the diseases. The most treatment against spot diseases and rust was *P. putida*, *P. fluorescens* in both crops. *Bacillus subtilis* was effective against powdery mildew. *Rhodotorula glutinis* and *Trichoderma harzianum* were

more effective against head blight in wheat. At the same time, Our results clearly show that the polyamines markedly enhanced resistance to diseases control. The control plant developed a strong disease incidence and severity. This may be to increase plant defense compounds against pathogens. Overall, methyl jasmonate holds great promise for protection.

Application of methyl jasmonate significantly increased yield compared with ascorbic acid. Putrescence significantly increased yield in comparison with other compounds. Moreover, data revealed that spraying plants with ascorbic acid, had positive effect on increasing yield / plant (Table 4). Methyl jasmonate, and Chitosan and *Pseudomonas putida* resulted in a significantly greater decrease in the diseases incidence under none and saline soils in addition increased of total phenols, peroxidase, chitinase and total soluble protein in wheat as well as plant growth and yield.

Most sustainable and environmentally acceptable control may be achieved using biocontrol agents (BCAs) (Haggag Wafaa *et al.*, 2014). Though bio-control with PGPR is an acceptable green approach, the proportion of registration of biocontrol agents for commercial availability is very slow. This differential effectiveness is clearly illustrated by results from three PGPR strains (Haggag, Wafaa *et al.*, 2013). *Pseudomonas putida*, in different plant-pathogen systems. Bacterial determinants involved in the elicitation of ISR by these strains are the fluorescent siderophore pseudobactin, the outer membrane lipopolysaccharide (LPS), and the flagella of *P. putida* WCS358 (Meziane *et al.*, 2005); the pseudobactin, other iron-regulated metabolites and LPS of *P. putida* WCS374 and the LPS and iron-regulated metabolites other than pseudobactin for WCS417. The genus *Pseudomonas* belongs to the subclass of the Proteobacteria and includes mostly fluorescent Pseudomonads as well as a few non-fluorescent species. *P. putida* was more effective against die back and phomopsis. Their mode of action is extremely varied: they may directly start a lethal biological process or only suppress the bio-aggressor by competition. Sometimes they induce resistance factors in the plant. Many studies involve these bacteria. The broad spectrum nature of the isolate, even though targeted could also have potential negative outcomes. *P. putida* as that are able to improve plant growth and plant health and are implicated in the natural suppressiveness of plant pathogens (Meziane *et al.*, 2005). Because of these changes in the pathogen population, therefore, recent efforts have been directed primarily toward identifying new control measures that could be effective, reliable and safe for the environment. Recently, it has been proposed to reduce Ochratoxin A producing *Aspergillus niger* by means of biocontrol yeast, *Rhodotorula glutinis* Harrison (Yuan *et al.*, 2004). Several isolates of *R. glutinis* reduced sporulation and established pathogen and disease severity (Haggag Wafaa and, Abdall, 2012). Several yeast species are known to accumulate carotenoid pigments as secondary metabolites. *Trichoderma* spp. is biological control agent for certain fungal plant diseases (Dorner *et al.*, 2003). *Trichoderma* spp involves a complementary action of antibiosis, nutrient competition and cell wall degrading enzymes such as 1, 3-glucanases, chitinases and proteases (Choudhary, 2008). Jasmonic acid (3-oxo-2-[2'-*cis*-pentyl]-cyclopentane-1-acetate; JA) and its methyl ester, methyl jasmonate (MJ) are widespread natural regulators involved in many processes during plant

development (Thatcher, *et al.*, 2005). Jasmonate, seems to play a dual role in plant development and defense. In fact, jasmonate induced protein include antifungal protein as phenylalanine ammonia lyase, and thionin, hydroxyproline- and proline -rich cell wall proteins. Another line of evidence for the role of jasmonates in disease resistance comes from their stimulatory effect on secondary metabolite production including alkaloids, terpenes, and phenolics and polyamines (Martin *et al.*, 2002). In a previous article, a positive relationship between bean rust disease suppress and the over-accumulation of free and acid-soluble polyamine conjugates induced by a precursor of polyamines was proposed (Haggag, Wafaa, 2005). Results in this research indicate that the higher increasing in free and conjugated forms polyamines was obtained with MJ at 20 and 30 mM of both treatments which increased free and conjugated putrescine, spermidine and spermine compared with untreated control. In this respect (Walters *et al.*, 2002) Jasmonates are known to enhance the amount of phenolic compounds, the conjugation partners for polyamines, by stimulating the phenylpropanoid pathway. Thus, treatment of wheat from various concentrations led to an increase in total phenol compared with untreated plants. Increase in phenolic content in plants has been correlated with resistance to pathogens. At the same time, all treatments increased the plant height, dry weight and grains yield of wheat plants. Jasmonates are involved in many plant processes such as seed germination, fertility, senescence, and yield, as well as responses to environmental stresses, such as drought and salinity (Martin, *et al.*, 2002). In addition, methyl jasmonate can up-regulate the genes that are involved in secondary metabolism, cell-wall formation, and jasmonate biosynthesis, whereas the genes involved in photosynthesis are down-regulated (Cheong and Yang, 2003) reported that the major forms of polyamines are putrescine, spermidine and spermine, in addition to methyl jasmonate alters polyamine metabolism and induces systemic protection against powdery mildew infection in barley seedlings (Haggag 2005). Also, all concentrations of methyl jasmonate treatments increased PR- protein. In this respect, many PR proteins exhibit antifungal activity *in vitro* (Velazhahan, *et al.*, 2000). For instance,  $\beta$ -1,3-glucanases (PR-2) and chitinases (PR-3) are able to hydrolyze  $\beta$ -1,3-glucan and chitin, respectively, the major components of fungal cell walls and accumulation of PR-proteins has been correlated with systemic resistance in plants. Our research studies applying bioproducts become more integrated into management strategies in protection and curative of wheat from biotic stress. The results of this study suggested it may be possible to replace conventional chemical fungicides with natural compounds, it is safe for human, environment and thus provided both economic and ecological efficacy.

#### 4. Acknowledgments

This project was supported financially by the National Research Centre under title: **Strategies for Enhancement of wheat and Barley production under biotic and abiotic stress in Sinai**, from 2013- 2016; PI. Wafaa M. Haggag

**Table 1:** In vitro, inhibition of wheat fungal growth by natural products

Treatments	Pathogens germination %					
	Necrotrophic fungi (Blotches)				Obligate fungi	
	Net blotch	Leaf blotch	Spot blotch	Septoria	Powdery mildew (PM)	Leaf rust
<i>Acremonium</i>	0.0	0.0	3.31	0.0	2.54	6.43
<i>Streptomyces,</i>	0.87	0.0	0.93	0.0	2.40	0.0
<i>Bacillus subtilis</i>	0.0	2.5	0.0	0.0	0.0	0.0
<i>B. megaterium</i>	4.4	2.7	5.9	2.7	3.0	4.9
<i>Pseudomonas putida</i>	0.0	0.0	0.0	0.0	0.0	0.0
<i>P. fluorescens</i>	0.0	0.0	0.95	0.0	0.88	0.81
<i>Rhodotorula glutinis</i>	0.0	0.0	0.0	0.0	0.0	2.3
<i>Trichoderma harzianum</i>	0.0	0.0	0.0	0.0	0.0	0.89
Methyl jasmonate	0.0	0.0	0.0	0.0	0.0	0.0
Chitosan	0.0	0.0	0.0	0.0	0.0	0.0
Ascorbic acid	5.25	6.5	5.9	5.2	3.25	4.8
Putrescine	2.47	0.94	3.88	1.74	0.90	0.4
<b>LSD</b>	0.057	0.03	0.06	0.06	0.07	1.6

**Table 2:** Diseases severity and Concentrations of antifungal in Leaves of wheat plants treated with different products, 10 days after inoculation with pathogenic fungi

Treatments	Necrotrophic fungi (Blotches)							
	Diseases severity %				Phenol content (mg catechol/g F.W.)	Enzymes activities		Soluble protein (mg g <sup>-1</sup> F.W.)
	Net blotch	Leaf blotch	Spot blotch	Septoria		Peroxidase	chitinase	
<i>Acremonium</i>	0.0	0.0	1.35	0.0	21.8	16.5	5.7	21.9
<i>Streptomyces,</i>	0.87	0.0	0.23	0.0	19.9	17.7	4.9	22.3
<i>Bacillus subtilis</i>	0.0	2.0	0.0	0.0	28.9	20.8	5.3	21.8
<i>B. megaterium</i>	4.4	2.0	3.7	2.7	18.8	19.7	5.0	15.6
<i>Pseudomonas putida</i>	0.0	0.0	0.0	0.0	36.8	24.7	7.8	31.5
<i>P. fluorescens</i>	0.0	0.0	0.65	0.0	33.8	22.8	6.5	30.9
<i>Rhodotorula glutinis</i>	0.0	0.0	0.0	0.0	17.8	16.8	5.5	24.5
<i>Trichoderma harzianum</i>	0.0	0.0	0.0	0.0	30.8	21.8	6.8	30.8
Methyl jasmonate	0.0	0.0	0.0	0.0	41.9	26.6	7.8	36.8
Chitosan	0.0	0.0	0.0	0.0	40.8	25.7	7.3	35.5
Ascorbic acid	4.65	3.5	5.0	4.2	23.8	11.8	6.4	27.7
Putrescine	0.97	0.34	2.68	0.96	31.0	18.7	6.6	30.8
Untreated control	3.0	2.0	0.65	0.67	12.8	8.8	4.4	12.7
Infected control	80.87	70.67	90.0	50.98	8.78	4.6	3.2	7.9
<b>LSD</b>	0.07	0.02	0.05	0.06				

**Table 3:** Diseases severity and Concentrations of antifungal in Leaves of wheat plants treated with different products, 10 days after inoculation with pathogenic fungi

Treatments	Obligate fungi					
	Diseases severity %		Phenol content (mg catechol/g F.W.)	Enzymes activities (PM)		Soluble protein (mg g <sup>-1</sup> F.W.)
	Powdery mildew (PM)	Leaf rust		Peroxidase	chitinase	
<i>Acremonium</i>	2.66	3.36	21.8	16.5	5.7	21.3
<i>Streptomyces,</i>	0.77	0.0	19.9	16.6	4.6	21.0
<i>Bacillus subtilis</i>	0.0	0.0	28.9	20.3	5.1	22.8
<i>B. megaterium</i>	3.0	4.4	18.8	18.7	5.1	16.3
<i>Pseudomonas putida</i>	0.0	0.0	36.8	23.7	7.3	31.5
<i>P. fluorescens</i>	0.88	0.79	33.8	21.8	6.5	31.9
<i>Rhodotorula glutinis</i>	0.0	0.97	17.8	15.3	5.3	25.5
<i>Trichoderma harzianum</i>	0.0	0.65	30.8	20.4	6.8	31.8
Methyl jasmonate	0.0	0.0	41.9	25.4	7.3	36.0
Chitosan	0.0	0.0	40.8	25.2	7.3	35.6
Ascorbic acid	3.25	4.3	23.8	10.8	6.2	28.7
Putrescine	0.60	0.36	31.0	18.7	6.6	31.5
Untreated control	26.8	19.65	12.8	8.3	4.3	12.6
Infected control	80.87	90.0	8.78	4.2	3.2	7.9
<b>LSD</b>	0.07	1.4				

**Table 4:** Growth and yield of wheat plants treated with natural products

	Mean plant height (cm)	Mean plant dry weight (g)	Grain dry weight of 100 years (g)
<i>Acremonium</i>	712	28.6	82.5
<i>Streptomyces</i>	83.4	39.8	88.6
<i>Bacillus subtilis</i>	85.6	47.9	96.4
<i>B. megaterium</i>	80.4	39.6	60.2
<i>Pseudomonas putida</i>	80.3	39.6	88.1
<i>P. fluorescens</i>	90.2	37.6	80
<i>Rhodotorula glutinis</i>	89.5	32.5	88.6
<i>Trichoderma harzianum</i>	85.4	29.5	86.5
Methyl jasmonate	95.3	43.5	104.5
Chitosan	96.5	36.9	92.4
Ascorbic acid	82.3	31.8	80.6
Putrescine	85.4	37	88
Untreated Control	69.7	17.6	51.6
<b>L.S.D</b>	5.42	3.45	6.05

## References

- [1] A.O. A. C. (1975). Official Methods of Analysis of the Association of Official Agricultural Chemists (12th ed). Washington DC. pp 1042 Agric. Sci. 29: 892-912.
- [2] Allam A (1972). Sulfide inhibition of oxidases in rice roots. *Phytopathology* 62:634–639
- [3] Bollag DM and Eldelstein SJ (1992). Protein extraction. In: Protein Methods. Bollag DM and Eldelstein SJ (eds). Wiley-Liss Inc, New York. pp 27-42
- [4] Boller T and Mauch F (1988). Colorimetric assay for chitinase. *Methods Enzymology* 161:430-435
- [5] Cheong JJ, Choi YD (2003). Methyl jasmonate as a vital substance in plants. *Trends Genet* 19: 409–413
- [6] Choudary, D.A., Reddy, K.R.N., Reddy, M.S.(2007). Antifungal activity and genetic variability of *Trichoderma harzianum* isolates. *J. Mycol. Plant Pathol.* 37(2): 295-300.
- [7] Cotterill PJ, Rees RG, Platz GJ, Dill-Macky R (1992). Effects of leaf rust on selected Australian barleys. *Aust J Exp Agric* 32:747–75
- [8] Coventry DR, Gupta RK, Yadav A, Poswal RS, Chhokar RS, Sharma RK, Yadav VK, Gill SC, Kumar A, Mehta A, Kleemann SGL, Bonamano A, Cummins JA (2011). Wheat quality and productivity as affected by varieties and sowing time in Haryana. *India F. Cr. Res.* 123: 214–225.
- [9] Dornier, J., Cole, R., Connick, W., Daigle, D.(2003). Evaluation of biological control formulations to reduce aflatoxin contamination in peanuts, *Biological Control*, 26:318–324.
- [10] Haggag Wafaa, Abdall AM. (2012). Evaluation of *Streptomyces Aureofaciens* and *Rhodotorula glutinis* Against Ochratoxin A Producing *Aspergillus nigrin* Grapevines. *Journal of Microbiology Research* 2(6): 170-175
- [11] Haggag WM (2005). Polyamines: induction and effect on rust disease control of bean. *Plant Pathol Bull* 14:89-102
- [12] Haggag, Wafaa and Salme Timmusk (2008). Colonization of peanut roots by biofilm-forming *Paenibacillus polymyxa* initiates biocontrol against crown rot disease. *Journal of Appl Microbiol (UK)*. 104 ( 4 ): 961-969.
- [13] Haggag, Wafaa M. Wheat Diseases in Egypt and its management. *Journal of Applied Sciences Research*, 9(1): 2013, 46-50, 2013
- [14] Haggag, Wafaa, M. and Abd-El-Kareem, F. (2009). Methyl jasmonate stimulates polyamines biosynthesis and resistance against leaf rust in wheat plants. *Archives Journal of Phytopathology and Plant Protection*. German 42(1): 16–31
- [15] Haggag, Wafaa M., Malaka A. E. Saleh, Inas Mostafa, Noran Adel (2013). Mass production, fermentation, formulation of *Pseudomonas putida* for controlling of die back and phomopsis diseases on grapevine. *Advances in Bioscience and Biotechnology*, 2013, 4, 741-750.
- [16] Haggag Wafaa, Lashin, S. and Sabrey R. (2014). Development of Bioagents as Fungicides for Plant Disease Control. *International Journal of Engineering Research and Management (IJERM)*. V. 1, No.6, 195-205.
- [17] Martin D, Tholl D, Gershenzon J and Jörg J (2002). Methyl jasmonate induces traumatic resin ducts, terpenoid resin biosynthesis and terpenoid accumulation in developing xylem of Norway spruce stems. *Plant Physiol* 129:1003-1018
- [18] Meziane H., Vander Sluis I., VanLoon LC, Hofte M and Bakker P.A.H.M. (2005). Determinants of *Pseudomonas putida* WCS358 involved in inducing systemic resistance in plants. *Molecular Plant Pathology* 6: 177–185.
- [19] Mwale, V.M. E.H.C. Chilembwe, H.C. Uluko (2004). Wheat powdery mildew (*Blumeria graminis* f. sp. tritici): Damage effects and genetic resistance developed in wheat (*Triticum aestivum*). *International Research Journal of Plant Science (ISSN: 2141-5447)* Vol. 5(1) pp. 1-16, February, 2014
- [20] Rosielle, A. A. (1972). Sources of resistance in wheat to speckled leaf blotch caused by *Septoria tritici*. *Euphytica* 21: 152-161.
- [21] Thatcher JS, Owen B, Higgins VJ. 2005 The role of the jasmonate response in plant susceptibility to diverse pathogens with a range of lifestyles. *Plant Physiol.* 135(1):530-8.
- [22] Velazhahan, R., Samiyappan, R., Vidhyasekaran, P.(2000). Purification of an elicitor-inducible antifungal chitinase from suspension-cultured rice cells. - *Phytoparasitica* 28: 131-139.
- [23] Walters, D., Cowley, T. and Ann Mitchell (2002). Methyl jasmonate alters polyamine metabolism and induces systemic protection against powdery mildew infection in barley seedlings. *Journal of Experimental Botany*, 53( 369): 747-756.
- [24] Xin M, Wang X, Peng H, Yao Y, Xie C, Han Y, Ni Z, Sun Q (2012). Transcriptome comparison of susceptible and resistant wheat in response to powdery mildew infection. *Genom. Prote. Bioinfo.* 10: 94–106.
- [25] Yuan F, ShiPing T, Zheng Q, Young X, MeiXiang C.( 2004). Effect of storage environment on the growth of *Rhodotorula glutinis* and biocontrol of *Alternaria alternate* on sweet cherries. *Journal of Fruit Science*, 21: 113-115.