

Synthesis, Characterization and Biological Studies on Some Mixed Ligand La (III) Complexes

Maharudra Kekare¹, Vikas Vaidya², Jagdish Thakur³, Sunil Patil⁴, Bhushan Langi⁵

¹S.S. & L.S. Patkar College, Goregaon (W), Mumbai, Pin.-400062, Maharashtra, India

²Ramanarain Ruia College, Matunga, Mumbai, Pin.-400019, Maharashtra, India

³S.S. & L.S. Patkar College, Goregaon (W), Mumbai, Pin.-400062, Maharashtra, India

⁴C.K. Thakur A.C.S. College, New Panvel, Navi Mumbai, Pin.-410206, India

⁵Dnyanasadhana College, Thane, Pin-400604, Maharashtra, India

Abstract: Synthesis of mixed ligand Lanthanum(III) complexes of the type $[M(Q)(L)\cdot 2H_2O]$ have been carried out by using 8-hydroxy quinoline (HQ) as a primary ligand and N- and/or O- donor amino acids (HL) such as L-proline, L-valine and L-serine as secondary ligands. The characterization of metal complexes has been done on the basis of elemental analysis, electrical conductance, room temperature magnetic susceptibility measurements, spectral and thermal studies. The electrical conductance studies of the complexes indicate their non-electrolytic nature. Room temperature magnetic susceptibility measurements specify that lanthanum (III) complexes are diamagnetic in nature. Electronic absorption spectra of the complexes show intra-ligand, charge transfer and d-d transitions respectively. The thermal analysis data of the complexes indicates the presence of coordinated water molecules. FTIR spectra show bonding of the metal ion through N- and O- donor atoms of the ligand molecules. Tube dilution and Agar cup methods were implemented for study of antibacterial activity of the complexes against *S. aureus*, *C. diphtheriae*, *S. typhi* and *P. aeruginosa*. The antibacterial study was carried out by using the tetracycline as a standard antibacterial compound and it was found that, the complexes show mild activity against selected strains of micro-organisms as compared to standard tetracycline.

Keywords: mixed ligand lanthanum complexes, synthesis, characterization and biological studies

1. Introduction

Comprehensive research has been carried out for the study of mixed ligand complexes and their importance in various biological processes [1], [2]. It has been found that many ternary complexes of some metals are significant for activation of enzymes and they are used for storage as well as for transport of active materials [3]. The correlation between the stability of the metal-ligand complexes with their antimicrobial activity has been studied [4]-[7]. Antitumor activity of some mixed ligand complexes also have been reported [8]. Complexes of many metals with 8-hydroxyquinoline have been studied for their biological activity [09]-[11]. Metabolic enzymatic activities for many metal complexes of amino acids have been reported [12]. Many researchers have studied characterization, antimicrobial and toxicological activity of mixed ligand complexes of transition metals and actinide metal ions [13]-[18]. Synthesis and characterization of some transition metal complexes derived from amino acids have been reported [19].

Lanthanum complexes have been assayed for anticancer activity *in vitro* against HL-60 (human leukocytoma) cells, PC-3MIE8 (human prostate carcinoma) cells, BGC-823 (human stomach carcinoma) cells, MDA-MB-435 (human galactophore carcinoma) cells, Bel-7402 (human liver carcinoma) cells and Hela (human cervix carcinoma) cells [20]-[22]. Petra Heffetera and *et.al* also carried out the anticancer properties of the new lanthanum compound [tris(1,10-phenanthroline) lanthanum (III)] trithiocyanate

(KP772; FFC24) [23]. Irena Kostova evaluated the cytotoxic activity of the novel lanthanide complexes on HL-60 myeloid cells revealed that they are potent cytotoxic agents [24].

The author therefore thought to undertake the study of mixed ligand complexes of lanthanum, with 8-hydroxy quinoline (HQ) as a primary ligand and different amino acids (HL) such as, L-proline, L-valine and L-serine as secondary ligands. The metal complexes have been characterized by elemental analysis and various physico-chemical techniques such as molar conductance, magnetic susceptibility, electronic spectra, IR spectra and thermal studies.

2. Experimental

2.1 Materials

Analytical grade lanthanum(III) chloride heptahydrate was used as such without further purification. Amino acids, L-proline, L-valine and L-serine were obtained from E. Merck. Solvents like DMF and DMSO whenever used were distilled and purified according to standard procedures [25]-[27].

2.2 Preparation of mixed ligand complexes

The mixed ligand lanthanum(III) complexes were prepared from lanthanum chloride heptahydrate with primary ligand (HQ) and L-proline, L-valine and L-serine as secondary ligands (HL) in 1:2:1 proportion. To an aqueous solution (10 ml) of lanthanum(III) chloride heptahydrate (371 mg, 1mmol), ethanolic solution (20 ml) of 8-hydroxy quinoline

(290 mg, 2mmol) was added. The mixture was stirred and kept in a boiling water bath for 10 minutes. To this hot solution, an aqueous solution (10 ml) of amino acids (1 mmol) was added with constant stirring. The mixture was again heated in a water bath till the temperature reached to 50°C. The complexes were precipitated by raising pH of the reaction mixture by adding diluted ammonia solution. The mixture was cooled and solid complex obtained was filtered, washed with water followed by ethanol. The complexes thus prepared were dried under vacuum and were used for further studies.

2.3 Instrumentation

The C, H and N elemental analysis of the lanthanum(III) complexes were carried out on Thermo Finnigan Elemental Analyzer, Model No. FLASH EA 1112 Series at Department of Chemistry, I.I.T., Mumbai. Metal content was estimated complexometrically using standard procedure [28], [29]. Complexes were dissolved in DMF (10^{-3} M) to measure molar conductance values on an Equip-tronics Autoranging Conductivity Meter Model No.-EQ-667 with a dip type conductivity cell fitted with platinum electrodes (cell constant = 1.0 cm^{-1}). Room temperature magnetic susceptibility measurements of the complexes were made by the Guoy's method using $\text{Hg}[\text{Co}(\text{SCN})_4]$ as calibrant at Department of Chemistry, I.I.T., Mumbai. The electronic absorption spectra of all the complexes in DMF solution (10^{-4} M) in the ultraviolet and visible region were recorded on Shimadzu UV/VIS-160 Spectrophotometer. All the synthesized complexes were verified by infrared spectra in KBr disc on a Perkin-Elmer FT-IR Spectrophotometer Model 1600 in the region $4000\text{-}400 \text{ cm}^{-1}$. All the necessary precautions were taken to avoid moisture while preparing the pellets. The instrument calibration with respect to wave number and percent transmission was confirmed by recording the spectrum of standard polystyrene film. By interpreting the spectra, the various functional groups were assigned the different frequencies. Thermal Analysis (TG and DTA) were carried out in controlled nitrogen atmosphere on a Perkin-Elmer Diamond TG-DTA Instrument at Department of Chemistry, I.I.T., Mumbai by recording the change in weight of the complexes on increasing temperature up to 900°C at heating rate of 10°C per minute.

2.4 Antibacterial screening

2.4.1 Agar Cup Method

Antibacterial screening of a single compound against number of organisms or a given organism against different concentrations of the same compound can be carried out by using Agar Cup method. The authors studied the complexes by using this method as it is applicable for semisolid or liquid samples. In agar cup method, a plate of sterile nutrient agar with the desired test strain was poured to a height of about 5mm, allowed to solidify and a single cup of 8 mm diameter was cut from the center of the plate with a sterile cork borer. Thereafter the cup was filled with the sample solution of $1000 \mu\text{g}/\text{cm}^3$ concentration. The test solution was allowed to diffuse in surrounding agar by keeping in refrigerator for 10 min and the plate was incubated at 37°C for 24 hrs. The extent of inhibition of growth from the edge of the cup was

considered as a measure of the activity of the given compound. By using several plates simultaneously, the activities of several samples could be qualitatively studied.

2.4.2 Tube Dilution Method

The test compounds were subjected to *in vitro* screening against *Staphylococcus aureus*, *Corynebacterium diphtheriae*, *Salmonella typhi* and *P. aeruginosa* using Muller Hinton broth as the culture medium.


The test compound (10 mg) was dissolved in DMSO (10 cm^3) so as to prepare a stock solution of concentration $1000 \mu\text{g}/\text{cm}^3$. From this stock solution, aliquots of 5, 10, 15, 20 to , $250 \mu\text{g}/\text{cm}^3$ were obtained in test broth. Bacterial inoculums were prepared in sterilized Muller Hinton broth and incubated for 24 hrs. at 37°C . The aliquots were dispensed (5 cm^3) in each borosilicate test tube ($150 \times 20 \text{ mm}$). The bacterial inoculums 0.1 cm^3 of the desired bacterial strain (*S. aureus*, *C. diphtheriae*, *S. typhi* and *P. aeruginosa*) containing 10^6 bacteria/ cm^3 was inoculated in the tube. The tubes were incubated at 37°C for 24 hrs. and then examined for the presence or absence of the growth of the test organisms.

The lowest concentration which showed no visible growth was noted as minimum inhibitory concentration (MIC). Tetracycline was used as standard drug against Gram-positive and Gram-negative bacteria by similar screening procedure. The solvent DMSO was also tested as control to see that it did not affect the growth of the culture. MIC of tetracycline was found to be $1.5 \mu\text{g}/\text{cm}^3$ against *S. aureus*, $2.0 \mu\text{g}/\text{cm}^3$ against *C. diphtheriae*, $1.5 \mu\text{g}/\text{cm}^3$ against *S. typhi* and $8.0 \mu\text{g}/\text{cm}^3$ against *P. aeruginosa*.

3. Results and Discussion

3.1 Characterization of metal complexes

The synthesis of mixed ligand lanthanum(III) complexes may be represented as follows:



(Where, HQ is 8-hydroxy quinoline and HL is an amino acid)

All the complexes are coloured, non-hygroscopic and thermally stable solids (Table 1 and 2), indicating a strong metal-ligand bond. The complexes are insoluble in common organic solvents such as ethyl alcohol, acetone, etc. but are partially soluble in DMF and DMSO. The elemental analysis data (Table 3) of metal complexes is consistent with their general formulation as 1:2:1, mixed ligand complexes of the type $[\text{La}(\text{Q})_2(\text{L}) \cdot 2\text{H}_2\text{O}]$. The molar conductance values of the complexes in DMF at 10^{-3} M concentration are found to be $0.0012\text{-}0.0018 \text{ mhos cm}^2 \text{ mol}^{-1}$ indicating their non-electrolytic nature [30].

Table 1: Empirical Formula, Molecular Weight and Colour

Complex	Empirical Formula	Molecular Weight	Colour
---------	-------------------	------------------	--------

[La(Q) ₂ (Pro)2H ₂ O]	LaC ₂₃ H ₂₄ O ₆ N ₃	577.36	Brown
[La(Q) ₂ (Val)2H ₂ O]	LaC ₂₃ H ₂₆ O ₆ N ₃	579.09	Brown
[La(Q) ₂ (Ser)2H ₂ O]	LaC ₂₁ H ₂₂ O ₇ N ₃	567.32	Brown

Table 2: Decomposition Temperature and pH

Complex	Decomposition Temperature (°C)	pH
[La(Q) ₂ (Pro)2H ₂ O]	230	7.03
[La(Q) ₂ (Val)2H ₂ O]	225	7.07
[La(Q) ₂ (Ser)2H ₂ O]	240	6.97

Where, Q represents the deprotonated primary ligand 8-hydroxy quinoline whereas Pro, Val and Ser represent deprotonated secondary ligands, L-proline, L-valine and L-serine respectively.

Table 3: Elemental Analysis Data and Molar Conductance of Lanthanum Complexes

Complex	Elemental Analysis Found (Calculated)				Molar Conductance (Mhos cm ² mol ⁻¹)
	% M	% C	% H	% N	
[La(Q) ₂ (Pro)2H ₂ O]	24.08 (24.06)	47.80 (47.85)	4.15 (4.19)	7.24 (7.28)	0.0015
[La(Q) ₂ (Val)2H ₂ O]	23.91 (23.98)	47.60 (47.68)	4.49 (4.52)	7.29 (7.25)	0.0012
[La(Q) ₂ (Ser)2H ₂ O]	24.40 (24.48)	44.10 (44.05)	3.97 (3.91)	7.35 (7.41)	0.0018

3.2 Magnetic studies

The magnetic moment (Table 4) of the complexes was calculated from the measured magnetic susceptibilities after employing diamagnetic corrections and revealed their diamagnetic nature [31].

Table 4: Magnetic susceptibility data of Lanthanum complexes (-10^{-6} c.g.s. units)

Complex	X_g	X_m	μ_{eff} (B.M.)
[La(Q) ₂ (Pro)2H ₂ O]	-6.80×10^{-7}	-3.92×10^{-4}	Diamagnetic
[La(Q) ₂ (Val)2H ₂ O]	-6.42×10^{-7}	-3.72×10^{-4}	Diamagnetic
[La(Q) ₂ (Ser)2H ₂ O]	-6.59×10^{-7}	-3.74×10^{-4}	Diamagnetic

3.3 Electronic absorption spectra

The electronic spectra of the metal complexes in DMF were recorded in the UV-visible region. The spectra show three transitions in the range 271-275 nm ($36364-36900$ cm⁻¹), 334-339 nm ($28985-29494$ cm⁻¹) and 390-396 nm ($25253-26641$ cm⁻¹) ascribed to $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$ and charge transfer transitions from the ligands to the metal respectively [32].

3.4 Infra-red spectra

The FTIR spectra of the metal complexes were recorded in KBr discs over the range 4000-400 cm⁻¹. These spectra were complex due to presence of numerous bands with varying intensities, making the task quite difficult. However, an attempt has been made to assign some of the important bands on the basis of reported infrared spectra of several N and / or O donor ligands, 8-hydroxy quinoline and their metal complexes. An important feature of infrared spectra of the

metal complexes is the absence of band at ~ 3440 cm⁻¹ due to the O-H stretching vibration of the free O-H group of HQ. This observation leads to the conclusion that complex formation takes place by deprotonation of the hydroxyl group of HQ moiety. A strong $\nu(\text{CO})$ band observed in the range of 1112-1110 cm⁻¹ in the spectra of the complexes, indicates the presence of the 8-hydroxyquinolate group in the complexes co-ordinating through its nitrogen and oxygen atoms as uninegative bidentate ligand. The $\nu(\text{C}=\text{N})$ mode observed at 1580 cm⁻¹ in the spectrum of free HQ ligand is found to be shifted to lower wave number in the range of 1469-1460 cm⁻¹ in the spectra of the complexes, suggesting co-ordination through the tertiary nitrogen donor of HQ. The in-plane and out-of-plane deformation modes observed at ~ 500 cm⁻¹ and ~ 780 cm⁻¹ respectively, in the spectrum of HQ are shifted to higher wave numbers 410 cm⁻¹ and in the range 790-785 cm⁻¹ respectively, confirming co-ordination through the nitrogen atom of HQ with the metal ion.

A broad band observed in the region between 3282-3196 cm⁻¹ due to asymmetric and symmetric O-H stretching modes and a weak band in the range of 1580-1573 cm⁻¹ due to H-O-H bending vibrations indicating presence of a coordinated water molecule [33]-[35], further confirmed by thermal studies. The N-H asymmetric and N-H symmetric vibrations observed at ~ 3042 and ~ 2965 cm⁻¹, respectively, in the free amino acids are shifted to higher wave numbers i.e. in the range 3199-3092 cm⁻¹ and 3058-3054 cm⁻¹, respectively, in the spectra of the complexes, suggesting co-ordination of the amino group through nitrogen with the metal ion. The $\nu_{\text{asymmetric}}(\text{COO}^-)$ band of the free amino acids i.e. ~ 1590 cm⁻¹ is shifted to higher wave number, i.e. in the range 1607-1605 cm⁻¹ and the $\nu_{\text{symmetric}}(\text{COO}^-)$ mode observed at ~ 1400 cm⁻¹ in the spectra of free amino acids is found to be shifted to lower wave number in the range of 1372-1370 cm⁻¹, in the spectra of complexes indicating the co-ordination of the carboxylic acid group via oxygen with the metal ion. Coordination through the amino group of the amino acids has been further confirmed by the C-N symmetrical stretching frequency. It is observed at ~ 950 cm⁻¹ in the spectra of free amino acids and found to be shifted to lower wave numbers in the range of 915-913 cm⁻¹ in the spectra of the complexes. Some new bands of weak intensity observed in the regions around 612-608 cm⁻¹ and 410 cm⁻¹ may be ascribed to the M-O and M-N vibrations respectively. It may be noted that these vibrational bands are absent in the infra-red spectra of HQ as well as amino acids.

3.5 Thermal studies

The TG and DTA studies of the complexes have been recorded in the nitrogen atmosphere at the constant heating rate of 10 °C / minute.

Thermal study on mixed ligand lanthanum complexes in controlled nitrogen atmosphere was carried out to understand stages of decomposition and temperature range of decomposition. The most probable decomposition pattern of the complexes is proposed on the basis of the careful examination of TG and

DTA curves. The thermo analytical data is summarized in (Table 5,6 and 7).

The TG of the complexes shows that, they are thermally quite stable to varying degree. The complexes show gradual loss in weight due to decomposition by fragmentation with increasing temperature. The complexes L-proline, L-valine and L-serine show similar behaviour in TG and DTA studies. The thermogram of these complexes shows the loss in weight corresponding to two water molecules in the temperature range 122-164°C, followed by weight loss due to amino acid moiety in the temperature range 225-415°C. The final step of decomposition observed in the temperature range 610-820°C corresponds to two molecules of 8-hydroxyquinoline in the complex.

Thermal decomposition of all the lanthanum(III) complexes in inert atmosphere produces finely divided metal powder which gets transformed to their oxides spontaneously even in the presence of traces of oxygen present in nitrogen gas used in the experiment. The constant weight plateau in TG of lanthanum(III) after 820°C indicates completion of the reaction.

Table 5: Thermal Data of Lanthanum Complexes showing % Loss due to Water Molecules

Sr. No.	Complex	% Weight Loss due to Water Molecules		
		Temperature Range (°C)	Found	Calculated
1	[La(Q) ₂ (Pro) 2H ₂ O]	125-160	6.30	6.24
2	[La(Q) ₂ (Val) 2H ₂ O]	122-164	6.24	6.21
3	[La(Q) ₂ (Ser) 2H ₂ O]	130-164	6.40	6.35

Table 6: Thermal Data of Lanthanum Complexes showing % Loss due to Amino Acid

Sr. No.	Complex	% Weight Loss due to Amino Acid		
		Temperature Range (°C)	Found	Calculated
1	[La(Q) ₂ (Pro) 2H ₂ O]	230-410	19.85	19.77
2	[La(Q) ₂ (Val) 2H ₂ O]	225-410	20.10	20.05
3	[La(Q) ₂ (Ser) 2H ₂ O]	240-415	18.45	18.35

Table 7: Thermal Data of Lanthanum Complexes showing % Loss due to 8HQ Molecules

Sr. No.	Complex	% Weight Loss due to 8HQ Molecules		
		Temperature Range (°C)	Found	Calculated
1	[La(Q) ₂ (Pro) 2H ₂ O]	620-800	50.00	49.94
2	[La(Q) ₂ (Val) 2H ₂ O]	610-820	49.80	49.76
3	[La(Q) ₂ (Ser) 2H ₂ O]	630-820	50.95	50.82

On the basis of the physicochemical studies, the bonding and structure for the lanthanum complexes may be represented as.




Figure 1: Proposed Structures of [La(Q)₂(Pro) 2H₂O]




Figure 2: Proposed Structures of [La(Q)₂(Val) 2H₂O]




Figure 3: Proposed Structures of [La(Q)₂(Ser) 2H₂O]

3.6 Biological studies

All the metal complexes were screened against *Staphylococcus aureus*, *Corynebacterium diphtheriae*, *Salmonella typhi* and *P. aeruginosa*.

The studies based on agar cup method revealed that the complexes are more sensitive against *S. aureus* and *S. typhi* while less sensitive against *C. diphtheria* and *P. aeruginosa* (Table 8). The minimum inhibitory concentration (MIC) of ligand and the metal salts ranges between 50 and 300 µg/mL while that of metal complexes ranges between 5 and 35 µg/mL (Table 9). The complexes are found to be more active against *S. aureus* and *S. typhi* as compared to *C. diphtheria* and *P. aeruginosa*. As compared to standard antibacterial compound tetracycline, the complexes show moderate activity against selected strains of microorganisms.

The results show that, as compared to the activity of metal salts and free ligand, the metal complexes show higher activity. The activity of metal complexes is enhanced due to chelation. The chelation reduces considerably the polarity of the metal ions in the complexes, which in turn increases the hydrophobic character of the chelate and thus enables its permeation through the lipid layer of microorganisms [36].

Table 8: Antibacterial Activity (mm) of Lanthanum Complexes by Agar Cup Method

Complex	Antibacterial Activity (mm) with			
	<i>S. aureus</i>	<i>C. diphtheriae</i>	<i>S. typhi</i>	<i>P. aeruginosa</i>
[La(Q) ₂ (Pro) 2H ₂ O]	24	15	23	12
[La(Q) ₂ (Val) 2H ₂ O]	26	16	23	13
[La(Q) ₂ (Ser) 2H ₂ O]	28	14	22	13

Table 9: MIC (µg/mL) Data of Lanthanum Complexes

Complex	MIC ($\mu\text{g/mL}$)			
	<i>S. aureus</i>	<i>C. diphtheriae</i>	<i>S. typhi</i>	<i>P. aeruginosa</i>
[La(Q) ₂ (Pro) 2H ₂ O]	10	20	15	25
[La(Q) ₂ (Val) 2H ₂ O]	10	25	15	35
[La(Q) ₂ (Ser) 2H ₂ O]	05	20	20	35
8-hydroxyquinoline	50	200	150	250
Tetracycline	1.5	2.0	1.5	8.0

4. Conclusions

Based on the above results, the following conclusions may be drawn. The higher decomposition temperatures of the complexes indicate a strong metal-ligand bond and electrical conductance studies show non-electrolytic nature of the complexes, respectively. Magnetic studies indicate diamagnetic nature of the complexes. Electronic absorption spectra of the complexes show intra ligand and charge transfer transitions, respectively. IR spectra show bonding of the metal ion through N- and O-donor atoms of the two ligands. Thermal analysis confirms the presence of two coordinated water molecules. On the basis of the above results, coordination number eight is proposed for lanthanum complexes. The antibacterial study shows that complexes are found to be more active against *S. aureus* and *S. typhi* as compared to *C. diphtheria* and *P. aeruginosa*. Compared to standard antibacterial compound, tetracycline, the complexes show mild activity against the selected strains of microorganisms.

5. Future Scope

The present study can be useful in pharmaceutical field by studying cytotoxic activities of these complexes.

References

- [1] D.P. Meller and L. Maley, "Order of Stability of Metal Complexes," Nature (London), (161), pp. 436-437, 1948.
- [2] P.V. Khadikar, R. Saxena, T. Khaddar and M.A. Feraqui, "Salicylhydroxamic acid and their metal chelate as antifungal agents," Journal of Indian Chemical Society, (56), pp. 215-219, 1994.
- [3] M.N. Hughes, Comprehensive Coordination Chemistry, G. Wilkinson, R.D. Gillard and J.A. McCleverty Eds., Pergamon Press, Oxford, (6), pp. 541, 1987.
- [4] M.S. Islam, M.B. Hossain and M.Y. Reza, "Antimicrobial Studies of Mixed Ligand Transition Metal Complexes of Maleic acid and Heterocyclic Amine Bases," Journal of Medicinal Science, 3 (4), pp. 289-293, 2003.
- [5] M. M. Mashaly, T. M. Ismail, S. B. El-Maraghy, and H. A. Habib, "Heteronuclear complexes of oxorhenium(V) with Fe(III), Co(II), Ni(II), Cu(II), Cd(II) and UO₂(VI) and their biological activities," Journal of Coordination Chemistry, 57 (13), pp. 1099-1123, 2004.
- [6] S.I. Mostafa, Mixed ligand complexes with 2-piperidine-carboxylic acid as primary ligand and ethylene diamine, 2,2'-bipyridyl, 1,10-phenanthroline and 2(2'-pyridyl)quinoxaline as secondary ligands: preparation, characterization and biological activity Transition Metal Chemistry, 32 (6), 769, 2007.
- [7] Z.H. Abd El-Wahab, "Mixed ligand complexes of nickel(II) and cerium(III) ions with 4-(-3-methoxy-4-hydroxybenzylideneamino)-1,3-dimethyl-2,6-pyrimidine-dione and some nitrogen/oxygen donor ligands," Journal of Co-ordination of Chemistry, 61, 3284, 2008.
- [8] M. Galanski, M.A. Jakupcic and B.K. Keppler, "Update of the preclinical situation of anticancer platinum complexes: novel design strategies and innovative analytical approaches," Current Medicinal Chemistry, 12 (18), pp. 2075-2094, 2005.
- [9] E. Howard-Lock and C.J.L. Lock, "Comprehensive Coordination Chemistry", G. Wilkinson, R.D. Gillard and J.A. McCleverty Eds., Pergamon Press, Oxford, (6), pp. 755, 1987.
- [10] J.R. Thakkar and N.V. Thakkar, "Synthesis and Characterization of Chiral Mixed Ligand Co(II) Complexes of Isonitrosopropiophenone and Amino Acids," Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry, 30, 1871, 2000.
- [11] V.S. Shivankar and N.V. Thakkar, Acta Poloniae Pharmaceutica - Drug Research, 60, 45, 2003.
- [12] D.D. Perrin and R.P. Agarwal, "Metal Ions in Biological Systems," Ed. H.C. Sigel, Marcel Dekker, New York, (2), pp. 167, 1973.
- [13] M.R. Mahmoud, A.A. Abdel Gaber, A.A. Boraie and E.M. Abdalla, "Divalent transition metal ion mixed ligand complexes with aliphatic dicarboxylic acids and imidazoles Transition Metal Chemistry," 19 (4), pp. 435-438, 1994.
- [14] P.R. Reddy and A.M. Reddy, "Synthesis and characterization of mixed ligand complexes of bio-metals with pyrimidine nucleoside (uridine) and amino acids," Indian Academy of Sciences (Chemical Sciences), 112, (6), pp. 593-600, 2000.
- [15] A. Romerosa, P. Bergamini and V. Bertolasi, "Biologically active platinum complexes containing 8-thiotheophylline and 8-(methylthio)theophylline", Inorganic Chemistry, 43 (3), pp. 905-913, 2004.
- [16] R.K. Agarwal and S. Prasad, "Synthesis, spectral and thermal investigations of some mixed ligand complexes of thorium(IV) derived from semicarbazones and diphenyl sulfoxide", Journal of Iranian Chemical Society, 2 (2), pp. 168-175, 2005.
- [17] R.S. Joseyphus, C.J. Dhanaraj and M.S. Nair, "Synthesis and characterization of some Schiff base transition metal complexes derived from vanillin and L(+)-alanine", Transition Metal Chemistry, 31 (6), pp. 99-702, 2006.
- [18] S.I. Mostafa and N. Hadjiliadis, "New biologically active transition metal complexes of 2-mercapto-4,6-diamino-5-hydroxypyrimidine," Inorganic Chemistry: An Indian Journal, 2(3), pp. 186, 2007.
- [19] S.A. Lahsani, R.A. Ammar, M.F. Amin and E.M. Shoukry, "Mixed-ligand complex formation of Cu(II) with 1,2-diphenylethylenediamine as primary ligand and amino acids as secondary ligands", International Journal of Electrochemical Science, 7, pp. 7699 - 7711, 2012.
- [20] Zhong-Ming Wang, Hua-Kuan Lin, Shou-Rong Zhu, Tian-Fu Liu, Yun-Ti Chen, "Spectroscopy, cytotoxicity and DNA-binding of the lanthanum(III) complex of an l-

- valine derivative of 1,10-phenanthroline”, *Journal of Inorganic Biochemistry*, 89 (1–2), pp. 97-106, 2002.
- [21] Feng-Hua Lia, Guang-Hua Zhao, c, , Hong-Xing Wu, Hai Linb, Xiang-Xia Wu, Shou-Rong Zhua, Hua-Kuan Lina, “Synthesis, characterization and biological activity of lanthanum(III) complexes containing 2-methylene-1,10-phenanthroline units bridged by aliphatic diamines”, *Journal of Inorganic Biochemistry* 100 (1), pp. 36-43, 2006.
- [22] Guanghua Zhao, Fenghua Lia, Hai Linb, Huakuan Lin, “Synthesis, characterization and biological activity of complexes of lanthanum(III) with 2-(1'-phenyl-2'-carboxyl-3'-aza-*n*-butyl)-1,10-phenanthroline and 2-(1'-*p*-phenol-2'-carboxyl-3'-aza-*n*-butyl)-1,10-phenanthroline”, *Bioorganic & Medicinal Chemistry*, 15 (1), pp. 533-540, 2007.
- [23] Petra Heffetera, Michael A. Jakupcb, Wilfried Körnerc, Stefan Wildd, Nikolai Graf von Keyserlingke, Leonilla Elblinga, Haralabos Zorbasd, Alla Korynevskaf, Siegfried Knasmüllera, Hedwig Sutterlütya, Michael Mickschea, Bernhard K. Kepplerb, Walter Bergera, “Anticancer activity of the lanthanum compound [tris(1,10-phenanthroline)lanthanum(III)]trithiocyanate (KP772; FFC24)”, *Biochemical Pharmacology* 71 (4), pp. 426-440, 2006.
- [24] Irena Kostova, Raina Kostova, Georgi Momekov, Natasha Trendafilova, Margarita Karaivanova, “Antineoplastic activity of new lanthanide (cerium, lanthanum and neodymium) complex compounds”, *Journal of Trace Elements in Medicine and Biology*, 18 (3), pp. 219-226, 2005.
- [25] Vogel A I, *Quantitative Inorganic Analysis*, 4th Ed. ELBS, 1965.
- [26] Vogel A I, *Textbook of Practical Organic Chemistry*, Longmans Green and Co. Ltd., London, 5th Ed., 1989.
- [27] Vogel A I, *Textbook of Quantitative Inorganic Analysis*, Longmans Green and Co. UK Ltd., 5th Ed., 1989.
- [28] M. Ionashiro, CAF Graner and J.Z. Netto, “Complexometric titration of lanthanides and yttrium”, *Ecletica Quimica*, 8, pp. 29-32, 1983.
- [29] Z. Yaquin and Y. Binsheng, “Spectral studies on the interaction between lanthanum ion and the ligand: *N,N*-ethylenebis-[2-(*o*-hydroxyphenolic)glycine]”, *spectrochimica Acta Part A*, 62, pp. 641-644, 2005.
- [30] Geary W J, “The use of conductivity measurements in organic solvents for the characterization of coordination compounds”, *Coordination Chemistry Reviews*, 7, pp. 81-122, 1971.
- [31] G. A. Thakur and M. M. Shaikh, “Synthesis, characterization, antibacterial and cytotoxicity studies on some mixed ligand Th(IV) complexes,” *Acta Poloniae Pharmaceutica*, 63 (2), pp. 95–100, (2006).
- [32] H. Beraldo, S.M. Kainsler, J.D. Turner, I.S. Billeh, J.S. Ives, D.X. West, *Transition Metal Chemistry*, Vol. 22, Ed R. L. Carlin, Marcel Decker Inc., New York, (1997).
- [33] K. Nakamoto, *Lattice Water and Aquo and Hydroxo Complexes in Infrared and Raman Spectra of Inorganic and Co-ordination Compounds*, 4th Ed. John-Wiley and Sons, New York, 1986.
- [34] G.A. Thakur, M.M. Shaikh, “Synthesis, characterization, antibacterial and cytotoxicity studies on some mixed ligand Th(IV) complexes”, *Acta Poloniae Pharmaceutica – Drug Research*, 63 (2), pp. 95-100, 2006.
- [35] G.A. Thakur, S.R. Dharwadkar, M.M. Shaikh, *Thermal Study on Mixed Ligand Thorium (IV) Complexes*, *Proceedings of the 15th National Symposium on Thermal Analysis (THERMANS 2006)*, pp. 399, 2006.
- [36] Z.H. Chohan, A.K. Misbahul, and M. Moazzam, “Synthesis, characterization and antimicrobial studies of Co(II) and Ni(II) complexes with some pyrazoles,” *Indian Journal of Chemistry A*, (27), p. 1102, 1988.