





















		<i>Cynodon dactylon</i> (L.) Garikhe hullu (Poaceae)	Plant extract is taken orally once in every 20min, also applied onto snake bite region.
		<i>Calotropis procera</i> (Aiton) Biliekke (Asclepiadaceae)	Latex is applied on snake bite region and also 21 young leaves are taken orally immediately after snake bite.
		<i>Sapindus laurifolia</i> (Vahl) Antuvala mara (Sapindaceae)	Fruits are crushed in water and taken orally once in every 30min.
		<i>Adhatoda zeylanica</i> (Medikus) Aadusoge (Acanthaceae)	Root paste 12-24g is mixed with water and taken orally once in 60min.
		<i>Rauwolfia serpentina</i> (L.) Sarpagandhi (Apocyanaceae)	Root extract is mixed with lemon juice and taken orally and also used to apply on snakebite region.
65	Scorpion sting	<i>Strychnos potatorum</i> (L.f.) Chillada mara (Strychnaceae)	Seed paste is applied on scorpion bite.
		<i>Madhuca longifolia</i> (L.) Hippe mara, ippe mara (Sapotaceae)	Flowers are crushed in water to make paste and applied on scorpion bite.
		<i>Ocimum gratissimum</i> (L.) Kadu Tulsi / rama tulsi (Lamiaceae)	Leaf extract is applied on scorpion bite.
		<i>Achyranthus aspera</i> (L.) Utrani (Amaranthaceae)	Leaf paste of <i>A. aspera</i> and seed powder of <i>S. potatorum</i> is mixed and applied on scorpion bite.
		<i>Strychnos potatorum</i> (L.f.) Chillada mara (Strychnaceae)	
		<i>Aloe vera</i> (L.) Kathligida, lolesara (Liliaceae)	Root is crushed inside the mouth.
66	Rat bite	<i>Strychnos potatorum</i> (L.f.) Chillada mara (Strychnaceae)	Seed powder is mixed with butter milk and taken orally, once in a day early in the morning for 1 week.
67	Dog bite	<i>Calotropis procera</i> (Aiton) Biliekke (Asclepiadaceae)	Latex is mixed with equal amount of jaggery and lin seed oil and applied on dog bite region, once daily for 1 week.
		<i>Calotropis procera</i> (Aiton) Biliekke (Asclepiadaceae)	Latex is applied on dog bite region and rubbed hardly.
		<i>Aloe vera</i> (L.) Kathligida, lolesara (Liliaceae)	Leaf paste is mixed with salt and applied on dog bite region for 7 days.
68	Menstruation problems	<i>Asperagus racemosus</i> (Willd.) Shatavari (Liliaceae)	Root is crushed to make decoction and equal amount of cow's milk is added and taken orally twice a day for 10-14 days
		<i>Ficus racemosa</i> (L.) Atthi mara (Moraceae)	Fruit extract (48g), <i>A. racemosus</i> (24g) mixed well and crushed along with (10g) sugar and taken orally twice a day for a week, after milk should be drink. Or
			Root is crushed along with <i>F. racemosa</i> bark and add little amount of honey, taken orally once in a day at night for 3-4 weeks.
		<i>Acacia ferruginea</i> (DC.) Banni mara (Mimosaceae)	Leaves (50g) are crushed in cow's milk and filtered then (6g) <i>Cuminum cyminum</i> , (6g) stone sugar, taken orally daily early in the morning once, for 7-14 days.
69	Leucorrhia	<i>Aegle mormelos</i> (L.) Bilwapatre (Rutaceae)	Leaf extract 30-60ml mixed with cow's milk taken orally before breakfast, twice a day for 2 weeks.
70	Macorrhagia	<i>Adhatoda zeylanica</i> (Medikus) Aadusoge (Acanthaceae)	Leaf extract 12-18g is mixed with sugar taken orally twice a day until cure.
71	Dysmenorrhea	<i>Corallocarpus epigaeus</i> (Rottler) Gili moothi gedde (Cucurbitaceae)	<i>Corallocarpus epigaeus</i> Corm is crushed and made into small pills taken twice a day every month thrice.
72	Bearing for male child	<i>Limonia acidissima</i> L. Belada mara (Rutaceae)	Fruit pulp 38-46g is mixed with 200ml of cow's milk taken orally twice a day from the third month of pregnancy.
		<i>Ficus benghalensis</i> (L.) Alada mara (Moraceae)	Root is grown in north direction is collected, and then tied to the shoulder of pregnant.
73	Soma roga	<i>Phyllanthus amarus</i> (Schumach.) Nelanelli, kallenchi (Euphorbiaceae)	Leaf extract (60ml) is mixed with 100ml of cow's milk taken orally twice a day for 7-9 days.
74	Infertility	<i>Ficus racemosa</i> (L.) Atthi mara (Moraceae)	3-4 fruits are taken orally along with honey after sugared milk is taken orally for 40 days.
		<i>Diplocyclos palmatus</i> (L.) Mahalinganaballi (Cucurbitaceae)	Seed powder is taken orally along with water for 20-30 days.
75	Veginal problem	<i>Stachytarpheta jamaicensis</i> (L.) Kari utrani (Verbanaceae)	Root (12g) is crushed in 12g of ghee taken orally twice a day for 1-2 weeks.
76	For abortion	<i>Ficus racemosa</i> (L.) Atthi mara (Moraceae)	Root (24g) bark decoction is mixed with honey, taken orally early in the morning for 2-3 weeks.
77	Garbha hulu nivarane	<i>Aristolochia bracteolata</i> (Lam.) Katthe kirubana gida (Aristolochiaceae)	Leaf is crushed with ginger, taken orally along with honey for 3-4 weeks.
78	Labour pain	<i>Aegle mormelos</i> (L.) Bilwapatre (Rutaceae)	Fruit pulp is mixed with ginger with equal proportion to make decoction, taken orally twice a day, for 7-8 days.
79	During pregnancy	<i>Balanites roxburghii</i> (Planchon) Ingalara (Simaroubaceae)	Root bark of the tree is crushed with children urine for 3-5 days.
80	Contraception	<i>Abrus precatorius</i> (L.) Gulagangi (Fabaceae)	Seed (4g) powder is mixed with butter milk and taken orally after 3 days of periods.

81	Lactation problems (to increase breast milk)	<i>Phoenix sylvestris</i> (L.) Eechalu mara (Aracaceae)	Copra is taken orally.
		<i>Olax scandens</i> (Roxb.) Nakkare, nakkaraki (Olacaceae)	Roots and leaves are crushed and taken orally.
82	To stop lactation	<i>Azadirachta indica</i> (A.Juss.) Bevina mara (Meliaceae)	Leaf paste is applied over the breast nipples.
83	Sperm dysfunction	<i>Achyranthus aspera</i> (L.) Utrani (Amaranthaceae)	Root powder (2.5g) of plant is mixed with sugar and milk, taken orally twice a day until cure.
		<i>Ficus racemosa</i> (L.) Atthi mara (Moraceae)	Bark (6g) of <i>F.racemosa</i> , twig of <i>F.bengalensis</i> and (6g) of stone sugar is mixed in cow's milk taken orally early in the morning for 40 days.
		<i>Ficus benghalensis</i> (L.) Alada mara (Moraceae)	
		<i>Tinospora cordifolia</i> (Willd.) Amruta balli (Menispermaceae)	Leaf extract (30ml) is mixed with (30ml) of honey, taken orally twice a day for 3-4 weeks.
84	Male infertility	<i>Ocimum gratissimum</i> (L.) Kadu Tulsi / rama tulsi (Lamiaceae)	Equal quantity of Seed powder and jiggery mixed well and taken orally twice a day, for 21-42 days.
85	Anti aging	<i>Tinospora cordifolia</i> (Willd.) Amruta balli (Menispermaceae)	Plant powder is mixed with jiggery, honey; ghee mixture is taken orally twice a day along with cow's milk.
86	Early ejaculation	<i>Tamarindus indica</i> L. Hunase mara (Caesalpinaceae)	Seeds (960g) are soaked in water for 4 days, then it is fried and remove seed coat, then it is finely powdered, to that equal amount of sugar is added, taken orally 3-6g along with honey and ghee.
87	Vaajikarana	<i>Terminalia chebula</i> (Retz.) Alalekai (Combretaceae)	Seed powder is taken orally along with sugar or ginger or salt or honey.
88	Sleeplessness	<i>Rauvolfia serpentina</i> (L.) Sarpagandhi (Apocyanaceae)	Rice is mixed with water and kept for overnight, then that water is mixed with leaves of <i>R.serpentina</i> and crushed, taken orally once in a day for 1-2 weeks.
		<i>Centella asiatica</i> (L.) Ondelaga (Apiaceae)	Root is crushed and the paste is applied over top of the head and tied with the help of cloth until cure.
89	High Bp	<i>Rauvolfia serpentina</i> (L.) Sarpagandhi (Apocyanaceae)	Root is powdered and taken orally twice a day along with lemon extract and honey for 3-4 weeks.
		<i>Cassia auriculata</i> (L.) Honnava (Caesalpinaceae)	Flower paste is made into small pills and taken orally twice a day.
90	Vomiting	<i>Ocimum gratissimum</i> (L.) Kadu Tulsi / rama tulsi (Lamiaceae)	15ml of leaf extract and 2g of cardamom powder is mixed and taken orally until cure.
		<i>Tinospora cordifolia</i> (Willd.) Amruta balli (Menispermaceae)	120g of cold decoction of plant is mixed with 24g of honey taken orally twice a day for 1 week.
91	Blood vomiting	<i>Adhatoda zeylanica</i> (Medikus) Aadusoge (Acanthaceae)	Root decoction is cooled and honey is mixed, taken orally until cure.
92	Emetic	<i>Murraya koenigii</i> (L.) Gandhabevu, karibevu (Rutaceae)	Bark or the leaf extract is mixed with water and taken orally about 200-300ml.
93	Herpes	<i>Pterocarpus santalinus</i> (L.f.) Rakta chandana (Fabaceae)	Bark is crushed to make decoction and taken orally about 12g for 20days.
		<i>Cynodon dactylon</i> (L.) Garike hullu (Poaceae)	Plant is crushed in cold water and applied on affected region until cure.
94	Impotency	<i>Securinega virosa</i> (Willd.) Bilihuli (Euphorbiaceae)	Leaf is crushed and make into small pills, taken orally twice a day for 15 days.
		<i>Ipomea batatas</i> (L.) Adavi genasu (Convolvulaceae)	Tubers are taken orally.
95	Nerve weakness	<i>Holoptelea intergrifolia</i> (Roxb.) Tapsi mara (Ulmaceae)	Oil extracted from the bark is applied on affected area and gently massaged.
96	Arthritis	<i>Opuntia vulgaris</i> (Miller) Mullugalli (Cactaceae)	Latex is applied on affected area.
		<i>Abutilon indicum</i> (L.) Shreemudre (Malvaceae)	Oil is applied on painful region and gently massaged.
97	Cancer	<i>Hemidesmus indicus</i> (L.) Haluballi, halikudi (Periplocaceae)	Whole plant is crushed mixed with equal amount of sugar, taken orally along with crystals of salt.
98	Lungs problem	<i>Boswellia serrata</i> (Roxb.) Sambranimara, lobana gida (Burseraceae)	Dried latex is put into the fire and the smoke is inhaled through nose, daily once a day in early morning.

#### 4. Acknowledgement

The authors sincerely acknowledge the kind support of Dr. Gayathri Devaraja, Associate professor, DOS in Microbiology, Davangere University, India for encouraging to this work. The authors express their sincere thanks to the

local herbal practitioners. Authors are highly obliged with the help rendered by Panchavati Research Academy for Nature, Kalamangi, Karnataka and valuable suggestions.

## References

- [1] Anonymous. 2002. *Wealth of India*. 3<sup>rd</sup> ed., No.3, C.S.I.R., New Delhi, p.267.
- [2] Bhandary, M.J., Chandrashekar, K.R. and Kaveriappa, K.M. 1995. Medical ethnobotany of the Siddis of Uttara Kannada district, Karnataka, India. *Journal of Ethnopharmacology*, 47: 149-158.
- [3] Chopra, L.C. and Nayar, M.C. 1956. Glossary of Indian medicinal plants. Council of scientific and industrial research, New Delhi; 1956. *Current Science*, 43(23): 749-750.
- [4] Das, A. and Tag, H. 2006. Ethnomedical studies of Khamti tribe of Arunachal Pradesh. *Indian Journal of traditional Knowledge*, 5(3): 317-322.
- [5] Dubey, N.K., Kumar, R. and Tripathi, P. 2004. Global promotion of herbal medicine: India's opportunity, *Current Science*, 86 (1): 37-41.
- [6] Gamble, J.S. 1995. *Flora of the presidency of Madras*. Pages 2017, 3 Volumes, Mahendra Pal Singh Publications, Dehra Dun.
- [7] Harsha, V.H. Hebbar, S.S., Hegde, G.R. and Shripathi, V. 2002. Ethnomedical Knowledge of Plants used by Kunabi Tribe, Karnataka. India. *Fitoterapia*, 73: 281-287.
- [8] Harsha, V.H., Hebbar, S.S., Shripathi, V. and Hegde, G.R. 2003. Ethnomedicobotany of Uttar Kannada District in Karnataka, India. Plants in Treatment of Skin Disease. *Journal of Ethnopharmacology*, 84: 37-40.
- [9] Harsha, V.H., Shripathi, V. and Hegde, G.R. 2005. Ethnoveterinary practices in Uttar Kannada District of Karnataka. *Indian Journal of Traditional Knowledge*, 4: 235-258.
- [10] Kirtikar, K.R. and Basu, B.D. 1996. Indian medicinal plants, 2<sup>nd</sup> ed., International Book Distributors, Dehra Dun, India, 2: 791.
- [11] Parinitha, M., Harish, G.U., Vivek, N.C., Mahesh, T. and Shivanna, M.B. 2004. Ethno-botanical wealth of Bhadra Wildlife Sanctuary in Karnataka. *Indian journal of Traditional knowledge*, 3: 37-50.
- [12] Parinitha, M., Shrinivasa, B.S. and Shivanna, M.B. 2005. Medicinal plant wealth of local communities in some villages in Shimoga district of Karnataka, *Indian Journal of Ethnopharmacology*, 98: 307-312.
- [13] Pei. 2001. Ethnobotanical approaches of traditional medicine studies on some experiences from Asia. *Pharmacology and Biology*, 39: 74-79.
- [14] Pradeep B, Ganesh H. R, Gurumurthi H, Gangadhar M, S. 2014. Ethnomedicinal Plants to cure skin Diseases- an Account of the Traditional Knowledge in the Coastal Parts of Central Western Ghats, Karnataka, India. *Journal of Ethnopharmacology*, 151: 493-502.
- [15] Prakash B.N. and Unnikrishnan P.M. (2013). Ethnomedical Survey of Herbs for the Management of Malaria in Karnataka, India. *Journal of Ethnobotany Research and Applications*, 11 :289-298.
- [16] Rajkumar, N. and Shivanna, M.B. 2010. Traditional herbal medicinal knowledge in Sagar taluk of Shimoga district, Karnataka, India. *Indian journal of Natural Product Resources*, 1(1): 102-108.
- [17] Rajkumar, N. and Shivanna, M.B. 2012. Traditional veterinary healthcare practices in Shimoga district of Karnataka, India. *Indian Journal of traditional Knowledge*. 11(2): 283-287.
- [18] Raju Kanti and Parashurama T.R. 2014. Ethno-Botanical Knowledge Health care in Hangal Taluk of Haveri district, Karnataka. *International Journal of Innovative Pharmaceutical Sciences and Research*. 2(10):2344-235.
- [19] Ramachandra Naik, M., Vaishnavi V, Preethi K, and Krishnamurthy, Y.L. (2012). Ethnoveterinary uses of Medicinal Plants among the Lambani Community in Chitradurga District, Karnataka, India. *Asian Pacific Journal of Tropical Biomedicine*, S470-S476.
- [20] Shivanna, M.B., Mangala, K.R. and Parinitha M. 2008. Ethno-medicinal knowledge of Lambani community in Chikmagalur district of Karnataka, India. *Journal of Medicinal and Aromatic Plant Sciences*. 30: 105-108.