

On The Non Homogeneous Heptic Equation with Five Unknowns $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^5$

P. Jayakumar¹, K. Sangeetha²

¹Department of Mathematics, A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur (District) – 613 503, Tamilnadu India

Abstract: The non homogeneous Diophantine equation of degree seven with five unknowns represented by $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^5$ is analyzed for its non - zero distinct integer solutions. Employing suitable linear transformations and applying the method of cross multiplication, four different patterns of non-zero distinct integer solutions to the heptic equation under consideration are obtained. A few interesting relation between the solutions and special numbers namely Polygonal numbers, Pyramidal numbers, Centered Pyramidal numbers, Star numbers and Stella octangular numbers are exhibited.

Keywords: The non homogeneous Diophantine equation, Heptic equation with five unknowns, integral solutions, special numbers, a few interesting relation

2010 Mathematics Subject Classification: 11D09

Notations: P_n^m = Pyramid number of rank n with size m
 $T_{m,n}$ = Polygonal number of rank n with size m
 gn_a = Gnomonic number of rank a
 So_n = Stella octangular number of rank n
 Pr_n = Pronic number of rank n
 $CP_{m,n}$ = Centered Pyramidal number of rank n with size m.

1. Introduction

The Diophantine equations, homogeneous and non-homogeneous have aroused the interest of numerous mathematicians since antiquity as can be seen from [1, 9] the problem of finding all integer solutions of a Diophantine equation with three or more variables and degree at least three, in general presents a good deal of difficulties. There is vast general theory of homogeneous quadratic equations with three variables [1-4, 9]. Cubic equations in two variables fall in to the theory of elliptic curves which is a very developed theory but still an important topic of current research [5-7]. A lot is known about equations in two variables in higher degrees. For equations with more than three variables and degree at least three very little is known. It is worth to note that undesirability appears in equations, even perhaps at degree four with fairly small coefficients. In [1, 8 and 9] a few higher order equations are considered for integral solutions. In this communication a seventh degree non-homogeneous equation with five variables represented by $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^5$ is considered and in particular a few interesting relations among the solutions among the solutions are presented.

2. Method of Analysis

The Diophantine equation representing the heptic equation with five unknowns under consideration is $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)z^5$ (1)

Introduction of the linear transformation

$$x = u + v, y = u - v, X = 2u + v, Y = 2u - v \quad (2)$$

in (1) leads to $u^2 + 17v^2 = 21z^5$ (3)

Now we solve (3) through different methods and thus obtain different patterns of solutions to (1)

2.1 Pattern: 1

Assume $z = z(a, b) = a^2 + 17b^2$ (4)

Where a and b are non zero distinct integers

Write 21 as

$$21 = (2 + i\sqrt{17})(2 - i\sqrt{17}) \quad (5)$$

Using (4) and (5) in (3) and applying the method of factorization, define

$$(u + i\sqrt{17}v) = (2 + i\sqrt{17})(a + i\sqrt{17}b)^5$$

Equating real and imaginary parts, we get

$$u = u(a, b) = 2a^5 - 85a^4b - 340a^3b^2 + 2890a^2b^3 + 2890ab^4 - 4913b^5$$

$$v = v(a, b) = a^5 + 10a^4b - 170a^3b^2 - 340a^2b^3 + 1445ab^4 + 578b^5$$

Hence in view of (2) the corresponding solutions of (1) are given by

$$x = x(a, b) = 3a^5 - 75a^4b - 510a^3b^2 + 2550a^2b^3 + 4335ab^4 - 4335b^5$$

$$y = y(a, b) = a^5 - 95a^4b - 170a^3b^2 + 3230a^2b^3 + 1445ab^4 - 5491b^5$$

$$X = X(a, b) = 5a^5 - 160a^4b - 850a^3b^2 + 5440a^2b^3 + 7225ab^4 - 9248b^5$$

$$Y = Y(a, b) = 3a^5 - 180a^4b - 510a^3b^2 + 612a^2b^3 + 4335ab^4 - 10404b^5$$

$$z = z(a, b) = a^2 + 17b^2$$

Properties:

1. $3X(a, 1) - 5Y(a, 1) = 420Pr_{a^2} - 367T_{4, 2a} + 24276$
2. $x(a, 1) - 3y(a, 1) - 210Pr_{a^2} \equiv 12138 \pmod{94620}$
3. $10z(a, a)$ a Nasty number
4. $x(a, 1) - 3y(a, 1) + 10490T_{4, 3a} - 30gn_{a^2} \equiv 168 \pmod{210}$
5. $x(1, b) - 3y(1, b) + 47205So_b - 1557T_{b, 2} \equiv 0 \pmod{12138}$

2.2 Pattern: 2

Rewrite (3) as $u^2 + 17v^2 = 21z^5 * 1$ (6)

Write 1 as $1 = \frac{1}{81} (8 + i\sqrt{17})(8 - i\sqrt{17})$ (7)

Following the procedure similar to Pattern-I, the corresponding non-zero distinct integral solutions of (1) are found to be

$$x = x(a, b) = 1240029a^5 - 6200145a^4b - 210804930a^3b^2 + 210804930a^2b^3 + 1791841905ab^4 - 358368381b^5$$

$$y = y(a, b) = 964467a^5 - 17222625a^4b - 163959390a^3b^2 + 585569250a^2b^3 + 1393654815ab^4 - 995467725b^5$$

$$X = X(a, b) = 2342277a^5 - 17911530a^4b - 398187090a^3b^2 + 608992020a^2b^3 + 3384590265ab^4 - 1035286434b^5$$

$$Y = Y(a, b) = 2066715a^5 - 28934010a^4b - 351341550a^3b^2 + 983756340a^2b^3 + 2986403175ab^4 - 1672385778b^5$$

$$z = z(a, b) = 81a^2 + 1377b^2$$

Properties:

1. $z(a, a) - 639gn_{a^2} - 639$ Nasty number
2. $\frac{1}{51254532} [7x(1, 1) - 9y(1, 1)]$ a Nasty number
3. $\frac{1}{7x(a, 1)} - 9y(a, b) + 105402465 T_{4, 6a} \equiv -22320522 \pmod{111602610}$
4. $15X(a, 1) - 17Y(a, 1) - 55801305gn_{2a^4} \equiv -2276693244 \pmod{7588977480}$
5. $15X(1, b) - 17Y(1, b) - 77407570296CP_{b^2, b} - 2656142118So_b + 771196355622 T_{2, b} = 0$

2.3: Pattern: 3

Instead of (5) write 21 as

$21 = \frac{1}{81} (1 + i10\sqrt{17})(1 - i10\sqrt{17})$ (8)

Following the procedure similar to Pattern-I, and performing a few calculations, the corresponding non-zero distinct integral solutions of (1) are found to be

$$x = x(a, b) = 72171a^5 - 5544045a^4b - 12269070a^3b^2 + 190728270a^2b^3 + 104287095ab^4 - 320445801b^5$$

$$y = y(a, b) = -59049a^5 - 5609655a^4b + 10038330a^3b^2 + 188497530a^2b^3 - 85325805ab^4 - 324238059b^5$$

$$X = X(a, b) = 78732a^5 - 11120895a^4b - 13384440a^3b^2 + 380341170a^2b^3 + 113767740ab^4 - 642787731b^5$$

$$Y = Y(a, b) = -52488a^5 - 11186505a^4b + 8922960a^3b^2 + 378110430a^2b^3 - 75845160ab^4 - 646579989b^5$$

$$z = z(a, b) = 81a^2 + 1377b^2$$

Properties:

1. $9x(a, 1) + 11y(a, 1) + 31000725 - T_{4, 6a} \equiv 1129423662 \pmod{37900027260}$
2. $\frac{1}{28877148} [9x(1, 1) - 11y(1, 1)]$ a Nasty number
3. $\frac{1}{27} z(a, a)$ a Nasty number
4. $8X(a, 1) + 12Y(a, 1) + 1339231320CP_{a, a} - 3678424650gn_{a^2} \equiv 416394254 \pmod{1339231320}$

2.4 Pattern: 4

Instead of (7) write 1 as

$1 = \frac{1}{1089} (1 + i8\sqrt{17})(1 - i8\sqrt{17})$ ---- (9)

Following the procedure similar to Pattern-III, the corresponding non-zero distinct integral solutions of (1) are found to be

$$x = x(a, b) = 224139069a^5 - 16810430175a^4b - 38103641730a^3b^2 + 571554625950a^2b^3 + 323880954705 - 971642864115b^5$$

$$y = y(a, b) = -174330387a^5 - 17059473585a^4b + 29636165790a^3b^2 + 580022101890a^2b^3 - 251907409215ab^4 - 986037573213b^5$$

$$X = X(a, b) = 249043410a^5 - 33745382055a^4b - 42337379700a^3b^2 + 1147342989890a^2b^3 + 3598677274 - 1950483082779b^5$$

$$Y = Y(a, b) = -149426046a^5 - 33994425465a^4b + 25402427820a^3b^2 + 1155810465810a^2b^3 - 215920636470ab^4 - 1964877791877b^5$$

$$z = z(a, b) = 1089a^2 + 18513b^2$$

Properties:

1. $7x(a, 1) + 9y(a, 1) - 54241546980T_{3, a^2-1} \equiv 3308740936578 \pmod{9492289572150}$
2. $\frac{1}{363} [z(a, a)]$ a Nasty number
3. $7x(1, b) + 9y(1, b) \equiv 0 \pmod{18080551566}$
4. $6X(a, 1) + 10Y(a, 1) + 7203829673960T_{3, a^2-1} \equiv 7420123880230gn_{a^2} - 23931552535214$

3. Conclusion

In linear transformations (2), the variables X and Y may also be represented by $X=2uv +1$, $Y= 2uv -1$. Applying the procedure similar to that of pattern I-IV choices of integral

solutions to (1) are obtained. To conclude, one may search for other patterns of solutions and their corresponding properties.

References

- [1] Dickson, L.E., History of Theory of numbers, Vol.2, Chelsea publishing company. New York, 1952.
- [2] Gopalan, M.A & Vidhalakshmi.S and Devibala.S, Integral solutions of $49x^2 + 50y^2 = 51z^2$, Acta cincia indica,XXXIIM(2),839-840,2006. I
- [3] Gopalan M.A. and Sangeetha.G, A remarkable observations on $y^2 = 10x^2 + 1$ Impact.J.sci.Tech, 4(1), 103-106,2010.
- [4] Gopalan M.A. Note on the Diophantine equation $x^2 + xy + y^2 = 3z^2$, Acta cincia indica, XXXVIM(3), 265-266,2000.
- [5] Gopalan M.A. and Sangeetha.G, On the ternary Diophantine equation $y^2 = Dx^2 + z^3$. Archimedeas Journal of Mathematics,1(1),7-14,2010.
- [6] Gopalan M.A., and Manju Somanath and Vanitha.N, Ternary Cubic Diophantine equation $x^2 + y^2 = 2z^3$, Advances in Theoretical and Applied Mathematics, I(3),227-231,2006.
- [7] Gopalan M.A., and Manju Somanath and Vanitha.N, Ternary Cubic Diophantine equation $2^{2a-1} \cdot (x^2 + y^2) = z^3$, Acta cincia Indica,XXXIVM(3),1135-1137,2008.
- [8] Gopalan M.A. and Sangeetha.G, Integral Solution of Ternary non homogeneous biquadratic equation $x^4 + x^2 + y^2 - y = z^2 + z$., Accepted in acta ciencia Indica, XXXVII M(4),799-803,2011.
- [9] Mordell, L.J., Diophantine equations, Academic press, London (1969).

Author Profile

P. Jayakumar received the B. Sc , M.Sc degrees in Mathematics from University of Madras in 1980 and 1983 and the M. Phil., Ph. D degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 1988 and 2010.He is now working as Associate Professor of Mathematics, A.V.V.M Sri Pushpam College Poondi, (Autonomous), Thanjavur (District) – 613 503, Tamil Nadu, India.