



who had taken antibiotics or antihelminthic a week earlier; or declined to give consent were excluded.

**Data collection tools:** Qualitative data was collected from 300 respondents at Bungoma County Hospital from June 2010 to February 2011 using patients' questionnaires. The questionnaires were distributed to patients in presence of the medical practitioners working at the CCC. For quantitative data; a single fresh stool sample of (10-20 gms) was collected from each study participants. 1 g of the stool was preserved in 3 ml of 10% formalin and specimens examined by direct saline, iodine wet mount preparation and formol ether sedimentation methods following the standard procedure. Concentration technique and modified Ziehl Neelsen staining techniques were used for the detection of oocyst of several intestinal parasites [11] under light microscope ( $\times 40$  magnification). Parasite species were identified by their different morphological forms: cysts, flagellates, ciliates, larvae and eggs; with their characteristic identifying features.

**Ethical Consideration:** This study was conducted with the approval of the Institutional Research and Ethics Committee (IREC) of Moi University. Informed written consent was obtained from each study participant. Participants were also informed that they were free to withdraw consent any time; and that their stool specimens were examined by qualified persons. Moreover, all personal information of the participants was treated with strict confidentiality. Study participants positive for intestinal parasites were treated free of charge using standard drugs at the Comprehensive Care Clinic (CCC).

**Data analysis:** All data were analyzed using SPSS V.17. Data were summarized using frequency, means and standard deviation. Differences in prevalence were analyzed using Pearson Chi-square test. Significant factors causing observed differences in the prevalence and intensity of the parasites were analyzed using chi-square test. In all analysis, results were considered significant at  $p < 0.05$ .

### 3. Results

Socio-demographic characteristics of the respondents are provided in Table 1. Most of the HIV/AIDS patients were female 138(57.5%). More than half of the HIV/AIDS patients (125, 56.2%) were aged between 36-50 years. Most of the HIV/AIDS patients had secondary level of education 142(59.2%). With regard to occupation, 109(45.4%) HIV/AIDS(+ve) and 26(43.3%) HIV(-ve) were self employed.

**Table 1:** Socio-demographic characteristics of the respondents

Attribute	Characteristics	HIV(+ve)		HIV(-ve) (control)	
		Frequency	%	Frequency	%
Gender	Male	102	42.5	27	45
	Female	138	57.5	33	55
Age	<18	15	6.3	7	11.7
	18-35 years	36	15.0	12	20
	36-50 years	135	56.2	26	43.3
	>50 years	54	22.5	15	25
Level of education	None	13	5.4	6	10
	Primary	65	27.1	17	28.3
	Secondary	142	59.2	23	38.4
	Tertiary	20	8.3	14	23.3
Occupation	Unemployed	78	32.5	21	35
	Self-employed	109	45.4	26	43.3
	Formal	53	22.1	13	21.7

The prevalence of intestinal parasitic infections among the HIV/AIDS patients was 33.4% while among the healthy individual it was 19.3%. Protozoans and helminthes were identified in the stool samples (Table 2). Among the protozoa highest prevalence was reported for *Entamoeba histolytica* (41.2%), followed by *E. coli*, *Giardia lamblia* and *Cryptosporidium* spp. In the helminth groups, *Ascaris lumbricoides*, *Trichura trichura*, *Ancylostoma duodenale* and *Strongyloides stercoralis* were of importance. The prevalence of intestinal protozoa was significantly higher in the HIV/AIDS patients as compared to the non-infected patients ( $p < 0.05$ ). However, there was no significant difference in the prevalence of intestinal helminthes between the two groups ( $p > 0.05$ ).

**Table 2:** Prevalence of various intestinal parasitic infections among the study subjects

Parasite	Group		Chi-square	P-value
	HIV(+ve)	HIV(-ve)		
<i>Entamoeba histolytica</i>	99 (41.2)	7 (11.7)	18.385	<0.001
<i>Entamoeba coli</i>	86 (35.8)	4 (6.7)	19.44	<0.001
<i>Giardia lamblia</i>	80 (33.3)	6 (13.3)	9.262	0.002
<i>Cryptosporidium</i> spp	30 (12.5)	1 (1.7)	6.080	0.014
<i>Ascaris lumbricoides</i>	33 (13.8)	3 (5)	3.480	0.062
<i>Strongyloides stercoralis</i>	16 (6.7)	1 (1.7)	2.245	0.134
<i>Ancylostoma duodenale</i>	24 (10)	3 (5)	1.465	0.226
<i>Trichuris trichura</i>	28 (10.7)	1 (1.7)	4.245	0.017

A total of eight factors (Age, gender, marital status, levels of education, income levels, smoking habits dietary habits and drinking habits) were analyzed and how they influence the prevalence of intestinal parasites in HIV/AIDS patients. Among the analyzed factors, it was established that all the factors except marital status affected the prevalence of intestinal parasitic infections among the HIV/AIDS patients (Table 3). Prevalence of IPI increased significantly ( $P < 0.05$ ) with increasing age of the patients. Significantly ( $p < 0.05$ ) higher proportion of the females than males had IPI. The general trend in decline of the IPI with levels of education and income levels was also significant ( $p < 0.05$ ). It was also noted that smokers had significantly ( $p < 0.05$ ) higher prevalence of intestinal parasitic infections than the non-smokers. Patients who attested to eating more fatty food (meat) recorded higher prevalence of intestinal parasitic infections than vegetarians ( $p < 0.05$ ). People consuming

alcohol had lower prevalence of IPI than those who did not engage in the drinking.

**Table 3:** Factors affecting the prevalence of intestinal parasitic infections among HIV/AIDS patients at Bungoma County Hospital

Attributes/ Characteristic		Frequency	Prevalence of IPI	$\chi^2$	P-value
Age	<18	26	17.8	19.443	<0.001
	18-35	23	36.9		
	36-55	19	39.7		
	> 55	12	51.6		
Gender	Male	63	22.8	21.311	<0.001
	Female	17	44.4		
Marital status	Married	19	34.5	0.811	0.623
	Single	21	35.1		
	Divorced	21	32.3		
	Widowed	19	32.7		
Levels of education	None	31	49.3	25.442	<0.001
	Primary	18	45.4		
	Secondary	13	33.9		
	College	14	24.2		
	University	4	13.2		
Income levels	< 1500	27	51.1	31.222	<0.001
	1501-5000	18	35.4		
	5001-10000	15	32.5		
	10001-	13	24.9		
	> 20001	7	22.4		
Smoking habits	Smoker	34	45.2	22.133	0.001
	Non-	46	21.3		
Dietary habits	Eat fatty	25	44.2	9.442	0.022
	Vegetarian	8	23.2		
	Eat at home	49	34.3		
	Eat in	11	35.4		
Drinking habits	Drinker	33	27.5	19.233	0.001
	Non-	47	39.3		

#### 4. Discussion

Despite increasing interest in the pathologic interactions between infections, few epidemiologic studies have assessed the occurrence of IPI and associated predictive factors in HIV-infected persons. Moreover, general prevalence surveys are more commonly done in children and none, to our knowledge, have examined risk factors for intestinal helminthic infection among HIV-infected adults in an urban SSA setting. In the present study, the overall prevalence of intestinal parasitic infections (IPI) among HIV/AIDS populations was (33.4%). It was higher than in other areas of Africa such as those reported in Senegal (15.3%) [12], Ivory Coast (17.9%) [13] and Congo (19%) [14], and relatively much higher than in developed countries such as: Netherlands (0.9%), Germany (1.1%), and USA (1.4%) [1]. However prevalence of (33.4%) were lower, compared to those reported in Nigeria (49%) [15]. This was probably due to all the species of parasites caused stress to the immune system and stimulated HIV to be more active, coupled with low level of sanitation [16].

Protozoan infections were more prevalent among the HIV/AIDS patients. Hayashi [17] reported *E. histolytica*'s positive rates of 35.1% in Machakos, 34.1% in Naivasha, almost similar to that reported in this study, but higher than 28.8% in Kitui, 31.4% in Nandi Hills and 27.9% in Taveta. Also reported *E. coli* positive rates of 44.2% in Naivasha, 46.4% in Kitui, 63.3% in Taveta; which were higher than those reported in our study. It is possible that human body already weakened by HIV/AIDS infection becomes an important site for parasitic infections. The resulting defects in cellular and humoral defense mechanism predispose the body to spectrum parasitic pathogens [18]. The study also determined lower helminth infections compared to protozoans. Low prevalence of helminthic infections has been reported in other tropical areas such as Thailand and especially among the HIV/AIDS individuals at different immune status [19]. Odhiambo reported high prevalence of amoebiasis in Kisumu County and observed that out of 1432 stool specimens 338 (about 26.3%) were positive for intestinal parasites [20]. The author attributes the widespread distribution of intestinal protozoa among HIV(+ve), especially *E. histolytica* and *E. coli*, in the County due to poor sanitation.

In this study, eight pre-disposing factors were used to determine the factors likely to influence the prevalence of the intestinal parasitic infections among the HIV/AIDS patients. First, it was established that prevalence of IPI increased with age. According to the results, HIV/AIDS patients aged 18 to 35 years had significantly lower prevalence of intestinal parasitic infections than those aged over 55 years. This observation is similar with that of [21]. The overall high infection rate with intestinal parasites recorded in the elderly could be due to reduced immunity during senescence [22]. Meanwhile significantly higher proportion of females had intestinal parasitic infections than their male counterparts. The higher prevalence of IPI among the females compared to males in these studies can be attributed to the fact that females in the study area engage in water and food preparation for the family, thus leaving them more exposed to infective agents of IPI than men. Alternatively it is also possible that more females visited the hospital not only for medical treatment but also for other services such as antenatal services and family planning in the study area. This is contrary to [23] observation that reported opposing trend and attributed it to the fact that males have fewer restrictions than females. The findings that parasite infections declined with increased level education of the study participants indicated the overall improvement of hygienic conditions and sanitation with knowledge of self deworming. It has been shown that such a relation between increase in educational level and lower prevalence of intestinal parasite infection [24]. Higher income levels was also found to result in reduced prevalence of IPI mainly because, people with high income have ability to afford drugs that will reduce the prevalence of IPI than those without any disposable income. Smoking and poor dietary habits were all found to increase the prevalence of IPI among the HIV/AIDS patients. Smoking, poor dietary habits and excess drinking have been associated with reduced immune response and increased pathogens in the body [25] and can presumably explain the high prevalence of these

pathogens among the patients who smoked, non-vegetarians or consumed less alcohol.

## 5. Future Prospects

Numerous studies have shown that multiple factors can each influence the proportion of parasites infections in the current study, HIV status of the individual has been highlighted yet as another factor that may affect the prevalence of the IPI. In management of the IPI, this study support determination of HIV/AIDS status of the patients before commencement of treatment for IPI.

## References

- [1] WHO (2010). Prevalence of preventable diseases in the world. Report of the World health Organization, Geneva. Switzerland.
- [2] Chacon-Cruz, E. (2009). Intestinal Protozoal Diseases among HIV/AIDS patients. *eMedicine J.* **3(5)**: 1–11.
- [3] Wiwanitkit, V. (2005). Intestinal parasitic infections in HIV-infected patients with different immunity status. *BMC Gastroenterol.* **1**:1–3.
- [4] Jemaneh, L. (1998). Comparative prevalence of some common intestinal helminthes infections in different altitudinal regions in Kenya. *Kenya Med. J.* **36**:1–8.
- [5] Adamu, H., Endeshaw, T., Teku, T. et al. (2009). Prevalence of intestinal parasites in paediatric diarrhoeal and non-diarrhoeal patients in Sub-Saharan Africa, with special emphasis on opportunistic parasitic infections. *Ethiop. J. Health. Dev.* **20(1)**: 39–46.
- [6] Ikpeba, A.C. and Ojololo, L.P. (2005). The intrigue between HIV/AIDS infection and prevalence of intestinal parasitic infections. *Afr. J. HIV Manag.* **103**: 145–148.
- [7] Joshi, M., Chowdhary, A.S., Dalal, P.J. et al. (2002). Parasitic diarrhea in patients with AIDS. *Nat. Med. J. India.* **15**: 72–74.
- [8] MacGowan S.J. (2007). Advances in mucosal immunology. *Gastro. Clin. North Am.* **26**:145–173.
- [9] Cimerman, S., Cimerman, B. & Salomo-Lewi, D. (1999). Prevalence of Intestinal Parasitic Infections (IPI) in patients with Acquired Immunodeficiency Syndrome in Brazil. *Int. J. Infect. Dis.* **3**: 203-206.
- [10] Mugenda, O.M & Mugenda, A.B.(1999). *Research Methods. Quantitative and Qualitative Approaches.* Nairobi: Acts Press
- [11] Cheesbrough, M. (1998). County laboratory practice in tropical countries. Tropical health technology, UK. Part I: 192–198.
- [12] Niang, M.E. (2008). Prevalence and intensity of intestinal parasites in Northern Senegal city of Dakar. *Afr. J. Biotech.* **7**: 4141–4144.
- [13] Mamadou, K.P. (2010). Prevalence and risk factors associated with intestinal parasites in Ivory Coast. *J. Parasite. Trop.* **12**: 133–141.
- [14] Nzenga, M.P., Joskey, M.P., Mbola, M.P. et al. (2009). Prevalence of intestinal parasitic infections in Katanga region of Zaire. *Afr. J. Paras. Infect.* **10**: 11–19.
- [15] Awolaju, B.A. & Morenikeji, O.A. (2009). Prevalence and intensity of intestinal parasites in five communities in south-west Nigeria. *Afr. J. Biotech.* **8**: 4542–4546.
- [16] Ortega, .Y & Adam, R. (1997). Giardia: an overview and update. *Clin. Infect. Dis.* **25(3)**: 545-549.
- [17] Hayashi, K.P. (2007). Prevalence of opportunistic parasitic infections among the HIV/AIDS positive individuals in selected towns in Kenya. *J. Trop. Dis.* **23**: 133–145.
- [18] Legesse, M. and Erko, B. (2003). Prevalence of intestinal parasites among school children in a rural area close to south east of Lake Langano, Ethiopia. *Ethiop. J. Health Develop.* **18**: 116–120.
- [19] Stringer, EM, Sinkala, M, Kumwenda, R. et al. (2004). Personal risk perception, HIV knowledge and risk avoidance behavior, and their relationships to actual HIV serostatus in Thailand obstetric population. *J. Acquir. Immune Def. Syndr.* **35**: 60–66.
- [20] Odhiambo, K.O. (2001). Prevalence of opportunistic infections among the HIV positive individuals and the levels of immunodeficiency. *J. Opport. Infect.* **23**: 133-145.
- [21] Sarfati, C., Bourgeois, A., Menotti, J. et al. (2006). Prevalence of Intestinal Parasites including Microsporidia in Human Immunodeficiency Virus–Infected Adults in Cameroon: A Cross-Sectional Study. *AJTMH.* **74(1)**: 162–164.
- [22] George-Morris, R.T. (2004). Immunodereulatory pathology during senescence period in a cohort study. *Nature, London.* **133**: 2423-2432.
- [23] Awogun I.A. (2011). The prevalence of intestinal parasitic infections among HIV/AIDS infected children living in Igboora, Oyo State, Nigeria. *West Afr. J. Med.* **4(1)**: 16-21.
- [24] Bern, C., Hernandez, B., Lopez, M.B. et al. (2000). The Contrasting epidemiology of *Cyclospora* and *Cryptosporidium* among outpatients in Guatemala. *AJTMH.* **63(5, 6)**: 231–235.
- [25] Foudraine, N.A., Weverling, G.J. & Van Gool, T. (1998). Improvement of chronic diarrhoea in patients with advanced HIV-1 infection during potent antiretroviral therapy. *AIDS.* **12**: 35–41.

## Author Profile



**Dr. Fredrick Wabwile Wanyama** has a Masters of Science in Parasitology (University of Eldoret); Bachelor of Science (Moi University) and a higher National Diploma in Clinical Medicine (Paediatrics KMTC Nairobi) with a Diploma in Clinical Medicine and Surgery (Port Reitz School of Clinical Medicine). He currently works at the University of Eldoret Health Services PhD in Aquatic Ecology and Ecotoxicology from the University of Amsterdam, The Netherlands. He has published several publications in top ranking Toxicology Journals. He has diverse interested in Environmental Biology, Environmental Health, Fisheries Science and Ecotoxicology.



**Mwongula W. Albert** is a graduate with MSc in Microbiology and BSc in Microbiology from Moi University. He is a researcher and scientist in the field of Microbiology. Currently he is a lecturer at University of Eldoret at the Department of Biological Sciences and Mount Kenya University (Kakamega Campus) at the Department of Health Sciences.



**Dr. Elijah Oyoo-Okoth** has a PhD in Aquatic Ecology and Ecotoxicology from the University of Amsterdam, The Netherlands. He has published several publications in top ranking Toxicology Journals. He has diverse interested in Environmental Biology, Environmental Health, Fisheries Science and Ecotoxicology.