

- www.uptodate.com/.../minor-head-trauma-in-infants-and-children. [Retrieved on Aug 13, 2014].
- [168] Tabish SA, Lone N. A, Wani M. Afzal, Bhat A S (2007). Clinical Management of Patients with Minor Head Injuries. International Journal of Health Sciences, Qassim University, 1(1)
- [169] Finkelstein EA, Corso PS, Miller TR (2006), Associates. Incidence and economic burden of injuries in the United States. New York, NY: Oxford University Press.
- [170] Todd W. Vitaz, Laura McIlvoy, George H. Raque, David Spain, and Christopher B (2001). Development and Implementation of a Clinical Pathway for Severe Traumatic Brain Injury. J Trauma 51:369–375
- [171] Gordon D. Murray, Isabella Butcher, Gillian S. Mchugh, Juan Lu, et al (2006). Multivariable Prognostic Analysis in Traumatic Brain Injury: Results from the IMPACT Study. Journal of Neurotrauma. 24, (2), 2007:329–337. DOI: 10.1089/neu.2006.0035
- [172] [Guideline] Society of Critical Care Medicine (2012). Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents. Pediatr Crit Care Med. 13(1 Suppl):S1-82.
- [173] Khanna S, Davis D, Peterson B, et al (2000). Use of hypertonic saline in the treatment of severe refractory posttraumatic intracranial hypertension in pediatric traumatic brain injury. Crit Care Med. 28(4):1144-51.
- [174] Faul M, Wald MM, Rutland-Brown W, Sullivent EE, Sattin RW (2007). Using a cost-benefit analysis to estimate outcomes of a clinical treatment guideline: testing the Brain Trauma Foundation guidelines for the treatment of severe traumatic brain injury. J Trauma. 63(6):1271-8.
- [175] Andrew I R Maas, Gordon D Murray, Bob Roozenbeek et al (2013). Advancing care for traumatic brain injury: findings from the IMPACT studies and perspectives on future research. The Lancet Neurology, 12(12):1200 - 1210
- [176] Shruti V. Kabadi and Alan I. Faden. (2014). Neuroprotective Strategies for Traumatic Brain Injury: Improving Clinical Translation. Int. J. Mol. Sci. 15(1), 1216-1236
- [177] Maas AI, Stocchetti N, Bullock R (2008). Moderate and severe traumatic brain injury in adults. Lancet Neurol. 7(8):728-41. doi: 10.1016/S1474-4422(08)70164-9.
- [178] John Bruns, Jr., and W. Allen Hauser (2003). The Epidemiology of Traumatic Brain Injury: A Review. Epilepsia, 44(Suppl. 10):2–10
- [179] http://www.cdc.gov/FieldTriage/pdf/EMS_Guide-a.pdf. [Accessed 27 August 2014].
- [180] Faden A.I., Knoblach S.M., Cernak I., Fan L., Vink R., Araldi G.L., Fricke S.T., Roth B.L., Kozikowski A.P. (2003). Novel diketopiperazine enhances motor and cognitive recovery after traumatic brain injury in rats and shows neuroprotection *in vitro* and *in vivo*. J. Cereb. Blood Flow Metab. 2003;23:342–354.
- [181] Faden A.I., Movsesyan V.A., Knoblach S.M., Ahmed F., Cernak I. (2005). Neuroprotective effects of novel small peptides *in vitro* and after brain injury. Neuropharmacology. 49:410–424.
- [182] Guan J., Mathai S., Harris P., Wen J.Y., Zhang R., Brimble M., Gluckman P. (2007). Peripheral administration of a novel diketopiperazine, NNZ 2591, prevents brain injury and improves somatosensory-motor function following hypoxia-ischemia in adult rats. Neuropharmacology. 53:749–762.
- [183] Simard J.M., Woo S.K., Bhatta S., Gerzanich V. Drugs acting on SUR1 to treat CNS ischemia and trauma. Curr. Opin. Pharmacol. 2008;8:42–49.
- [184] Simard J.M., Kilbourne M., Tsymbalyuk O., Tosun C., Caridi J., Ivanova S., Keledjian K., Bochicchio G., Gerzanich V. Key role of sulfonyleurea receptor 1 in progressive secondary hemorrhage after brain contusion. J. Neurotrauma. 2009;26:2257–2267.
- [185] Shu-Hui Dai, Tao Chen, Yu-Hai Wang, Jie Zhu, Peng Luo, Wei Rao, Yue-Fan Yang, Zhou Fei, and Xiao-Fan Jiang. (2014). Sirt3 Protects Cortical Neurons against Oxidative Stress via Regulating Mitochondrial Ca²⁺ and Mitochondrial Biogenesis. Int J Mol Sci. 15(8): 14591–14609.
- [186] Nilgün Şenol and Mustafa Nazıroğlu (2014). Melatonin reduces traumatic brain injury-induced oxidative stress in the cerebral cortex and blood of rats. Neural Regen Res. 9(11): 1112–1116.
- [187] Wible E.F., Laskowitz D.T. (2010). Statins in traumatic brain injury. Neurotherapeutics. 7:62–73.
- [188] Chen G., Zhang S., Shi J., Ai J., Qi M., Hang C. (2009). Simvastatin reduces secondary brain injury caused by cortical contusion in rats: Possible involvement of TLR4/NF-kappaB pathway. Exp. Neurol. 216:398–406.
- [189] Tapia-Perez J., Sanchez-Aguilar M., Torres-Corzo J.G., Gordillo-Moscoso A., Martinez-Perez P., Madeville P., de la Cruz-Mendoza E., Chalita-Williams J. (2008). Effect of rosuvastatin on amnesia and disorientation after traumatic brain injury (NCT003229758) J. Neurotrauma. 25:1011–1017.
- [190] Okonkwo D.O., Büki A., Siman R., Povlishock J.T. (1999). Cyclosporin A limits calcium-induced axonal damage following traumatic brain injury. Neuroreport. 10:353–358. [PubMed]
- [191] Margulies S., Hicks R. (2009). Combination therapies for traumatic brain injury: Prospective considerations. J. Neurotrauma. 26:925–939.
- [192] Donkin J.J., Nimmo A.J., Cernak I., Blumbergs P.C., Vink R. (2009). Substance P is associated with the development of brain edema and functional deficits after traumatic brain injury. J. Cereb. Blood Flow Metab. 29:1388–1398.
- [193] Nimmo A.J., Cernak I., Heath D.L., Hu X., Bennett C.J., Vink R. (2004). Neurogenic inflammation is associated with development of edema and functional deficits following traumatic brain injury in rats. Neuropeptides. 38:40–47.
- [194] Arendt T. (2003). Synaptic plasticity and cell cycle activation in neurons are alternative effector pathways: The ‘Dr. Jekyll and Mr. Hyde concept’ of Alzheimer’s disease ordisease or the yin and yang of neuroplasticity. Prog. Neurobiol. 71:83–248.

- [195] Di Giovanni S., Movsesyan V., Ahmed F., Cernak I., Schinelli S., Stoica B., Faden A.I. (2005). Cell cycle inhibition provides neuroprotection and reduces glial proliferation and scar formation after traumatic brain injury. *Proc. Natl. Acad. Sci. USA.* 102:8333–8338.
- [196] Hilton G.D., Stoica B.A., Byrnes K.R., Faden A.I. (2008). Roscovitine reduces neuronal loss, glial activation, and neurologic deficits after brain trauma. *J. Cereb. Blood Flow Metab.* 28:1845–1859.
- [197] Bettayeb K., Oumata N., Echalié A., Ferandin Y., Endicott J.A., Galons H., Meijer L. (2008). CR8, a potent and selective, roscovitine-derived inhibitor of cyclin-dependent kinases. *Oncogene.* 27:5797–5807.
- [198] Kabadi S.V., Stoica B.A., Hanscom M., Loane D.J., Kharebava G., Murray M.G., II, Cabatbat R.M., Faden A.I. (2012). CR8, a selective and potent CDK inhibitor, provides neuroprotection in experimental traumatic brain injury. *Neurotherapeutics.* 9:405–421.
- [199] Loane D.J., Stoica B.A., Pajoohesh-Ganji A., Byrnes K.R., Faden A.I. (2009). Activation of metabotropic glutamate receptor 5 modulates microglial reactivity and neurotoxicity by inhibiting NADPH oxidase. *J. Biol. Chem.* 284:15629–15639.
- [200] Loane D.J., Stoica B.A., Byrnes K.R., Jeong W., Faden A.I. (2013). Activation of mGluR5 and inhibition of NADPH oxidase improves functional recovery after traumatic brain injury. *J. Neurotrauma.* 30:403–412.
- [201] Byrnes K.R., Loane D.J., Stoica B.A., Zhang J., Faden A.I. (2012). Delayed mGluR5 activation limits neuroinflammation and neurodegeneration after traumatic brain injury. *J. Neuroinflamm.* 9:43.
- [202] Clark R.S., Kochanek P.M., Watkins S.C., Chen M., Dixon C.E., Seidberg N.A., Melick J., Loeffert J.E., Nathaniel P.D., Jin K.L., et al. (2000). Caspase-3 mediated neuronal death after traumatic brain injury in rats. *J. Neurochem.* 2000;74:740–753.
- [203] Zhu C., Wang X., Huang Z., Qiu L., Xu F., Vahsen N., Nilsson M., Eriksson P.S., Hagberg H., Culmsee C., et al. (2007). Apoptosis-inducing factor is a major contributor to neuronal loss induced by neonatal cerebral hypoxia-ischemia. *Cell Death Differ.* 2007;14:775–784.
- [204] Piao C.S., Loane D.J., Stoica B.A., Li S., Hanscom M., Cabatbat R., Blomgren K., Faden A.I. (2012). Combined inhibition of cell death induced by apoptosis inducing factor and caspases provides additive neuroprotection in experimental traumatic brain injury. *Neurobiol. Dis.* 46:745–758.
- [205] Turturici G., Sconzo G., Geraci F. (2011). Hsp70 and its molecular role in nervous system diseases. *Biochem. Res. Int.* 2011:618127.
- [206] Sabirzhanov B., Stoica B.A., Hanscom M., Piao C.S., Faden A.I. (2012). Over-expression of HSP70 attenuates caspase-dependent and caspase-independent pathways and inhibits neuronal apoptosis. *J. Neurochem.* 123:542–554.
- [207] Zhao Z., Faden A.I., Loane D.J., Lipinski M.M., Sabirzhanov B., Stoica B.A. (2013). Neuroprotective effects of geranylgeranylacetone in experimental traumatic brain injury. *J. Cereb. Blood Flow Metab.* 33:1897–1908.
- [208] Lai Y., Hickey R.W., Chen Y., Bayir H., Sullivan M.L., Chu C.T., Kochanek P.M., Dixon C.E., Jenkins L.W., Graham S.H., et al. (2008). Autophagy is increased after traumatic brain injury in mice and is partially inhibited by the antioxidant gamma-glutamylcysteinyl ethyl ester. *J. Cereb. Blood Flow Metab.* 28:540–550.
- [209] Erlich S., Alexandrovich A., Shohami E., Pinkas-Kramarski R. (2007). Rapamycin is a neuroprotective treatment for traumatic brain injury. *Neurobiol. Dis.* 26:86–93.
- [210] Griesbach G.S., Gomez-Pinilla F., Hovda D.A. (2004). The upregulation of plasticity-related proteins following TBI is disrupted with acute voluntary exercise. *Brain Res.* 1016:154–162.
- [211] Griesbach G.S., Gomez-Pinilla F., Hovda D.A. (2004). The upregulation of plasticity-related proteins following TBI is disrupted with acute voluntary exercise. *Brain Res.* 2004;1016:154–162.
- [212] Piao C.S., Stoica B.A., Wu J., Sabirzhanov B., Zhao Z., Cabatbat R., Loane D.J., Faden A.I. (2013). Late exercise reduces neuroinflammation and cognitive dysfunction after traumatic brain injury. *Neurobiol. Dis.* 2013;54:252–263.
- [213] Traumatic Brain Injury: Hope Through Research. National Institute of Neurological Disorders and Stroke. National Institutes of Health. Bethesda, MD 20892. http://www.ninds.nih.gov/disorders/tbi/detail_tbi.htm. [Retrieved on 24 Oct 2014]
- [214] Rutland-Brown W, Langlois JA, Thomas KE, Xi YL (2006). Incidence of traumatic brain injury in the United States, 2003. *J Head Trauma Rehabil* 21:544.
- [215] Tagliaferri F, Compagnone C, Korsic M, et al (2006). A systematic review of brain injury epidemiology in Europe. *Acta Neurochir (Wien)* 148:255.
- [216] Brain Trauma Foundation, American Association of Neurological Surgeons, Congress of Neurological Surgeons, et al (2007). Guidelines for the management of severe traumatic brain injury. Introduction. *J Neurotrauma* 24 Suppl 1:S1.
- [217] Patel HC, Bouamra O, Woodford M, et al (2005). Trends in head injury outcome from 1989 to 2003 and the effect of neurosurgical care: an observational study. *Lancet* 366:1538.
- [218] Davis DP, Peay J, Sise MJ, et al (2010). Prehospital airway and ventilation management: a trauma score and injury severity score-based analysis. *J Trauma* 2010; 69:294.