

The Relationship between Depth and Diameter of Human Acetabulum in Dry Hip Bone of Indian Population

Rajkumar¹, Anant Sachan², Kalu Ram Meena³

¹Lecturer in Govt. Medical College, Jalaun, U.P., India

²Lecturer in S.M.M.H. Medical College, Saharanpur, U.P., India

³Lecturer in Arba Minch College of Medicine & Health Science Arba Minch Ethiopia

Abstract: The acetabulum (Latin Acetabulum-Shallow Vinegar Cup) is an approximately hemispherical cavity central on the lateral aspect of the innominate bone, facing anteroinferiorly and surrounded by an irregular margin deficient inferiorly at the acetabular notch. The acetabular fossa forms the central floor and is rough and non-articular. The articular lunate surface is widest above (the 'dome'), where weight is transmitted to the femur. All three innominate elements contribute to the acetabulum, but unequally. The pubis forms the anterosuperior fifth of the articular surface, the ischium forms the floor of the fossa and rather more than the posteroinferior two-fifths of the articular surface, and the ilium forms the remainder. In the present study consists of 154 dry human hip bone of unknown sex from Teerthankar Mahaveer Medical College & Research Centre, Moradabad. To determine the depth and width of human acetabulum in dry hip bone of Indian population, using the Stainless steel sliding caliper and Slide Scale, The mean \pm S.D value of depth and diameter of acetabulum were 27.14 ± 3.50 mm, 47.57 ± 3.99 mm. The maximum and minimum measurements of acetabulum depth were 38.0 mm; 13.0 mm and maximum and minimum measurements of acetabulum diameter were 56.0 mm, 39.0 mm respectively. Morphological study on adult hip bone is useful for anatomists, anthropologists, experts in Forensic medicine, and orthopedics for performing surgical procedures in this area.

Keywords: Depth of acetabulum, Diameter of acetabulum, dry hip bone, Stainless steel sliding caliper and Slide Scale

1. Introduction

The acetabulum is a deep cup-shaped cavity on the lateral aspect of the hip-bone about its center, and is directed laterally, downwards and somewhat forwards. It is surrounded by an irregular projecting margin which is deficient inferiorly; this gap is termed the acetabular notch⁹.

The floor of the cavity is roughened and non-articular and is termed the acetabular fossa. The sides of the cup present a horseshoe-shaped articular surface which is widest superiorly; in this situation the weight of the trunk is transmitted to the femur in the erect attitude. In the recent state this strip is covered with articular cartilage and provides the surface on which the head of the femur moves within the hip-joint. All three elements of the hip-bone contribute to the formation of the acetabulum in human, but not in equal proportions⁹.

The pubis forms the upper and anterior fifth of the articular surface; the ischium forms the floor of the acetabular fossa and rather more than the lower and posterior two-fifths of the articular surface and the ilium forms the remainder of the articular surface⁹.

2. Material & Methods

The material for the present study consists of 154 dry human hip bone of unknown sex. All these bones were fully ossified bones and free from any pathological or congenital defect. These bones were obtained from various sources:-

1. Teerthankar Mahaveer Medical college & Research centre, Moradabad (U.P).
2. Chhatrapati Shahu ji Maharaj Medical College, Lucknow.
3. Shri Ram Murti Smarak Institute of Medical Sciences, Bareilly

Sample size: In the present study the sample no. are 154. It's estimation based on precision. The Precision consists of significance levels and the allowable error. In this study 5% significance level and various levels of significance are considered. Correlation between morphometrical parameters were investigated using z-test, $p < 0.0001$ was significant level.

From the study the Standard Deviation (S.D) for various parameters are as follows-

S.D. of Acetabulum-

Diameter- 0.399

Depth- 0.350

Measurements were taken using:-

1. Stainless steel sliding caliper
2. Slide Scale



Figure 1: Showing Used Instruments

The measurements were done on intact parts of normal bones. Bones showing wear and tear, fracture or any

pathology were not considered. Each linear recording was taken to the nearest millimeter.

Following parameters were used for measurement:

Acetabulum- Diameter and depth

Measurements for all the above indices were taken as per the norms described in anthropometry. The data obtained for all parameters were analyzed statistically to find range, z-test, mean and standard deviation (S.D.). Each variable was measured 2 times at 2 different sessions by the same observer and the mean value of the 2 measurements was calculated for each variable of each bone for the measurements of these variables a scale.

Depth of Acetabulum

A thin slide scale was placed across the diameter of the acetabulum. Depth of the acetabulum was measured in mm using vernier scale from the center of the acetabulum to the slide scale (fig.no-2). Measurements were recorded accurately to 1/10 of a millimeter by this scale.



Figure 2: Showing measurement of depth of acetabulum

Diameter of acetabulum

Maximum transverse diameter of the acetabulum was measured using vernier calipers (fig.no-3).



Figure 3: Showing measurement of diameter of the acetabulum

3. Result

Summarizes the means and standard deviations of depth and diameter of acetabulum of the hip bone. The result shown in below.

No. of steps	Depth	Diameter
Mean	27.14	47.57
Standard deviation	3.50	3.99
z-test	-96.2282	-147.9519
P value	< 0.0001	< 0.0001
Correlation(r)	0.48	

The mean ±S.D value of depth and diameter of acetabulum were 27.14 ± 3.50mm, 47.57 ± 3.99mm. The maximum and minimum measurements of acetabulum depth were 38.0 mm; 13.0 mm and maximum and minimum measurements of acetabulum diameter were 56.0 mm, 39.0 mm respectively. Distribution of anatomical parameters of hip bone and unknown sex related differences within various ethnic parameters are presented in Table. Statistical software namely SPSS 10.0 was used for analysis of data. Microsoft word and Excel have been used to generate graphs, tables, etc.

4. Discussion

In the present study effort has been made to find the dried human hip bone with the available data in relation to the various parameters. Total of about 154 hip bone have been studied. The parameters have been selected from the literature available in the well-known Anatomy textbooks and published articles. In these articles the study was on identification of validity of the parameters in the hip bone of known sex. In the following section the results of the present study when compared with that of previous ones show the following results.

Table 1: Comparison of present study with earlier studies

Acetabulum	Mean (Depth)	Mean (Diamention)
Present study	27.14mm	47.57mm
Funda Tastekin Aksu	29.49mm	54.29mm
Croft et al	14.4mm	Not available
Lau et al	11.8mm	Not available
Jeremic dejan et al	11.9mm	±2.8
Jeremic et al	11.9mm	±2.8

The mean(Funda Tastekin Aksu,2006) depth of acetabulum of the dried hip bone was 29.49mm. In the present study it is 27.14mm and is low when compared with the previous study. Whereas the range in the present study lies within the range observed in the previous study (refer Table No.1). The mean (Funda Tastekin Aksu, 2006) diameter of acetabulum of the dried hip bone was 54.29mm. In the present study it is 47.57mm and is low when compared with the previous study. Whereas the range in the present study lies within the range observed in the previous study (refer Table No.2). The whole of the study relates with all the relevant past studies except funda tuskien aksu whose study is greater than the observed in the present study. The acetabular depth is less than 9 mm so it may be considered as the acetabular dysplasia⁴.

Table 2: Comparison of present study with earlier studies

Acetabulum	Mean (Depth)	Mean (Diamention)
Present study	27.14mm	47.57mm
Funda Tastekin Aksu	29.49mm	54.29mm
Croft et al	14.4mm	Not available
Lau et al	11.8mm	Not available
Jeremic dejan et al	11.9mm	2.8mm

5. Conclusion

Most of the Previous Studies have been done on those parameters which determine the gender variations in the hip bones. After comparing the findings of the present study with the previous data available, the observations are similar while others are varying due to the formulas used for these calculations. Morphological study on adult hip bone is useful for anatomists, anthropologists, experts in Forensic medicine, and orthopedics for performing surgical procedures in this area. A radiological study may be added for further accuracy. The purpose of this work is to contribute to the scientific literature, providing anatomical data on the similarities and variations. The depth of acetabulum correlates with acetabular diameter so this information may be helpful during hip arthroplasty, treatment of joint fracture and in diagnosing congenital hip dysplasia.

References

- [1] Croft, P., Cooper, C., Wickham, C., and D. Coggon (1991). Osteoarthritis of the hip and acetabular dysplasia. *Ann Rheum Dis*, 50, 308-310.
- [2] Funda Tastekin Aksu, Nazli Gulriz Ceri, Candan Arman, Suleyman Tetik, Morphology and Morphometry of the Acetabulum, *Cilt 20, Sayi 3, (Eylul) 2006, S: 143 – 148.*
- [3] Lau, E.M.C., Lin, F., Lam, D., Silman, A., and P. Croft (1995). Hip osteoarthritis and dysplasia in Chinese men. *Ann Rheum Dis*, 54L, 965-969.
- [4] Murray, R.O. (1965). The etiology of primary osteoarthritis of the hip. *Br J Radiol*, 38, 810-824.
- [5] Dejan Jeremic, Ivana Zivanovic Macuzic, Maja Vulovic (2011). Sex dimorphism of postural parameters of the human Acetabulum among asymptomatic Serbian population. *Vojnosanit Pregl* 2011;68 (11):935-939.
- [6] D. Jeremic, B. Jonovic, Ivana Zivanovic- Macuzic, Gordana Dordevic, Maja Sazdanovic, P. Sazdanovic, Maja Volovic and J. tosevski sex dimorphism of postural parameters of human Acetabulum, *Arch. Biol. Sci. Belgrade*, 63(1), 137-143, 2011.
- [7] Kreyszig E, *Introductory Mathematical Statistics*, John Wiley, 1970, page 206.
- [8] Richard S. Snell, "Clinical anatomy by regions" ed. 8th 555-559.
- [9] Willams PL., Bannister LH, Martin MM Collins, P., Dyson, M. and Dussek, JE. MWJ, *Gray's Anatomy* (39ed.) Churchill Livingstone, London, pp-1421-1427.