International Journal of Science and Research (IJSR)
ISSN (Online): 2319-7064
Impact Factor (2012): 3.358

On Certain Inequalities Pertaining to I-Function

V.G.Gupta[1], Nawal Kishor Jangid[2]

[1]Department of Mathematics, University of Rajasthan, Jaipur - 302055, Rajasthan, India

[2]Department of Mathematics, Swami Keshvanand Institute of Technology, Management and Gramothan, Ramnagaria, Jaipur-302025, Rajasthan, India

Abstract: ***In this paper, we establish four inequalities pertaining to I-function . On account of general nature of the results established here, a number of known and new results follow as their particular cases on suitable specifications of the parameters involved there in. To illustrate, we have recorded some particular cases of our main results.***

Mathematics Subject classification: 33C60, 26D20

Keywords: I-function, Gamma function, Inequalities

1. Introduction

The I-function occurring in the present paper was studied by Saxena[3] and defined as

$$I^{M,N}_{P_i,Q_i:r}[z] = I^{M,N}_{P_i,Q_i:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i}\end{matrix}\right.\right]$$

$$= \frac{1}{2\pi\omega}\int_L \varphi(\xi)\, z^{\xi}\, d\xi \qquad (1)$$

where

$$\varphi(\xi) = \frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}} \qquad (2)$$

and $P_i(i=1,2,...,r)$, $Q_i(i=1,2,...,r)$, M, N are integers satisfying $0\le N\le P_i$, $0\le M\le Q_i$ $(i=1,2,...,r)$; r is finite, $\alpha_j,\beta_j,\alpha_{ji},\beta_{ji}$ are real and positive and a_j,b_j,a_{ji},b_{ji} are complex numbers such that $a_j(b_h+v)\ne\beta_h(a_j-1-k)$, for v,k = 1,2,…; h = 1,2,…,M; i =1,2,…,r.L is the contour running from $\sigma-i\infty$ to $\sigma+i\infty$ (σ is real) in the complex ξ-plane such that the points

$$\xi=(a_j-1-v)/a_{j,}\qquad j=1,2,...,N;v=0,1,2,...$$

$$\xi=(b_j+v)/\beta_j,\qquad j=1,2,...,M;v=0,1,2,...$$

lie to the left hand and right hand sides of L, respectively.
For the present study we shall use the following result due to [2]

$$\frac{\prod_{i=1}^{n}\Gamma(1+\alpha_i)}{\Gamma\left(\beta+\sum_{i=1}^{n}\alpha_i\right)} \le \frac{\prod_{i=1}^{n}\Gamma(1+\alpha_i x)}{\Gamma\left(\beta+x\sum_{i=1}^{n}\alpha_i\right)} \le \frac{1}{\Gamma(\beta)} \qquad (3)$$

where $x\in[0,1]$, $\beta\ge1$, $\alpha_i>0$, $n\in N$

2. Main Results

<u>Inequality 1</u> By putting $\alpha_i=\mu_i\xi-a_i$, $\beta=\upsilon+1+\mu\xi$ we get

Licensed Under Creative Commons Attribution CC BY

$$\frac{\prod_{i=1}^{n}\Gamma(1+\mu_i\xi-a_i)}{\Gamma\left(\upsilon+1+\mu\xi+\sum_{i=1}^{n}(\mu_i\xi-a_i)\right)} \le \frac{\prod_{i=1}^{n}\Gamma(1+(\mu_i\xi-a_i)x)}{\Gamma\left(\upsilon+1+\mu\xi+x\sum_{i=1}^{n}(\mu_i\xi-a_i)\right)} \le \frac{1}{\Gamma(\upsilon+1+\mu\xi)}$$

multipying all three terms with

$$\frac{1}{2\pi\omega}\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}$$ and then integrating along the contour L

$$\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{\prod_{i=1}^{n}\Gamma(1+\mu_i\xi-a_i)}{\Gamma\left(\upsilon+1+\mu\xi+\sum_{i=1}^{n}(\mu_i\xi-a_i)\right)}d\xi$$

$$\le\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{\prod_{i=1}^{n}\Gamma(1+(\mu_i\xi-a_i)x)}{\Gamma\left(\upsilon+1+\mu\xi+x\sum_{i=1}^{n}(\mu_i\xi-a_i)\right)}d\xi$$

$$\le\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{1}{\Gamma(\upsilon+1+\mu\xi)}d\xi$$

=

$$I^{M,N+n}_{P_i+n,Q_i+1:r}\left[z\left|\begin{matrix}(a_i,\mu_i)_{1,n};(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(-\nu+\sum_{i=1}^{n}a_i,\mu+\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\le I^{M,N+n}_{P_i+n,Q_i+1:r}\left[z\left|\begin{matrix}(a_ix,x\mu_i)_{1,n};(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(-\nu+x\sum_{i=1}^{n}a_i,\mu+x\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\le I^{M,N}_{P_i,Q_i+1:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(-\nu,\mu)\end{matrix}\right.\right]$$ (4) provided that

$x\in[0,1]$

Licensed Under Creative Commons Attribution CC BY

$$\mathrm{Re}\left[\nu+\min_{1\le j\le M}\mu\left(\frac{b_j}{\beta_j}\right)\right]>1,\ \mathrm{Re}\left[a_i+\min_{1\le j\le M}\mu_i\left(\frac{b_j}{\beta_j}\right)\right]>0\ ,\ |\arg(z)|<\frac{1}{2}\pi\phi,$$

and $j=1,\ldots,M,\ i=1,\ldots,n\ and\ n\in N$ (5)

<u>Inequality 2</u> By putting $\alpha_i=a_i-\mu_i\xi,\ \beta=\upsilon-\mu\xi$ we get

$$\frac{\prod_{i=1}^{n}\Gamma(1+a_i-\mu_i\xi)}{\Gamma\left(\upsilon-\mu\xi+\sum_{i=1}^{n}a_i-\mu_i\xi\right)}\le\frac{\prod_{i=1}^{n}\Gamma(1+(a_i-\mu_i\xi)x)}{\Gamma\left(\upsilon-\mu\xi+x\sum_{i=1}^{n}(a_i-\mu_i\xi)\right)}\le\frac{1}{\Gamma(\upsilon-\mu\xi)}$$

multipying all three terms with

$$\frac{1}{2\pi\omega}\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}$$

and then integrating along the contour L we get

$$\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{\prod_{i=1}^{n}\Gamma(1+a_i-\mu_i\xi)}{\Gamma\left(\upsilon-\mu\xi+\sum_{i=1}^{n}(a_i-\mu_i\xi)\right)}d\xi$$

$$\le\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{\prod_{i=1}^{n}\Gamma(1+(a_i-\mu_i\xi)x)}{\Gamma\left(\upsilon-\mu\xi+x\sum_{i=1}^{n}(a_i-\mu_i\xi)\right)}d\xi$$

$$\le\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{1}{\Gamma(\upsilon-\mu\xi)}d\xi$$

=

$$I^{M+n,N}_{P_i+1,Q_i+n:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(\nu+\sum_{i=1}^{n}a_i,\ \mu+\sum_{i=1}^{n}\mu_i)\\(1+a_i,\mu_i)_{1,n};(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i}\end{matrix}\right.\right]$$

$$\le I^{M+n,N}_{P_i+1,Q_i+n:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(\nu+x\sum_{i=1}^{n}a_i,\ \mu+x\sum_{i=1}^{n}\mu_i)\\(1+a_ix,\mu_ix)_{1,n};(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i}\end{matrix}\right.\right]$$

$$\le I^{M,N}_{P_i+1,Q_i:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(\nu,\mu)\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i}\end{matrix}\right.\right]$$ (6)

Licensed Under Creative Commons Attribution CC BY

provided that
$x \in [0,1]$

$$\operatorname{Re}\left[a_i - \min_{1\le j\le M}\left(\frac{b_j}{\beta_j}\right)\right] > 0,\ \operatorname{Re}\left[\nu - \min_{1\le j\le M}\left(\frac{b_j}{\beta_j}\right)\right] > 0\ ,\ |\arg(z)| < \frac{1}{2}\pi\phi,$$

For $j = 1,\ldots,M,\ i = 1,\ldots,n$ and $n \in N$ (7)

Inequality 3 By putting $\alpha_i = \mu_i\xi + a_i,\ \beta = \upsilon + 1 + \mu\xi$ we get

$$\frac{\prod_{i=1}^{n}\Gamma(1+\mu_i\xi+a_i)}{\Gamma\left(\upsilon+1+\mu\xi+\sum_{i=1}^{n}(\mu_i\xi+a_i)\right)} \le \frac{\prod_{i=1}^{n}\Gamma(1+(\mu_i\xi+a_i)x)}{\Gamma\left(\upsilon+1+\mu\xi+x\sum_{i=1}^{n}(\mu_i\xi+a_i)\right)} \le \frac{1}{\Gamma(\upsilon+1+\mu\xi)}$$

multipying all three terms with

$$\frac{1}{2\pi\omega}\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)\Gamma\left(\left(\upsilon+\sum_{i=1}^{n}a_i\right)-\left(\mu+\sum_{i=1}^{n}\mu_i\right)\xi\right)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}$$

and then integrating along the contour L

$$\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)\Gamma\left(\left(\upsilon+\sum_{i=1}^{n}a_i\right)-\left(\mu+\sum_{i=1}^{n}\mu_i\right)\xi\right)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}$$

$$\frac{\prod_{i=1}^{n}\Gamma(1+\mu_i\xi+a_i)}{\Gamma\left(\upsilon+1+\mu\xi+\sum_{i=1}^{n}(\mu_i\xi+a_i)\right)}d\xi$$

$$\le\frac{1}{2\pi\omega}\int_L\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)\Gamma\left(\left(\upsilon+\sum_{i=1}^{n}a_i\right)-\left(\mu+\sum_{i=1}^{n}\mu_i\right)\xi\right)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}$$

$$\frac{\prod_{i=1}^{n}\Gamma(1+(\mu_i\xi+a_i)x)}{\Gamma\left(\upsilon+1+\mu\xi+x\sum_{i=1}^{n}(\mu_i\xi+a_i)\right)}d\xi$$

Licensed Under Creative Commons Attribution CC BY

$$\leq \frac{1}{2\pi\omega}\int_L \frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)\Gamma\left(\left(\upsilon+\sum_{i=1}^{n}a_i\right)-\left(\mu+\sum_{i=1}^{n}\mu_i\right)\xi\right)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\frac{1}{\Gamma(\upsilon+1+\mu\xi)}d\xi$$

=

$$I^{M+1,N+n}_{P_i+n,Q_i+2:r}\left[z\left|\begin{matrix}(-a_i,\mu_i)_{1,n};(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\(\nu+\sum_{i=1}^{n}a_i',\mu+\sum_{i=1}^{n}\mu_i);(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(-\nu-\sum_{i=1}^{n}a_i,\mu+\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\leq I^{M+1,N+n}_{P_i+n,Q_i+2:r}\left[z\left|\begin{matrix}(-a_ix,x\mu_i)_{1,n};(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\\left(\nu+\sum_{i=1}^{n}a_i',\left(\mu+\sum_{i=1}^{n}\mu_i\right)\right);(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};\left(-\nu-x\sum_{i=1}^{n}a_i',\left(\mu+x\sum_{i=1}^{n}\mu_i\right)\right)\end{matrix}\right.\right]$$

$$\leq I^{M+1,N}_{P_i,Q_i+2:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i}\\\left(\nu+\sum_{i=1}^{n}a_i',\left(\mu+\sum_{i=1}^{n}\mu_i\right)\right);(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(-\nu,\mu)\end{matrix}\right.\right] \quad (8)$$

provided that

$x\in[0,1]$

$$\mathrm{Re}\left[\nu-\min_{1\leq j\leq M}\mu_i\left(\frac{b_j}{\beta_j}\right)\right]>1,\ \mathrm{Re}\left[a_i+\min_{1\leq j\leq M}\mu\left(\frac{b_j}{\beta_j}\right)\right]>0\ ,\ |\arg(z)|<\frac{1}{2}\pi\phi,$$

For $j=1,\ldots,M,\ i=1,\ldots,n$ *and* $n\in N$ (9)

<u>Inequality 4</u> By putting $\alpha_i=a_i-\mu_i\xi,\ \beta=\upsilon-\mu\xi$ we get

$$\frac{\prod_{i=1}^{n}\Gamma(1+a_i-\mu_i\xi)}{\Gamma\left(\upsilon-\mu\xi+\sum_{i=1}^{n}(a_i-\mu_i\xi)\right)}\leq\frac{\prod_{i=1}^{n}\Gamma(1+(a_i-\mu_i\xi)x)}{\Gamma\left(\upsilon-\mu\xi+x\sum_{i=1}^{n}(a_i-\mu_i\xi)\right)}\leq\frac{1}{\Gamma(\upsilon-\mu\xi)}$$

multipying all three terms with

$$\frac{1}{2\pi\omega}\frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\times\frac{1}{\Gamma(1+\upsilon+\sum_{i=1}^{n}a_i+\left(\mu-\sum_{i=1}^{n}\mu_i\right)\xi)}$$

and then integrating along the contour L

Licensed Under Creative Commons Attribution CC BY

$$\frac{1}{2\pi\omega}\int_L \frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\times\frac{1}{\Gamma(1+\upsilon+\sum_{i=1}^{n}a_i+\left(\mu-\sum_{i=1}^{n}\mu_i\right)\xi)}$$

$$\frac{\prod_{i=1}^{n}\Gamma\left(1+a_i-\mu_i\xi\right)}{\Gamma\left(\upsilon-\mu\xi+\sum_{i=1}^{n}a_i-\mu_i\xi\right)}d\xi$$

$$\leq\frac{1}{2\pi\omega}\int_L \frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\times\frac{1}{\Gamma(1+\upsilon+\sum_{i=1}^{n}a_i+\left(\mu-\sum_{i=1}^{n}\mu_i\right)\xi)}$$

$$\frac{\prod_{i=1}^{n}\Gamma\left(1+\left(a_i-\mu_i\xi\right)x\right)}{\Gamma\left(\upsilon-\mu\xi+x\sum_{i=1}^{n}\left(a_i-\mu_i\xi\right)\right)}d\xi$$

$$\leq\frac{1}{2\pi\omega}\int_L \frac{\prod_{j=1}^{M}\Gamma(b_j-\beta_j\xi)\prod_{j=1}^{N}\Gamma(1-a_j+\alpha_j\xi)}{\sum_{i=1}^{r}\left\{\prod_{j=M+1}^{Q_i}\Gamma(1-b_{ji}+\beta_{ji}\xi)\prod_{j=N+1}^{P_i}\Gamma(a_{ji}-\alpha_{ji}\xi)\right\}}\times\frac{1}{\Gamma(1+\upsilon+\sum_{i=1}^{n}a_i+\left(\mu-\sum_{i=1}^{n}\mu_i\right)\xi)}$$

$$\frac{1}{\Gamma\left(\upsilon-\mu\xi\right)}d\xi$$

=

$$I^{M+n,N}_{P_i+1,Q_i+n+1:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(-\nu-\sum_{i=1}^{n}a_i,\ \mu-\sum_{i=1}^{n}\mu_i))\\(1+a_i,\mu_i)_{1,n};(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(\nu+\sum_{i=1}^{n}a'_i,\ \mu+\sum_{i=1}^{n}\mu_i\end{matrix}\right.\right]$$

$$\leq I^{M+n,N}_{P_i+1,Q_i+n+1:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(-\nu-x\sum_{i=1}^{n}a_i,\ \mu-x\sum_{i=1}^{n}\mu_i)\\(1+a_ix,\mu_ix)_{1,n};(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(\nu+\sum_{i=1}^{n}a'_i,\ \mu+\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\leq I^{M,N}_{P_i+1,Q_i+1:r}\left[z\left|\begin{matrix}(a_j,\alpha_j)_{1,N};(a_{ji},\alpha_{ji})_{N+1,P_i};(\nu,\mu)\\(b_j,\beta_j)_{1,M};(b_{ji},\beta_{ji})_{M+1,Q_i};(\nu+\sum_{i=1}^{n}a'_i,\ \mu+\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right] \quad (10)$$

provided that $x\in[0,1]$

Licensed Under Creative Commons Attribution CC BY

$$\mathrm{Re}\left[a_i - \min_{1\le j\le M}\left(\frac{b_j}{\beta_j}\right)\right] > 0,\ \mathrm{Re}\left[\nu + \min_{1\le j\le M}\left(\frac{b_j}{\beta_j}\right)\right] > 1,\ |\arg(z)| < \frac{1}{2}\pi\phi,$$

For $j = 1,\ldots,M,\ i = 1,\ldots,n \text{ and } n \in N$ (11)

Particular cases :-

(i) On setting r = 1 in the inequalities, the I-function occurring therein reduces to H-function and we arrive at the results due to[1].

(ii) If we put $r = 1, \alpha_j = \beta_j = \alpha_{ji} = \beta_{ji} = \alpha$ in the inequalities, the I-function reduces to a Meijer's G-function[3].

$$G^{M,N+n}_{P+n,Q+1}\left[z\left|\begin{matrix}(a_i,\mu_i)_{1,n};(a_j,\alpha)_{1,N};(a_j,\alpha)_{N+1,P}\\(b_j,\alpha)_{1,M};(b_j,\alpha)_{M+1,Q};\ (-\nu+\sum_{i=1}^{n}a_i,\ \mu+\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\le G^{M,N+n}_{P+n,Q+1}\left[z\left|\begin{matrix}(a_i x,\mu_i x)_{1,n};(a_j,\alpha)_{1,N};(a_j,\alpha)_{N+1,P}\\(b_j,\alpha)_{1,M};(b_j,\alpha)_{M+1,Q};\ (-\nu+x\sum_{i=1}^{n}a_i,\ \mu+x\sum_{i=1}^{n}\mu_i)\end{matrix}\right.\right]$$

$$\le G^{M,N}_{P,Q+1}\left[z\left|\begin{matrix}(a_j,\alpha)_{1,N};(a_j,\alpha)_{N+1,P}\\(b_j,\alpha)_{1,M};(b_j,\alpha)_{M+1,Q};\ (-\nu,\mu)\end{matrix}\right.\right]$$

References

[1] Ronghe, A.K., Kulwant Kaur and Arati Baurase, Double inequalities for the Fox's H-function.Volume4, No. 3(2012), pp.313-318.

[2] Nguyen Van Vinh and An inequality for the gamma function. Ngo Phuoc Nguyen Ngoc(2009)international mathematical forum 4 no. 28, 1379-1382.

[3] Saxena V.P. The I-Function, Anamaya Publishers, New Delhi, (2008).

Licensed Under Creative Commons Attribution CC BY