

Travelling Salesman Problem (TSP) Using Fuzzy Quantifier

G. Nirmala¹, R. Anju²

¹Associate Professor of Mathematics, PG & Research, Dept. of Mathematics, K.N.G. Arts College for Women (Autonomous) Thanjavur-613007, Tamil Nadu, India

²Lecturer in Mathematics, Bharathidasan University Constituent College (W), Orathanadu, Tamil Nadu, India

Abstract: In this paper we have preferred the fuzzy quantifier and ranking method for finding the initial solution, also Nearest Neighbour Method (NN-Method) is applied for finding an optimal solution for Travelling Salesman Problem (TSP). This method requires least iterations to reach the optimality. Results are given to check the validity of the proposed method.

Keywords: TSP, Fuzzy quantifier, ranking function, NN-method, optimal solution.

1. Introduction

Travelling Salesman Problem (TSP) is a well known, popular and extensively studied problem in the field of conditional optimization and attracts computer scientists, mathematicians and others. Its statement is deceptively simple, but yet it remains one of the most challenging problems in operational research. It also an optimization problem of finding a shortest closed tour that visits all the given cities. It is known as a classical NP-Complete problem, which has extremely large search space and is very difficult to solve.

The definition of a TSP is : given N cities, if a salesman starting from his home city is to visit each city exactly once and then return home, find the order of a tour such that the total distances (cost) travelled is minimum. Cost can be distance, time, money, energy, etc. TSP is an NP-hard Problem and researchers especially mathematicians and scientists have been studying to develop efficient solving methods since 1950's. Because it is so easy to describe and so difficult to solve. The travelling salesman problem is widespread in engineering applications. It has been employed in designing hardware devices and radio electronic devices, in communications in the architecture of computational networks etc. In addition, some industrial problems such as machine scheduling, cellular manufacturing and frequency assignment problems can be formulated as a TSP. In this paper we investigate a TSP with fuzzy cost \tilde{C}_{ij} represented by fuzzy quantifier which are replaced by triangular or trapezoidal fuzzy numbers.

2. Preliminaries

In 1965, Zadeh introduced the concept of a fuzzy set as a mathematical way of representing impreciseness in real world problems.

2.1 Definition

A fuzzy set \tilde{A} in a universe of discourse X is defined by

$$\tilde{A} = \{ \langle x, \mu_{\tilde{A}}(x) \rangle / x \in X \} \text{ where } \mu_{\tilde{A}}(x) : X \rightarrow [0,1]$$

is called the membership function of \tilde{A} and $\mu_{\tilde{A}}(x)$ is the degree of membership to which $x \in \tilde{A}$.

2.2 Definition

A fuzzy set \tilde{A} on R is convex iff for any $x_1, x_2 \in X$, the membership function of \tilde{A} satisfies the inequality

$$\mu_{\tilde{A}}(\lambda x_1 + (1 - \lambda)x_2) \geq \min\{\mu_{\tilde{A}}(x_1), \mu_{\tilde{A}}(x_2)\}; 0 \leq \lambda \leq 1.$$

2.3 Definition

A fuzzy set \tilde{A} of the universe of discourse X is called a normal fuzzy set if there exists at least one $x \in X$ such that $\mu_{\tilde{A}}(x) = 1$.

2.4 Definition

A fuzzy set \tilde{A} , is defined on the universal set of real number R, is said to be a fuzzy number, if its membership function has the following characteristics

- (i) \tilde{A} , is convex
- (ii) \tilde{A} , is normal and
- (iii) $\mu_{\tilde{A}}(x)$ is piecewise continuous.

2.5 Definition

A fuzzy number $\tilde{A} = (a, b, c)$ is said to be a triangular fuzzy number if its membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} \frac{x-a}{b-a}, & a \leq x \leq b \\ 1 & x = b \\ \frac{c-x}{c-b}, & b \leq x \leq c \end{cases}$$

2.6 Definition

The α –cut set of a fuzzy set \tilde{A} is a crisp set defined by $\tilde{A}_\alpha = \{x \in X / \mu_{\tilde{A}}(x) \geq \alpha\}$.

2.7 Definition

For a convex fuzzy \tilde{a} , the Robust’s Ranking index is defined by

$$R(\tilde{a}) = \int_0^1 0.5(a_\alpha^L, a_\alpha^U) d \alpha$$

where (a_α^L, a_α^U) is the α -level cut of the fuzzy number \tilde{a} .

Robust’s ranking technique which satisfy compensation, linearity and additive properties and provides results which are consist human intuition.

3. Fuzzy Quantifier

Fuzzy quantifiers are fuzzy numbers that take part in fuzzy propositions. Fuzzy quantifiers that characterize linguistic terms such as about 10, much more than 100, atleast about 5 of the first kind and almost all, about half, most and so on of the second kind.

One of them is the form

P: There are Q_i ’s in I such that $v(i)$ of v is F,

$$\tilde{C}_{ij} = \begin{matrix} & D_1 & D_2 & D_3 & D_4 \\ \begin{matrix} O_1 \\ O_2 \\ O_3 \\ O_4 \end{matrix} & \begin{pmatrix} Nil & Substantially low & Low & Above average \\ Substantial & Nil & Below average & High \\ Above average & Substantially high & Nil & Average \\ Somewhat low & High & Somewhat high & Nil \end{pmatrix} \end{matrix}$$

Solution

The linguistic variables showing the qualitative data is converted into quantitative data using the following table. The linguistic variables are represented by triangular fuzzy numbers.

Table 1

Nil(Prohibited/not permitted)	∞
Substantially low	(-2,1,4)
Somewhat low	(-2,3,8)
low	(1,4,7)
Below average	(2,5,8)
Average	(5,6,7)
Above average	(6,7,8)
Substantial	(6,10,14)
High	(7,8,9)
Somewhat high	(8,9,10)
Substantially high	(10,12,14)

Now from table 1 we have

$$\tilde{C}_{ij} = \begin{matrix} & D1 & D2 & D3 & D4 \\ \begin{matrix} O_1 \\ O_2 \\ O_3 \\ O_4 \end{matrix} & \begin{pmatrix} \infty & (-2,1,4) & (1,4,7) & (6,7,8) \\ (6,10,14) & \infty & (2,5,8) & (7,8,9) \\ (6,7,8) & (10,12,14) & \infty & (5,6,7) \\ (-2,3,8) & (7,8,9) & (8,9,10) & \infty \end{pmatrix} \end{matrix}$$

where $v(i)$ of v is a variable , Q is a fuzzy number expressing linguistic term, F is a fuzzy set of variable v .

4. Nearest Neighbour Method

Step 1: Locate the smallest element in the cost matrix (break ties arbitrarily) circle it, and include the corresponding link in the itinerary.

Step 2: If the newly circled element is C_{pq} , replace all other elements in the p^{th} row and all other element in the q^{th} column, as well as the transposed element C_{pqq} , by a prohibitively large number.

Step 3: Locate the smallest uncircled element in the latest cost matrix. Tentatively adjoin its corresponding link to the (incomplete) itinerary. If the resulting itinerary is feasible, circle the designated cost and go to step 5.

Step 4: If the resulting itinerary is infeasible, remove the latest link from the itinerary and replace its corresponding cost by a prohibitively large number. Go to step 3.

Step 5: Determine whether the itinerary is complete. If so, accept it as the near-optimal one. If not, go to step 2.

5. Result: TSP- A Logistic Problem

Consider the following Travelling Salesman Problem. Here the distance travelled (cost) are represented by fuzzy quantifiers which characterize the linguistic variables are replaced by fuzzy numbers. The problem is then solved by using the proposed method.

Now, using Robust Ranking technique the above fuzzy TSP is

$$\begin{matrix} & 1 & 2 & 3 & 4 \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{pmatrix} \infty & 1 & 4 & 7 \\ 10 & \infty & 5 & 8 \\ 7 & 12 & \infty & 6 \\ 3 & 8 & 9 & \infty \end{pmatrix} \end{matrix}$$

and it is given in the following figure.




Figure 1: TSP – A Logistic problem

After applying NN Method, the fuzzy optimal total cost is $X_{12}= 1, X_{23}= 5, X_{34}= 6, X_{41}= 3$

$$\begin{aligned} \text{i.e., } & R(-2,1,4) + R(2,5,8) + R(5,6,7) + R(-2,3,8) = R(3,15,27) \\ & = 1 + 5 + 6 + 3 \\ & = \text{Rs. } 15 \end{aligned}$$

Therefore the near-optimal solution is

$$\begin{aligned} & 1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 4, 4 \rightarrow 1 \text{ (or)} \\ & 1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 1 \end{aligned}$$

$$\text{i.e., } O_1 \rightarrow D_2, O_2 \rightarrow D_3, O_3 \rightarrow D_4, O_4 \rightarrow D_1$$

which is complete and it is given in the above figure.

6. Conclusion

In this paper the travelling costs are considered as fuzzy quantifiers that characterize linguistic variables are represented by triangular fuzzy numbers. Moreover, the fuzzy TSP has been transformed into crisp TSP using Robust's Ranking indices. We proposed a Nearest Neighbour Method to find an optimal solution for fuzzy TSP. Numerical example is solved using the proposed method and the results are obtained optimally. Thus the proposed method carries systematic procedure and very easy to understand. It can be extended to assignment problem and transportation problems to get optimal solution. The proposed method is an important tool for the decision makers when they are handling various types of logistic problems to make decision optimally.

References

[1] Zadeh, L.A.,(1965), Fuzzy sets, Information and Control, Vol.8, 338-353.
 [2] Zimmermann H.J., Fuzzy Programming and linear programming with several objective functions, Fuzzy sets and systems: (1978), 45-55.
 [3] Zimmermann H.J., Fuzzy Mathematical programming, computer science and operations Research Vol.10, No.4,(1983) 291-298.

[4] Zimmermann H.J., (1985) Application of Fuzzy set theory to Mathematical Programming.
 [5] T. Bektas., The multiple travelling salesman problem an overview of formulations and solution procedures omega 34, 209-219, 2006.
 [6] W.J. Cook, “ In Pursuit of the travelling salesman: Mathematics at the Limits of computation”, Princeton University Press, Princeton, 2011.
 [7] H. Fort, M. Kornbluth and F. Zypman, “Travelling Salesman Problem for Finite-Size cities”, Mathematical Structures in computer science, Vol. 20, No.6, 2010, pp.1067-1078.
 [8] Gilad Barach, Hugo Fort, Yoni Mehlman, Fredy Zypman, Information in the Travelling Salesman Problem, Applied Mathematics, 2012,3, 926-930.
 [9] S. Lin, “Computer solutions of the Travelling Salesman Problem”, Bell systems Technical Journal, Vol.44, 1965, pp.2245-2269.
 [10] David S Johnson : Travelling Salesman Problem, DIMACS Implementation challenge, AT & T (2001).
 [11] G. Bezael and S. Kizhanathan : An optimal solution method for large – scale multiple travelling salesman problems Operations Research, 34(5): (1986). 698-718.
 [12] Lawler E.L., Lenstra J.K., Rinooy Kan A.H.G., and Shmoys D.B. eds., The Travelling salesman problem, John Wiley, Chichester UK, 1985.
 [13] G. Nirmala and R. Anju, Fuzzy Rule based inference decision making process, The PMU Journal of Humanities and Sciences, Vol.3, No.1, Jan-Jun 2012.ISSN:0976-1853.
 [14] G. Nirmala and R. Anju, An Application of Fuzzy quantifier in sequencing problem with fuzzy ranking method, Aryabhata Journal of Mathematics and Informatics, Vol. 6, No.1, Jan-July 2014.
 [15] G. Nirmala and R. Anju, Decision making of conducting remedial classes for weak students through fuzzy ingredients, Internation Journal of Scientific and Research Publications, Vol. 3, issue 6, June 2013.

Author Profile



Dr. G. Nirmala is currently working as Research Supervisor & convenor, Head & Associate Professor in the department of Mathematics, K.N. Govt. Arts College (W), Autonomous, Thanjavur-613 007, Tamil Nadu, India. She has got 30 years of teaching experience and 15 years of research experience. Her field of interest is Genetic Algorithm with Fuzzy Neural Network. She has published more than 110 research articles. She has received two national awards.



R. Anju is Lecturer in the department of Mathematics, BDUCC(W), Orathanadu, Thanjavur, Tamil Nadu, India. She has obtained her master degree in 2005 and M.Phil degree in 2007 from Bharathidasan University, Tiruchirappalli. She has got 8 years of teaching experience.