

# Population fluctuation of *Apis cerana* Fabr. (Hymenoptera: Apidae) Colonies on Policulture Plantations in West Sumatra

Jasmi<sup>1</sup>, Siti Salmah<sup>2</sup>, Dahelmi<sup>2</sup>, Syamsuardi<sup>2</sup>

<sup>1</sup>Department of Biology Education STKIP PGRI West Sumatra, Gunung Pangilun Street, Padang 25137, Indonesia

<sup>2</sup>Department of Biology, Faculty of Mathematics and Science, Andalas University, Limau Manis, Padang, West Sumatra, Indonesia

**Abstract:** Population fluctuation of *Apis cerana* Fabr. (Hymenoptera: Apidae) colonies on policulture plantations was studied in West Sumatra, Indonesia. Census method was carried out to determine species of plants used as nesting sites and colony population. Data were taken from low (altitude < 100 meter) and high (altitude > 1000 meter) lands, each with five plantations (each 10.000 m<sup>2</sup> width). Multiple regression was used to analyze data on population density of colonies and their relations with weather factors. Results indicated that 18 species of plants in 15 families were used as nesting sites. Colony populations of *Apis cerana* in lowlands were higher (4,06 ± 0,56 colonies per 10.000 m<sup>2</sup>) than the one in high lands (3,17 ± 0,54 colonies per 10.000 m<sup>2</sup>). Colony population densities fluctuated based on time, the highest population was found in June and the lowest was in December and January. Weather factors showed more influence on colony population densities in highlands (R<sup>2</sup> = 95,03%) compared to lowlands (R<sup>2</sup> = 54,97%).

**Keywords:** Honeybee, nesting diversity, altitude, colony fluctuation

## 1. Introduction

*Apis cerana* is honey bee originated from Asia distributed worldwide [1]. It has some superiorities, i.e. as good pollinators for coconut and coffee plants [2,3], easy to be reared [4] and it has organized self defence strategy from hymenopteran predators [5, 6]. This species of bee is still widely distributed in different habitats in West Sumatra [7] with the highest colony population density 22 nests/km<sup>2</sup> [8].

Polyculture plantations play important role in conserving the population of honey bee. They function as a habitat to carry out activities like making nests, foraging for food, and colony multiplications. On the other sides the presence of honey bee is very important to increase plant productions, mainly on *Coffea canephora* and *Coffea arabica* [3], apple [9], highbush blueberry (*Vaccinium corymbosum* L.) [10], and palm [11]. Due their importance, some studies on ecology, spatial distribution, diversity, and conservation of this bee and other bees have been gaining pace in recent years [8, 12, 13, 14, 15, 16].

Population colony density of different species of honey bee varies in every habitat [8, 12, 17]. The density is affected by various factors such as queen productivity [18], colony migrations [19, 20], food resources [21, 22, 23, 24, 25], pests and diseases [26, 27, 28, 29, 30, 31, 33] and environment factors [33].

Polyculture plantations is one of habitats of *Apis cerana* in West Sumatra [7]. The width of polyculture plantations is the third (5,93%) out of total land usage in West Sumatra [34]. Some good points of polyculture plantations are the presence of various species of plants, various planting

dates, low frequency of pesticide applications, very scarce rotation of major plants, and plantations are distributed in all altitudes. Plants diversity and weather factors at different altitudes affect the density of *A. cerana* colony population. This paper reports the diversity of nesting sites, density and fluctuation of colony population of *A. cerana* and the impact of weather factors, in tropical polyculture plantations in West Sumatra, Indonesia

## 2. Materials and Methods

### 2.1. Study Sites

The research was conducted in polyculture plantations in West Sumatra, from March 2011-March 2012. Observations were conducted in low and highlands (Fig. 1). The lowlands site was in Nagari Parik

Malintang Kec. 2x 11 Enam Lingkung, Kabupaten Padang Pariaman, astronomical position at 100°27' 00" east longitude and 0°50'30" south latitude, altitude < 100 m a.s.l., dominated with *Durio zibetinus*, *Spondias pinnata* Kurz, *Areca catecu*, *Cocos nucifera* dan *Theobroma cacao*. Average temperature was 25,7°C and monthly rain fall was 368,4 mm. The high land observation was conducted in Nagari Andaleh Kecamatan Batipuh, Kabupaten Tanah Datar, astronomical position at 100°22' " east longitude and 0°23'38" south latitude, altitude > 1000 m a.s.l., dominated with *Coffea canephora* and *Cinnamomum burmanii*. Average daily temperature 25,0°C and monthly rain fall was 549,00 mm [34]. The dominant weed species in highlands were *Bidens pilosa* and *Galiana soga parviflora*, while the ones in lowlands were *Cynodon dactylon* and *Mimosa pudica*.

## 2.2. Colonies Census

The data of *A. cerana* colonies were taken through census. Site criteria were: minimum width 10.000 m<sup>2</sup>, there were trees with minimum stem diameter > 20 cm., dominated with cultured plants for fruit productions purpose, at least there was minimum one colony of *A. cerana* found. The plot size for every sample point was 100 x 100 m. There were 10 plots, 5 for each lowlands and highlands. All trees with diameter > 10 cm were observed. Parameters observed were place and species of plants used for nesting sites and number of colonies per plot. The species of plants for nesting were identified in Laboratory Plants Taxonomy Andalas University, Padang.

## 2.3. Colony fluctuation observations

The observation was conducted through census. The number of colonies of *A. cerana* was counted once in a month for 13 month periode. Parameters measured were colony numbers (old and new nests) and absconding colonies. Secondary data like rain fall ( $x_1$ , mmHg), relative humidity ( $x_2$ , %), temperature ( $x_3$ , °C), wind flow ( $x_4$ , knot) and photoperiode ( $x_5$ , %) were obtained from Agency for Meterology, Climatology and Geophysics, station of Padang Panjang (highland) and Sicincin Padang Pariaman (lowland).

## 2.4. Data Analisis

Colonies fluctuations were analysed with Mann-Wit test. Correlation between colony population density (Y) with all weather factors, rain fall ( $x_1$ , mmHg), relative humidity ( $x_2$ , %), temperature ( $x_3$ , °C), wind flow ( $x_4$ , knot) and photoperiode ( $x_5$ , %) is presented in Equation 1. Degree of correlation is expressed as koefisien determination (R) with a rank 0-100 % [35].

$$Y = a + b_1x_1 + b_2x_2 + b_3x_3 + b_4x_4 + b_5x_5 \dots \dots \dots \text{(Equation 1)}$$

## 3. Result

### 3.1. Nesting Sites Diversity

Nesting sites of *A. cerana* were found in 18 species of plants (15 families) in polyculture plantations in West Sumatera. Fifty five colonies of *A. cerana* were found in tree cavities, 12 colonies in *Erythria variegata* (Dadap), 11 colonies in *Cocos nucifera* (coconut), 6 colonies in *Areca catechu* (Pinang), 4 colonies in *Toona sureni* (Surian), 3 colonies in *Peronema canescens* (Sungkai), 2 colonies in each of the following plants, *Durio zibethinus* (Durian), *E. acuminata* (Jirak), *G. mangostana* (Manggis), *Phithocellobium lithosperma* (Jengkol), *Spondias lutea* (Kedondong), and 1 colony in each of the following plants, *Arthrocarpus* sp. (Sukun), *Caryota rumphiana* (Rumbia), *Coffea robusta* (Kopi), *S. reticulata* (Kepundung hutan), *Lansium domesticum* (Duku), *Musca* sp. (Pisang), *Syzygium aqueum* (jambu air) and *Toxicodendron radicans* (Jelatang). Nests were also found

in house wall, rat hollow, used chair and electric pool made of cement. There was only one nest found on each of above sites. The highest number of nests were found in *E. variegata* (highland) and *C. nucifera* (lowland).

### 2.2. Colony Population Density

Average colony population density of *A. cerana* in polyculture plantations in West Sumatera ( $4,06 \pm 0,65$  colonies per 10.000 m<sup>2</sup>). Average colony population density in lowland polyculture plantations was higher ( $4,06 \pm 0,56$  colony per 10.000 m<sup>2</sup>) than the one in highlands ( $3,17 \pm 0,54$  colonies per 10.000 m<sup>2</sup>). The highest colony population density in lowlands was found in June ( $5,2 \pm 0,84$  colonies per 10.000 m<sup>2</sup>) and in highlands in May ( $4,4 \pm 0,55$  colonies per 10.000 m<sup>2</sup>) (Fig. 2a).

Average density colony population *A. cerana* in polyculture plantations in West Sumatera fluctuated throughout the year (Fig. 2). Colony population density in highlands increased from  $4.0 \pm 0.71$  colonies in April to  $4.4 \pm 0.55$  colonies per 10.000 m<sup>2</sup> in June and in lowlands increased from  $4.6 \pm 0.55$  colonies in April to  $5.2 \pm 0.84$  colonies per 10.000 m<sup>2</sup> in June.

The significant fluctuation of colony population density of *A. cerana* in polyculture plantations in highlands (Fig 2a) ( $p=0.01$ ) occurred from October 2011 ( $2.6 \pm 1.67$  colonies per 10.000 m<sup>2</sup>) to November 2011 ( $2.2 \pm 1.30$  colonies per 10.000 m<sup>2</sup>) and March 2012 ( $2.2 \pm 0.84$  colonies per 10.000 m<sup>2</sup>). Significant fluctuation of colony population density in lowlands (Fig. 2b) ( $p=0.01$ ) occurred from November ( $3.8 \pm 1.30$  colonies per 10.000 m<sup>2</sup>) to December 2011 ( $2.8 \pm 0.45$  colonies per 10.000 m<sup>2</sup>) and January 2012 ( $2.8 \pm 0.84$  colonies per 10.000 m<sup>2</sup>).

### 3.3. Correlation between colony population density and weather factors

There was highly significant correlation between colony population density of *A. cerana* and weather factors in highland polyculture plantations (Equation 2) and loose correlation in lowlands (Equation 3). Interaction of weather factors in high lands ( $R^2=95.03\%$ ) determined *A. cerana* colonies density more than the one in lowlands ( $R^2=54.97\%$ ). The difference of determination coefficient between the two locations due to different altitudes. The highlands were located >1000 meter a.s.l. and low lands < 100 meter a.s.l. Different altitude caused different weather conditions in the two locations (Table 1).

$$Y = 47.814 + 0.007x_1 - 0.664x_2 + 1.425x_3 - 4.365x_4 + 0.236x_5 \quad (R^2 = 0.95) \dots \dots \dots \text{(Equation 2)}$$

$$Y = 87.807 - 0.0005x_1 + 1.2776x_2 + 1.845x_3 - 1.360x_4 - 0.044x_5 \quad (R^2 = 0.356) \dots \dots \dots \text{(Equation 3)}$$

## 4. Discussion

The diversity of plant species used as nesting sites of *A. cerana* in polyculture plantations in West Sumatra was higher compared to the past finding either of the same species in Padang Sarai, Pasir Jambak and Bungo Pasang Padang [8] or different species of bee in Kalimantan Selatan [36]. The use of *C. nucifer* as nesting sites was also reported by Inoue *et al.* [8]. *Toxicodendron radicans* dan *P. lobatum* were also used as nesting site of *A. koschevnikovi* [36].

The highest number of *A. cerana* nests was found in *Erythria variegata*, i.e. 12 out of 24 nests found in high lands. The preference of *A. cerana* to place nests in above tree due to the availability of a lot of cavities on the trees and due to the abundance of the trees where food resources was available. *Erythria variegata* is planted as a shading plants for young coffee trees but the shading plants are let to grow naturally until dead. *Erythria variegata* has soft stem tissues easily destroy that produce cavities. According to Baum *et al.* [12] colony aggregations probably resulted from the distribution of resources, especially cavities.

The highest number of *A. cerana* nests found in *Cocos nucifera* due to their dominant number of trees found in polyculture plantations in low lands and they have relatively more cavities. The outer part structure of *C. nucifera* is hard while inner parts is soft. This condition facilitates the formation of cavities well protected by the hard outer parts. Other reason is *C. nucifera* also produces pollen and nectar throughout the year. Inoue *et al.* [8] also found 13 nests of *A. cerana indica* in tree cavity in Central Sumatra. Vaudo *et al.* [16] reported colonies nesting on the reserves may occur in greater densities than those nesting on livestock farms. Donaldson-Matasci & Dornhaus [37] also reported two important characteristics of honey bees' ancestral ecological habitat that could have selected for their unique ability to symbolically communicate resource location. Honey-bees of the genus *Apis* have been recorded as potential pollinators in a few cases such as *Serenoa repens* and in cultivated palms such as *Cocos nucifera* [38].

### 4.1 Colony population density of *A. cerana*

Average colony population density of *A. cerana* in polyculture plantations in West Sumatra ( $4,06 \pm 0.65$  colonies per 10.000 m<sup>2</sup>). This number was higher than the one found by Inoue *et al.* [8] (1990) which was 22 colonies of *A. cerana indica* per km<sup>2</sup> or equal to 2.2 colonies per 10.000 m<sup>2</sup>. Baum *et al.* [12] found 12.5 colonies of *A. mellifera* per km<sup>2</sup> or equal to 1.25 colonies per 10.000 m<sup>2</sup> in coastal prairie landscape. On horticultural plants in Tropical fruit research centre in southeast Thailand it was found 2 colonies *A. dorsata*, 5 colonies *A. cerana*, 12 colonies *A. andreniformis* dan 12 colonies *A. florea* per km<sup>2</sup> or equal to 0.2 colonies *A. dorsata*, 0.5 colonies *A. cerana*, 1.2 colonies *A. andreniformis*, 1.2 colonies *A. florea* per 10.000 m<sup>2</sup> [17].

Average colony population density in lowland polyculture plantations was higher than the one in highlands. The higher colony density in lowlands was because of supporting weather factors (Table 1). According to Conte and Navajas [33] environmental changes have a direct influence on honey bee development. Chagnon [39] reported the weather conditions have a direct influence on pollinator flight and an indirect effect on pollen and nectar production by the flowers selected by the pollinators for foraging. Temperature constitutes an important factor potentially limiting insect flight and pollen availability, but light intensity, rain and relative humidity also play a role.

Average density colony population *A. cerana* in polyculture plantations in West Sumatra fluctuated throughout the year (Fig. 2). Colony population density in highlands and lowlands increased from April to June. The increase of colony population until June was assumed that some colonies multiplied in the observed locations. In four months there had been colonies formed gyna. When a gyna emerged in a colony, thus the colony would swarm. Time needed by a colony to form gyna was around one month. According to Rhodes *et al.* [40] the survival of honey bee *Apis mellifera* queens to 14 days and 15 weeks after introduction into an established bee colony increases with increasing age of the queen at introduction. Survival rates increased strongly to high levels for queen bees introduced between 7 and 24 days of age and at a slower rate for queens introduced at ages up to 35 days.

The increase of *A. cerana* colony numbers in polyculture plantations could also be due to immigration. During observation period three colonies of *A. cerana* were found to move to observation areas in May. Migration colony was aimed at looking for new nest sites supported by abundance food sources. The colony rested on branch or parts of tree waiting for the signal from the guard bees. If guard bees found suitable cavities they would move to new nests. According to Ruttner [1] seasonal migration of bee colonies is a common characteristic of all tropical honey bee. Woyke *et al.* [20] reported after environmental conditions deteriorate, all the bees with their queens abscond and migrate to alternate seasonal nesting sites. The next season, the swarms do not return to their original reproductive natal sites, but to those sites they occupied the previous season lately, where from they absconded.

The fluctuation of daily temperature followed the weather changes in tropical area. From March to August it is a dry season in tropical areas and the changes are also followed by the increase of daily temperature in the observation areas. The increase in average daily temperature has positive impact on the availability of food either quantity or quality. Quantitatively the plants having blooming period in dry season would produce more flowers in certain period. Qualitatively the temperature increase would increase the nectar sugar concentration that produce better nectar quality. As a result it would be preferred by bees. Bee colonies would

take advantage from the available food sources to increase the queen productivity and to swarm. According to Kjølhl *et al.* [41] Weather change can be simulated by distributing experimental plots along natural climatic gradients or by creating different climatic conditions in artificial environments such as laboratory or greenhouse experiment. Le Conte dan Navajas [33] reported tropical weathers may evolve towards more distinct seasons with dry periods. In this case, Asian honey bees would need to rapidly step up their honey-harvesting strategy to amass sufficient stores to survive periods without flowers.

The significant fluctuation of colony population density of *A. cerana* in Polyculture plantations in high lands (Fig. 2a) occurred from October to November 2011 and March 2012. The significant decrease of average colony population density in high lands due to various things such as man activities, weather condition and problem of pests and diseases. From September until October 2011 four colonies were found absconding because of herbicide application in high land polyculture plantations. Arzone [42] reported the use of commercial herbicides as recommended by spraying the bees can cause death after 12 hours of use. High toxicity of insecticides besides generally harmful and also directs selective insect emergence.

Significant fluctuation of colony population density in low lands (Fig. 2b) occurred from November to December 2011 and January 2012. Decrease colony population density in low lands due to man activities like burning tree colonies. Kremen *et al.* [43] reported human impacts have modified the landscape through fragmentation, degradation and destruction of natural habitats and the creation of new anthropogenic habitats. Changes in land-use and landscape structure influence pollinators, target plants and their interactions at individual, population and community scales.

Other factor decreasing colony population of *A. cerana* in polyculture plantations in December 2011 and January 2012 (Fig. 2) was the presence of pest, wax moths and ants. According to Gulati and Kaushik [26] among several limiting factors, honeybee enemies constitute a major factor. Wax moths and wasps cause heavy losses to beekeepers throughout the world. Ellis *et al.* [32] reported wax moths remain a vexing problem for beekeepers and honey bee colonies around the globe. According to Gulati and Kaushik [26] ants are not usually serious pests in honeybee colonies. Ants are typically found between the inner and outer covers of the hive and in pollen traps. The small red household ant, *Dorylus labiatus* and small black ants, *Monomorium indicum*, *M. destructor* are some of the other ant species which visit bee colony for food purposes.

#### 4.2 Correlation between colony population density and weather factors

Wind was one of weather factors which highly significant influenced and had negative impact on colony population density in highlands (Table 1 and Equation 1). Strong wind broke the branches or the plants mainly the dead ones that caused the colonies to abscond. During observation there were 4 dead trees used for nest in high lands and one tree in low land that were broken by the strong wind. The strong wind also affected nests directly by shaking them thus it caused the colony to abscond. According to Chagnon [39] a strong wind will also make the flowers sway on the plant rendering access difficult for the pollinators. A wind force greater than six on the Beaufort scale interferes with bumble bee flight.

Besides affecting the nest sites, wind also had negative impact on the activity of *A. cerana* forager and the availability of pollen sources in high lands. The strong wind could change the flight direction of bees that they could not come back to their nests. The wind caused pollen to dry out and easily blown by wind. The disturbance on forager activity caused direct impact on the amount of stroge available in nests. The short supply of stroke caused the decrease on queen productivity. The limited amount of stroke in nests was one of the cause for colonies to abscond. The same results were reported by Woyke *et al.* [20] on *Apis dorsata* and *Apis laborisa* honey bees in India, Nepal and Bhutan.

Interaction of weather factors in low land polyculture plantations determined density population colony *A. cerana* as 54,97% (Table 1 and Equation 3). Besides weather factors other factor affecting colony population density of *A. cerana* was pest such as wax moths *G. mellonella* and ants and human activity. Three colonies were attacked by pest, wax moths, *Galeria mellonella* and one colony was attacked by ants that caused the colony to abscond. According to Gulati and Kaushik [26] among several limiting factors, honeybee enemies constitute a major factor. Wax moths and wasps cause heavy losses to beekeepers throughout the world. Ellis *et al.* [32] reported wax moths remain a vexing problem for beekeepers and honey bee colonies around the globe. According to Gulati and Kaushik [26] ants are not usually serious pests in honeybee colonies. Ants are typically found between the inner and outer covers of the hive and in pollen traps. The small red household ant, *Dorylus labiatus* and small black ants, *Monomorium indicum*, *M. destructor* are some of the other ant species which visit bee colony for food purposes.

#### 5. Conclusion

Eighteen plant species (15 families) and four other nonplant sites were found to be used by *A. cerana* to place their nests in West Sumatera. Colony population density of *A. cerana* fluctuated throughout the year with an average was  $4,06 \pm 0,65$  colonies per  $10.000 \text{ m}^2$ . The highest colony population density was found in June and lowest was in December and

January. Interaction of weather factors determined colony population density 95.03% in highlands and 54.97% in lowlands

## 6. Acknowledgement

We acknowledge Prof. Dr. Mansyurdin, M.S, and Dr. Henny Herwina, M.Sc, Biology Dept. Faculty of mathematic and Natural Science Andalas University for his great contribution to this paper. We also acknowledge the society of Andaleh and Parik Malintang for their information on the existence of colonies of *A. cerana*.

## References

- [1] F.Ruttner. *Bioqography and taxonomi of honeybees*. Springer-Verlag. Berlin.1988.
- [2] Munaan. The rule of honeybees on the pollination of coconut mixed-cropping. *Indonesian Agricultural Research & development Journal* 3 (3): 43-49. 1997.
- [3] M.Klein, I.Steffan-Dewenter, T.Tscharntke. Bee pollination and fruit set of *Coffea arabica* and *C. canephora* (Rubiaceae). *Ame J Bot* 90 (1): 153-157.2003.
- [4] Kuntadi. Pengaruh umur larva terhadap kualitas ratu yang dihasilkan pada penangkaran lebah ratu *Apis cerana* L. (Hymenoptera: Apidae) dengan teknik pencangkakan. *J Entomol Indonesia* 10 (1): 1-6.2013.
- [5] D. P. Abrol. Defensive behaviour of *Apis cerana* F. against predatory wasps. *J Apic Sci* 50 (2): 39-46. 2006.
- [6] K. Tan, Z. Wang, H.Li, S. Yang, Z. Hub, G. Kastberger, B. P. Oldroyd. An 'I see you' prey-predator signal between the Asian honeybee, *Apis cerana*, and the hornet, *Vespa velutina*. *Anim Behav* 30: 1-4. 2012.
- [7] S. Salmah, T.Inoue, and S.F.Sakagami. An analysis of aphid bee richness (Apidae) in Central Sumatra. In: *Natural History of Social Waps and Bees in Equatorial Sumatra*. Hokaido University Press. pp. 139-174.1990.
- [8] T. Inoue, Adri and S.Salmah. Nest site selection and reproductive ecology of the Asian honey bee, *Apis cerana indica* in central Sumatra. In: *Natural History of Social Waps and Bees in Equatorial Sumatra*. Hokaido University Press. pp. 219-2.1990.
- [9] M. R.Khan and M. R.Khan. The role honey bees *Apis mellifera* L. (Hymenoptera: Apidae) in pollination apple. *Pak J Bio Sci* 7 (3): 359-362.2004.
- [10] J. K.Tuell, J.S.Ascher and R. Isaacs. Wild bees (Hymenoptera: Apoidea: Anthophila) of the Michigan highbush blueberry agroecosystem. *Ann. Entomol. Soc. Am.* 102 (2): 275-287.2009.
- [11] F. N.Emuh and A. U.Ofuoku. Productivity of honeybees in oil palm integrated system in Niger-Delta of Nigeria. *J Agric and Environ* 12: 38-40.2011.
- [12] K. A.Baum, W. L.Rubink, M. A.Pinto and R. N.Coulson. Spatial and temporal distribution and nest site characteristics of feral honey bee (Hymenoptera: Apidae) colonies in a Coastal Praire Landscape. *Pop Eco* 34 (3): 610-618.2005.
- [13] R. N.Coulson, A. M.Pinto, M. D.Tchakerian, K. A.Baum, W. L.Rubink and J. S.Johnston. Feral honey bees in pine forest landscapes of east Texas. *For Eco and Mana* 215: 91-102. 2005.
- [14] S.Sheffield, P. G.Kevan, S. M.Westby and R. F.Smith. Diversity of cavity-nesting bees (Hymenoptera: Apoidea) within apple orchards and wild habitats in the Annapolis Valley, Nova Scotia, Canada. *Can Ento* 140(2):235-249. 2008.
- [15] T. E.Murray, M.Kuhlmann and S. G.Potts. Conservation ecology of bees: populations, species and communities. *Apidologie* 40: 211-236.2009.
- [16] D.Vaudo, J. D.Ellis, G. A.Cambray and M.Hill. The effects of land use on honey bee (*Apis mellifera*) population density and colony strength parameters in the Eastern Cape, South Africa. *J Insect Conserv*. 1-11.2011.
- [17] T.E. Rinderer, B. P. Oldroyd, L. I. de Guzman, W. Wattanachaiyingchareon and S.Wongsiri. Spatial distribution of the dwarf honey bees in an agroecosystem in Southeastern Thailand. *Apidologie*. 33: 539-543.2002.
- [18] N. H.Fefferman and P.T. Starks. A modeling approach to swarming in honey bees (*Apis mellifera*). *Insectes Sociaux* 53: 37-45.2006.
- [19] S. Kahono, T. Nakamura and M. Amir. Seasonal migration and colony behavior of the tropical *Apis dorsata* F. (Hymenoptera: Apidae). *Treubia* 31 (3): 285-299.2002.
- [20] J.Woyke, J.Wilde and M.Wilde. Swarming and migrating of *Apis dorsata* and *Apis laborisa* honey bees in India, Nepal and Bhutan. *J Apic Sci* 56 (1): 81-92.2012.
- [21] K.H. Bhuiyan, M.M.Hossain, M.N.Bari and M.R. Khan. Identification of bee plants and analysis of honey collected from different plant souerces. *Pakistan Journal of Biological Sciences* 5 (11): 1199-1201. 2002.
- [22] E.Mbah and A. O.Amao. Natural foods and feeding habits of the African honey bee *Apis mellifera adansonii* Latrielle (1804) in Zaria, Northern Nigeria. *Apidologie* 41: 264-277.2010.
- [23] A.Decourtye, E.Mader and N.Desneux. Landscape enhancement of floral resources for honey bees in agroecosystems. *Apidologie* 41: 264-277. 2010.
- [24] P.Tiwari, J. K. Tiwari and R. Ballabha. Studies on sources of bee-forage for rock bee (*Apis dorsata* F.) from Garhwal Himalaya, India: a Melissopalyno-gaical Approach. *Nature and Science* 8 (6): 5-15. 2010.
- [25] R.Shubharani, V.Sivaram and P. Roopa. Assessment of honey plant resources through pollen analysis in Coorg Honeys of Karnataka State. *Intern J. Plant Repro. Bio.* 4(1): 31-39. 2012.
- [26] R.Gulati and H. D.Kaushik. Enemies of honeybees and their management. *Agric. Rev.* 25 (3): 189-200.2004.
- [27] T. D. Selley. Honey bees of the Arnot Forest: a population of feral colonies persisting with Varroa destructor in the northeastern United States. *Apidologie* 38: 19-29.2007.
- [28] P.Abrol. Defensive behaviour of *Apis cerana* F. against predatory wasps. *J Apic Sci* 50 (2): 39-46.2006.

[29] N. S.M.Omran, M. H. Hussein, M. M. Khodairy and M. A.Awad. Predators of honeybee and its impact on activities of honeybee colonies under conditions of South Valley, Egypt. *J. Agri. and Bio. Sci.* 7(1): 79-88.2011.

[30] M.A. M.Ali and E. A.Taha. Bee-eating birds (Coraciiformes: Meropidae) reduce virgin honey bee queen survival during mating flights and foraging activity of honey bees (*Apis mellifera* L.). *Intern J Sci & Engi Res* 3 (6): 1-8.2012.

[31] J. D Ellis, J. R Graham and A.Mortensen. Standard methods for wax moth research. *J Apic Res* 52(1): 1-18.2013.

[32] Y.Le Conte and M.Navajas. Weather change: impact on honey bee populations and diseases. *Rev. Sci. Tech. Off. Int. Epiz* 27 (2): 499-510. 2008.

[33] BPS. *Statistik daerah West Sumatera tahun 2010*. Badan Pusat Statistik Provinsi West Sumatera. Padang.2010.

[34] R. G. D.Steel, and J.H. Torrie. *Principle and procedures of statistics*. McGraw-Hill, Inc. Gramedia Pustaka utama. Jakarta.1980.

[35] R. Raffiuddin, S.Hadisoesilo dan T.Atmowidi. Studi keragaman genetik dan morfologi lebah *Apis koschevnikovi* di Kalimantan Selatan. *Laporan penelitian*. Lembaga Penelitian dan Pemberdayaan Masyarakat. IPB, Bogor, Indonesia. 2005.

[36] M. C.Donaldson-Matasci and A.Dornhaus. How habitat affects the benefits of communication in collectively foraging honey bees. *Behav Ecol Socio* 1-10. 2011.

[37] S.Barfod, M.Hagen and F.Borchsenius. Twenty-five years of progress in understanding pollination mechanisms in palms (Arecaceae). *Ann Botany* 108: 1503-1516.2011.

[38] M.Chagnon. *Causes and effects of the worldwide decline in pollinators and corrective measures*. Canadian Wildlife Federation. Quebec Regional Office.2008.

[39] J. W.Rhodes, D. C.Somerville and H.Harden. Queen honey bee introduction and early survival-effects of queen age at introduction. *Apidologie* 35: 383-388.2004.

[40] M.Kjøhl, A. Nielsen and N.C. Stenseth. *Potential effects of weather change on crop pollinat.* Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biology, University of Oslo, Norway.2011.

[41] A.Arzone. Effect of pesticides on honeybees. Proceeding of the Seventh International Conference on Tropical Bees: *Management and Diversity and Fifth Asian apicultural Association Conference*. Chiang Mai, Thailand 19-25 March 2000.2000.

[42] C.Kremen, M.N. Williams, A. Marcelo, B. Gemmill-Herren, G. LeBuhn, R. Minckley, L. Packer, S. G. Potts, T. Roulston, I. Steffan-Dewenter, D. P. Va' zquez, R. Winfree, L. Adams, E. E. Crone, S. S. Greenleaf, T. HKeitt, A. M. Klein, J.Regetz and T. H.Ricketts., Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology Letters*. 10: 299-314. 2007.

a.Highland b.Lowland. M-M= times observation, started from March 2011 to March 2012.





Figure 2: Fluctuation of colony population density of *A. cerana* in polyculture plantations in West Sumatera

Table 1: Multiple regression analysis on colony population density and weather factors in polyculture plantations in West Sumatera

Variable	DB	Ft (0,05)	Ft (0,01)	Highlands	Lowlands
Regression	5	2,6	3,86	26,786	0,774
x1	1	4,24	7,77	4,473*	0,004 <sup>ns</sup>
x2	1	4,24	7,77	6,040*	2,504 <sup>ns</sup>
x3	1	4,24	7,77	1,253 <sup>ns</sup>	0,536 <sup>ns</sup>
x4	1	4,24	7,77	8,885**	0,021 <sup>ns</sup>
x5	1	4,24	7,77	4,170 <sup>ns</sup>	0,153 <sup>ns</sup>
Residu	7	-	-	-	-
Total	12	-	-	-	-
R	-	-	-	0,950	0,356

x<sub>1</sub>= rainfall (mmhg), x<sub>2</sub>= humidity (%), x<sub>3</sub>= temperatur (°C), x<sub>4</sub>= wind (knot), x<sub>5</sub>= photoperiode (%). <sup>ns</sup> = nonsignificant in 5% (α= 0,05) or 1% (α= 0,01). \* = significant difference 5% (α= 0,05) and \*\* = significant difference on level of confident 1% (α= 0,01).

Author Profile

Jasmi, M.S. Insect ecologist in Biology Department STKIP PGRI West Sumatera. Received B.S in Biology Dept. from Muhammadiyah University Padang Panjang in 1990 and taking Ph.D program at Andalas University, from 2009. He has become a lecturer since 1991 until now.

**Siti Salmah**, A Professor on Zoologi Taxonomy in Biology Dept. Faculty of Mathematic and Natural Science Andalas University, Padang

**Dahelmi**, A Professor on Entomology in Biology Dept. Faculty of Mathematic and Natural Science Andalas University, Padang.

**Syamsuardi**, A Professor on Plant Taxonomy in Biology Dept. Faculty of Mathematic and Natural Science Andalas University, Padang.

**Hidayani**, Lector on Department of Plant Protection, Agriculture Faculty, Andalas University, Padang.

