

# Prevalence of Hypertension and its Risk Factors among Urban Sikh Population of Amritsar

Amrinder Singh<sup>1</sup>, Shweta Shenoy<sup>2</sup>, Jaspal Singh Sandhu<sup>3</sup>

<sup>1</sup>Assistant Professor, Faculty of Sports Medicine & Physiotherapy, Guru Nanak Dev University, Amritsar, Punjab, India

<sup>2</sup>Associate Professor, Faculty of Sports Medicine & Physiotherapy, Guru Nanak Dev University, Amritsar, Punjab, India

<sup>3</sup>Dean, Faculty of Sports Medicine & Physiotherapy, Guru Nanak Dev University, Amritsar, Punjab, India

**Abstract:** *Hypertension is a major risk factor for cardiovascular disorders, stroke and myocardial infarction, and its burden is increasing disproportionately in developing countries as they undergo demographic transition. Hypertension is responsible for 57% of stroke deaths and 24% of coronary heart disease deaths in India. The sample size of 1089 subjects was calculated. This study focused on Urban Sikhs living in the Amritsar, Punjab, India as they constitute 70% of the total population of Amritsar according to 2011 census report. The prevalence of hypertension has increased during the last decade. The high prevalence of hypertension (35.9%) in this study, confirms this increasing trend. It is necessary to adopt appropriate preventive strategies and interventions in high risk individuals to curb the growing epidemic of hypertension.*

**Keywords:** Hypertension, Sikh, Urban, Cardiovascular, lifestyle, prevalence

## 1. Introduction

Hypertension is the most common cardiovascular disease and is assuming epidemic proportions in developing countries as well [1]. It affects nearly 26% of the population worldwide [2]. Hypertension exhibits an iceberg phenomenon where unknown morbidity exceeds the known morbidity. Hypertension is a major risk factor for cardiovascular disorders, stroke and myocardial infarction, and its burden are increasing disproportionately in developing countries as they undergo demographic transition [3, 4, 5, 6]. It is estimated that by 2020, CVDs will be the largest cause of mortality and morbidity in India [7]. The prevalence of hypertension is rapidly increasing in developing countries and is said to be one of the leading causes of death and disability among the elderly [8]. Hypertension is responsible for 57% of stroke deaths and 24% of coronary heart disease deaths in India. [9]

Using a cut-off of 140 mmHg or greater systolic blood pressure (BP), or 90 mmHg or greater diastolic BP, the age-standardized prevalence of hypertension worldwide in the year 2000 was estimated to be 26.6% in men (95% confidence interval, CI 26.0 to 27.2) and 26.1% in women (95% CI 25.5 to 26.6). This is estimated to rise to 29.0% in men (95% CI 28.6 to 29.4) and 29.5% in women (95% CI 29.1 to 29.9) by 2025 [10]

India, the world's largest democracy, is undergoing a rapid economic growth. This growth has been accompanied by demographic, lifestyle and cultural changes which have had a large impact on the health profile of India's citizens and placed a significant strain on the country's healthcare system [11, 12, 13]

It was estimated that around two-thirds of those with people with hypertension worldwide were living in developing countries (639 million) in 2000, and that this would rise to three-quarters living in developing countries (1.15 billion) by 2025. Previous authors have suggested prevalence rates

for hypertension in urban Indian population to be 29-45% in men and 25-38% in women [14, 15, 16, 17]

Indian studies have revealed that the prevalence of hypertension has increased by 30 times among the urban population over a period of 55 years and about ten times among the rural population over a period of 36 years. (18). Management of hypertension requires life-long medication with some lifestyle modifications. The only way to curb the problem of hypertension is by its prevention.

Several risk factors have been implicated in the etiology of hypertension. This includes geographic considerations, genetic socio-economic, socio-cultural and dietary, nutritional status etc. While the risk factors and their impact on hypertension is documented by well designed studies in the Western countries, systematically conducted studies using rigorous epidemiological techniques are lacking in India. Epidemiological studies to assess the prevalence of Hypertension are urgently needed in developing countries like India to determine the baseline against which future trends in risk factor levels can be assessed and preventive strategies planned to promote health. The urban population is very prone to development of cardiovascular diseases with Sidhu et al putting to figure at 20.15 % [19]. In spite of the prediction of this population to the development of CVD, there has been no Epidemiological in this population.

Thus the primary aim of this study was to identify the prevalence of hypertension in the Urban Sikh Population of Amritsar by means of a door-to-door survey. A secondary aim was to identify the risk factors for hypertension.

## 2. Materials and Methods

The sample size of 1089 subjects was calculated. The multi stage cluster randomized sampling was done using the probabilities proportional size (PPS) method. This study focused on Urban Sikhs living in the Amritsar, Punjab, India as they constitute 70% of the total population of Amritsar

according to 2011 census report. The area was arbitrarily divided into five equal zones namely (Z<sub>1</sub>, Z<sub>2</sub>, Z<sub>3</sub>, Z<sub>4</sub>, Z<sub>5</sub>). Each zone constituted of 13 Censes wards .The list provided the name, age and address of those eligible for voting (>18 years). The fieldwork was completed in a period of 20 months, starting in Jan 2012. The totals of 215 subjects from each zone were randomly selected. In the selection of the family in a particular ward the WHO method of sampling was followed to prevent any non uniformity in selection of the subjects. A model consent form was constructed to ensure compliance with ICMR guidelines regarding the use of human subjects in research. All protocols and consent documents were reviewed and approved by the Institutional Ethics Committee of Faculty of Sports Medicine & Physiotherapy, Guru Nanak Dev University, Amritsar. The complete detailed Performa of all the family members was filled by the investigator which included number of members in the family, age, educational status etc. From the list provided one male and one female member were randomly selected. These two family members were requested to fill in the self designed, professionally validated questionnaire Blood pressure was measured using a standard auscultatory method with the help of a pneumatically operated mercurial type random-zero sphygmomanometer in the sitting position with the back supported and the arm was kept at the level of fourth interspaced at sternum. Blood pressure was measured after the questionnaire was filled in i.e. nearly 5 min after the visit, by which time the emotional reaction of the subject had stabilized. Three casual readings were taken. Average of these readings was taken as the final reading.

Hypertension was defined on the basis of Joint National Committee (JNC) VII criteria. If blood pressure was > 120/80, the BP was checked again after minimum of 5 min. and only if BP was >140/90, was the person labeled to have HTN. Persons having systolic blood pressure (SBP) between 120-139 and / or diastolic blood pressure (DBP) between 80 - 89 were labeled to have pre-hypertension. Stage1 hypertension was taken as systolic BP between 140-159 and/ or diastolic BP between 90-99 mmHg. Stage 2 hypertension was taken as systolic BP > 160 and/ or diastolic BP > 100 mmHg. The fasting blood sample was taken for the analysis of biochemical variables. The blood was analyzed for serum cholesterol, (CHO), serum triglyerides (TG), (High density lipoprotein (HDL),(Low density lipoprotein (LDL), very low density lipoprotein (VLDL).The anthropometric readings were taken namely height, weight, waist (WC)and hip circumference(HC).

**Results:** All the Statistical analysis was done using Stata 11.2. The association of the categorical variables with outcomes was seen by the Chi Square test. p values less than 0.05 were taken as significant. Unadjusted odds ratio of relationship of each factor with outcomes was found using binary logistic regression test. Factors, which are significant in univariate analysis, were included in multivariate analysis using multi variable logistic regression test.

**Table 1:** Prevalence of hypertension in the urban sikh population with 95% CI

| Lifestyle disorder | Total Subject population | Frequency | Prevalence in %age ( 95% CI) |
|--------------------|--------------------------|-----------|------------------------------|
| Hypertension       | 1089                     | 359       | 35.9(33.0,38.7)              |

**Table 2:** Gender wise distribution of the biochemical parameters in hypertensive and non hypertensive individuals (Mean ± SD)

| Age              | Male         | Female       |
|------------------|--------------|--------------|
| Non hypertensive | 45.6 ± 14.3  | 46.1 ± 14.7  |
| hypertensive     | 55.1 ± 13.5  | 54.3 ±13.1   |
| CHO              |              |              |
| Non hypertensive | 199.3 ±46.9  | 193.2 ±42.1  |
| hypertensive     | 205.2 ±44.0  | 209.7 ±50.5  |
| TG               |              |              |
| Non hypertensive | 156.4 ±65.3  | 154.2 ±61.0  |
| hypertensive     | 167.9 ± 55.3 | 165.2 ± 60.9 |
| HDL              |              |              |
| Non hypertensive | 44.2 ± 10.3  | 44.0 ±9.7    |
| hypertensive     | 47.9 ± 8.5   | 44.2 ±11.0   |
| LDL              |              |              |
| Non hypertensive | 147.2 ±49.4  | 140.4 ±42.1  |
| hypertensive     | 154.0 ± 46.2 | 154.9 ±43.4  |
| VLDL             |              |              |
| Non hypertensive | 33.0 ± 23.8  | 31.3 ± 18.1  |
| hypertensive     | 33.4 ± 11.1  | 33.0 ± 12.3  |
| SBP              |              |              |
| Non hypertensive | 120.0 ±8.4   | 119.7 ±7.9   |
| hypertensive     | 136.4 ±11.6  | 139.4 ±13.0  |
| DBP              |              |              |
| Non hypertensive | 80.0 ±5.5    | 78.8 ± 5.3   |
| hypertensive     | 93.2 ± 5.8   | 94.1 ± 6.8   |
| BMI              |              |              |
| Non hypertensive | 26.1 ± 5.0   | 26.1 ± 5.2   |
| hypertensive     | 25.4 ± 5.0   | 26.1 ± 5.4   |
| WC               |              |              |
| Non hypertensive | 84.9 ± 13.0  | 84.4 ± 13.5  |
| hypertensive     | 80.8 ± 12.4  | 84.0 ± 13.5  |
| HC               |              |              |
| Non hypertensive | 97.6 ± 14.4  | 98.1 ± 13.0  |
| hypertensive     | 98.1 ± 13.4  | 97.4 ± 13.9  |
| WHR              |              |              |
| Non hypertensive | 0.89 ± 0.21  | 0.87 ± 0.19  |
| hypertensive     | 0.83 ± 0.18  | 0.87 ± 0.20  |

**Table 3:** Age Specific &Age Standardized Prevalence of Hypertension gender- wise in % age with 95% CI

| Age Group (in years) | Prevalence of Hypertension in % age with 95% CI |                                 |        |                                 |       |                                 |
|----------------------|---|---------------------------------|--------|---------------------------------|-------|---------------------------------|
|                      | Male  |                                 | Female |                                 | Total |                                 |
|                      | n   | Prevalence in % age with 95% CI | n      | Prevalence in % age with 95% CI | n     | Prevalence in % age with 95% CI |
| 20-29                | 30  | 10.0 (-1.4,21.4)                | 45     | 11.1 (1.6,20.6)                 | 75    | 10.7 (3.5,17.8)                 |
| 30-39                | 96  | 25.0 (16.2,33.9)                | 96     | 14.6 (7.4,21.8)                 | 192   | 19.8 (14.1,25.5)                |
| 40-49                | 136   | 39.0 (31.5,47.3)                | 127    | 25.9 (18.2,33.7)                | 263   | 32.7 (27.0,38.4)                |
| 50-59                | 147   | 39.4 (31.5,47.4)                | 104    | 43.3 (33.6,52.9)                | 251   | 41.0 (34.9,47.2)                |
| 60+                  | 178   | 54.5 (47.1,61.9)                | 130    | 45.4 (36.8,54.0)                | 308   | 50.6 (45.0,56.3)                |
| Total                | 587   | 40.0 (36.0,44.0)                | 502    | 31.0 (27.0,35.1)                | 1089  | 35.9 (33.0,38.7)                |

**Table 3:** Risk factors for Hypertension

| Variables                            | Hypertension              |                       | p-value | Unadjusted Odds Ratio with 95% CI | Adjusted Odds Ratio with 95% CI |
|--------------------------------------|---------------------------|-----------------------|---------|-----------------------------------|---------------------------------|
|                                      | Non hypertensive n (%age) | hypertensive n (%age) |         |                                   |                                 |
| Age (in years)                       |                           |                       |         |                                   |                                 |
| 20-29                                | 67 (9.6)                  | 8.0 (2.0)             | <0.00   | 1                                 | 1                               |
| 30-39                                | 154 (22.0)                | 38 (9.7)              |         | 2.0 (0.9,4.7)                     | 1.9 (0.8,4.4)                   |
| 40-49                                | 177 (25.3)                | 86 (21.9)             |         | 4.0 (1.9,8.9)                     | 3.5 (1.6,7.7)                   |
| 50-59                                | 148 (21.2)                | 103 (26.3)            |         | 5.8 (2.7,12.7)                    | 4.7 (2.1,10.4)                  |
| 60+                                  | 152 (21.7)                | 156 (39.9)            |         | 8.6 (4.0,18.5)                    | 6.9 (3.1,15.2)                  |
| Gender                               |                           |                       |         |                                   |                                 |
| Female                               | 346(49.5)                 | 156 (39.9)            | <0.00   | 1                                 | 1                               |
| Male                                 | 352 (50.4)                | 235 (60.1)            |         | 1.5 (1.1,1.9)                     | 1.0 (0.8,1.5)                   |
| BMI                                  |                           |                       |         |                                   |                                 |
| Underweight                          | 46 (6.5)                  | 27 (6.9)              | 0.48    | 1.2 (0.7,2.0)                     | -----                           |
| Normal                               | 239 (34.2)                | 116 (29.6)            |         | 1                                 | -----                           |
| Pre Obese                            | 251 (35.9)                | 149 (38.1)            |         | 1.2 (0.9,1.6)                     | -----                           |
| Obese                                | 162 (23.2)                | 99 (25.3)             |         | 1.2 (0.9,1.7)                     | -----                           |
| Waist Circumference                  |                           |                       |         |                                   |                                 |
| < 102 cm in males < 88 cm in females | 505(72.3)                 | 293(74.9)             | 0.35    | 1                                 | -----                           |
| ≥ 102 cm in males ≥ 88 cm in females | 193 (27.6)                | 98 (25.0)             |         | 0.8 (0.6,1.2)                     | -----                           |
| Physical Activity                    |                           |                       |         |                                   |                                 |
| Mild                                 | 29 (4.1)                  | 16 (4.0)              | 0.7     | 1                                 | -----                           |
| Moderate                             | 83 (11.8)                 | 40 (10.2)             |         | 0.9 (0.4,1.8)                     | -----                           |
| Heavy                                | 586 (83.9)                | 335 (85.6)            |         | 1.0 (0.5,2.0)                     | -----                           |
| Diet                                 |                           |                       |         |                                   |                                 |
| Vegetarian                           | 378 (54.1)                | 198 (50.6)            | 0.26    | 1                                 | -----                           |
| Non Vegetarian                       | 320 (45.8)                | 193 (49.3)            |         | 1.1 (0.9,1.5)                     | -----                           |
| Alcohol Intake                       |                           |                       |         |                                   |                                 |
| Non Alcoholic                        | 567 (81.2)                | 293 (74.9)            | 0.01    | 1                                 | 1                               |
| Alcoholic                            | 131 (18.7)                | 98 (25.0)             |         | 1.4 (1.0,1.9)                     | 1.3 (0.9,1.8)                   |
| Family History                       |                           |                       |         |                                   |                                 |

|   |            |             |       |               |               |
|---|------------|-------------|-------|---------------|---------------|
| No family history                           | 396 (56.7) | 209 (53.4)  | 0.46  | 1             | -----         |
| Family history of T2DM                      | 162 (23.2) | 92 (23.5)   |       | 1.0 (0.8,1.5) | -----         |
| Family history of hypertension              | 140 (20.0) | 90 (23.0)   |       | 1.2 (0.9,1.7) | -----         |
| Type of oil used                            |            |             |       |               |               |
| Refined                                     | 494 (70.7) | 267 (68.2)  | 0.22  | 1             | -----         |
| Desi Ghee                                   | 124 (17.7) | 65 (16.6)   |       | 1.0 (0.7,1.3) | -----         |
| Dalda                                       | 80 (11.4)  | 59 (15.0)   |       | 1.3 (1.0,2.0) | -----         |
| Triglycerides                               |            |             |       |               |               |
| <150 mg/dl                                  | 341 (48.8) | 144 (36.8)  | <0.00 | 1             | 1             |
| ≥ 150 mg/dl                                 | 357 (51.1) | 247 (63.1)  |       | 1.6 (1.3,2.1) | 1.2 (0.9,1.6) |
| HDL   |            |             |       |               |               |
| < 40 mg/dl for males < 50 mg/dl for females | 401 (57.4) | 199(50.9)   | 0.03  | 1             | 1             |
| ≥ 40 mg/dl for males ≥50 mg/dl for females  | 297 (42.5) | 192 (49.1)  |       | 0.8 (0.6,1.0) | 0.8 (0.6,1.0) |
| Cholesterol                                 |            |             |       |               |               |
| < 200 mg/dl                                 | 402 (57.5) | 173 (44.2)  | <0.00 | 1             | 1             |
| ≥ 200 mg/dl                                 | 296 (42.4) | 218 (55.7)  |       | 1.7 (1.3,2.1) | 1.2 (0.9,1.7) |
| LDL   |            |             |       |               |               |
| <130 mg/dl                                  | 251 (35.9) | 108 (27.6)  | <0.00 | 1             | 1             |
| ≥ 130 mg/dl                                 | 447 (64.0) | 283 (72.3)  |       | 1.5 (1.1,1.9) | 0.9 (0.7,1.3) |
| Diabetes                                    |            |             |       |               |               |
| Not present                                 | 563 (80.6) | 273 (69.8)  | <0.00 | 1             | 1             |
| Present                                     | 135 (19.3) | 118 (30.21) |       | 1.8 (1.3,2.4) | 1.4 (1.0,1.9) |
| Metabolic Syndrome                          |            |             |       |               |               |
| Not present                                 | 549 (78.6) | 166 (42.4)  | <0.00 | 1             | -----         |
| Present                                     | 149 (21.3) | 225(57.5)   |       | 5.0 (3.8,6.5) | -----         |

Table 5: BMI: WHR distribution in the sample population

| BMI         | WHR         | N   |
|-------------|-------------|-----|
| Normal      | 0.79 ± 0.15 | 355 |
| Underweight | 0.71 ± 0.2  | 73  |
| Pre Obese   | 0.91 ± 0.18 | 400 |
| Obese       | 1.01 ± 0.2  | 261 |


**Table 6:** Mean & SD values of the anthropometric physiological and biochemical parameters accordingly to age group

| Age (in years) | BMI        | Waist       | WHR         | TG           | CHO          |
|----------------|------------|-------------|-------------|--------------|--------------|
| 0-40           | 26.4 ± 5.3 | 85.7 ± 13.5 | 0.89 ± 0.19 | 150.7 ± 66.1 | 188.8 ± 46.2 |
| >40            | 26.3 ± 5.3 | 84.9 ± 13.7 | 0.88 ± 0.20 | 173.3 ± 66.5 | 210.8 ± 47.8 |

| Age (in years) | HDL         | LDL          | VLDL        | SBP          | DBP        |
|----------------|-------------|--------------|-------------|--------------|------------|
| 0-40           | 44.4 ± 10.3 | 135.4 ± 42.6 | 31.1 ± 11.5 | 121.3 ± 11.5 | 80.5 ± 8.1 |
| >40            | 43.4 ± 9.3  | 156.8 ± 14.6 | 34.8 ± 14.6 | 129.2 ± 13.2 | 85.7 ± 9.0 |

**Table 7:** Blood pressure classification in the subject population

| Blood pressure  | N   | % age |
|---|-----|-------|
| Normal (SBP ≤ 119, DBP ≤ 79 mm of Hg)                 | 133 | 12.2  |
| Pre Hypertensive (SBP 120-139 DBP 80-89 mm of Hg)     | 565 | 51.8  |
| Hypertension Stage 1 (SBP 140-159, DBP 90-99mm of Hg) | 276 | 25.3  |
| Hypertension Stage 2 (SBP ≥ 160, DBP ≥ 100mm of Hg)   | 115 | 10.6  |



**Figure 1:** Age wise Trend of Hypertension

Demographic & Clinical profile of subjects with and without T2DM. Table 3 shows the gender wise prevalence of hypertension across different age groups. The study also shows significantly higher rates of hypertension in the older age groups. The hypertension prevalence rates increase from 10 % in the age group of 20-29 years peaking to 50.6% in the age group of 60 + years.

Chi Square outputs of significant proportions with several clinical, demographic and biochemical parameters available to the present study comparing the subjects with and without hypertension are shown in Table 4. Proportionately more subjects with hypertension had diabetes (30.2%). Likewise proportionately more subjects, 63.1 % had hypertriglyceridemia, 50.9 % had low HDL levels, 55.7 % had hypercholesterolemia and 57.5% suffered from metabolic syndrome. Gender wise comparison shows that out of total hypertensive's 60 % are male's patients. 23% hypertensive's had a positive family history of hypertension. 63.4 % of the hypertensive's fall in category of pre-obese and obese category.

Detailed correlates of hypertension in univariate analysis and multivariate logistic regression method showing

significant predictors of hypertension are summarized in Table 4. Advancing age, gender, alcohol intake, T2DM, hypercholesterolemia, hypertriglyceridemia and low serum HDL levels significantly contributed to the increased hypertension risk. The increased serum cholesterol and serum triglycerides increased the risk of developing hypertension one and half times more.

### 3. Discussion

The prevalence of hypertension has increased during the last decade [20]. The high prevalence of hypertension (35.9%) in this study, confirms this increasing trend. Rapid urbanization, lifestyle changes are dietary changes are the factors attributable to this rising trend. Increased prevalence of hypertension observed in this study was similar to that reported in Himachal Pradesh with an overall prevalence of 35.89 % with 34.8% in males and 33.1 % in females. [21], in Central India, overall prevalence of hypertension is 19.04%, with 23.4% males and 14.4 % in females. (22) Similarly in Kerala overall prevalence was 36.7% with 36 % in males and 37.2 % in females [23].

Further we also observed that the findings of the current study are coherent with those reported in the study conducted among urban and rural adults of Lucknow [24]. Cross-sectional surveys, as well as prospective observational cohort studies, have consistently demonstrated a positive relation between age and blood pressure in most populations with diverse geographical, cultural and socioeconomic characteristics. Such changes of blood pressure with age might be due to changes in vascular system [25]

When the age- and sex-specific prevalence of hypertension was compared, The proportion of hypertension was higher among males (60.1%) compared to that in females (39.9 %). It is also seen that in each age group, the prevalence of hypertension was higher among males as compared to females except in 20-29 and 50-59 years of age group. This concurs with the observation made by a previous study that males (42.9%) had high risk of hypertension when compared to females (34.2%) among rural population of Davanagere [26]. This fact also confirm with the current study that women have lower systolic blood pressure (SBP) levels than men during early adulthood, while the opposite is true after the sixth decade of life [27]. Diastolic blood pressure (DBP) tends to be just marginally lower in women than men regardless of age. Similarly, in early adulthood, hypertension is less common among women than men. However, the incidence of hypertension increases more rapidly in women older than 60 years. The highest prevalence rates of hypertension were observed in elderly black women (>75%), aged > 75 years [27].

In the present study, significant positive association was found between age and prevalence of hypertension. In the target population of 20 years and above, the prevalence of T2DM increased with age from 10.7% in 20-29 years old age group through 19.8% in those aged 30-39 years, 32.7 % in age group 40 -49 years, 41 % in 50-59 year old age group and 50.6 % in those aged ≥ 60 years. Desai *et al* [28] also reported an increase in the prevalence of hypertension with the increasing age.



In this study we inferred that 30.2 % of the hypertensive's suffered from Type 2 Diabetes Mellitus. Type 2 Diabetes mellitus is considered as an important risk factor in the development of hypertension, i.e. subjects who suffer from diabetes had 1.4 times more risk of developing hypertension. The prevalence of coexistent hypertension and diabetes varies across different ethnic, racial, and social groups. Importantly, hypertension in patients with type 2 diabetes causes a significant increase in the risk of vascular complications in this population, and together both conditions predispose to chronic kidney disease. [29,30]. The overlap between hypertension and diabetes substantially increases the risk of ischemic cerebrovascular disease, retinopathy, and sexual dysfunction. [31]. Diabetes mellitus is an independent risk factor for coronary artery disease, and the risk is markedly increased when hypertension is present.

Another risk factor in development of hypertension is obesity. Very few subjects in this study were in the category of healthy BMI. Only 32.6 % (n= 355) of the subjects presented with a healthy BMI as per the WHO guidelines. 6.7% (n= 73) were reported underweight, whereas 36.7% (n= 400) were reported as pre obese and 24% (n=261) presented as obese. The subject population presented a picture of general obesity as central obesity was not significantly present in this population. General obesity is related to the abnormal lipid profile values i.e increased serum cholesterol, serum triglyride, LDL and decreased HDL values.

While observing the life style pattern of Sikh population in current study we found that 25 % of the hypertensive's in this population were alcoholics. Alcohol intake also presents as independent risk factor in the development of hypertension. A person who consumes alcohol has 1.3 times more risk of having hypertension in some part of his lifetime than the non alcoholic. The alcohol intake raises the risk of hypertension and the cardiovascular disorders and stroke.

Our study also clearly indicates that the Young Sikh adults < 40 years of age have similar high BMI, WC and WHR to that of the older adults >40 years of age (Table 6). It is necessary to adopt appropriate preventive strategies and interventions in high risk individuals to curb the growing epidemic of hypertension.

## References

- [1] Gupta R. Rethinking Diseases of Affluence; Coronary Heart Disease in Developing Countries. *South Asian Journal of Preventive Cardiology* 2006; 10(2): 65-78 and 2004; 8(1&2): 5-6.
- [2] Kearney PM, Whelton M, He J. Global Burden of Hypertension: analysis of worldwide data. *Lancet* 2005; 365:217-23
- [3] O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, *et al.* Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): A case-control study. *Lancet* 2010;376:112-23
- [4] Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, *et al.* Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): Case-control study. *Lancet* 2004;364:937-52
- [5] Murray CJL, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global burden of disease study. *Lancet* 1997; 349:1498-504.
- [6] Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: Analysis of worldwide data. *Lancet* 2005; 365:217-23.
- [7] World Health Organization. The World Health Report 2002: Reducing Risk, Promoting Healthy Life. Geneva, Switzerland: World Health Organization; 2002
- [8] Murray C L J, Lopez A D: Global Health statistics: Global burden of diseases and injury series. *Harvard school of Public Health: Boston* 1996;349:1436-42
- [9] Gupta R. Trends in hypertension epidemiology in India. *Journal of Human Hypertension* 2004; 18:73-78.
- [10] Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: Analysis of worldwide data. *Lancet* 2005; 365:217-23.
- [11] Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: Part 1: General considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* 2001;104:2746-53
- [12] World Health Organization. The World Health Report 2008: Primary Health Care - Now More than Ever. Geneva, Switzerland: World Health Organization; 2008
- [13] Horton R, Das P. Indian health: The path from crisis to progress. *Lancet* 2011; 377:181-3.
- [14] Gupta R, Gupta VP. Hypertension epidemiology in India: Lessons from Jaipur Heart Watch. *Curr Sci* 2009;97:349-55
- [15] Gupta R, Sharma AK, Gupta VP, Bhatnagar S, Rastogi S, Deedwania PC. Increased variance in blood pressure distribution and changing hypertension prevalence in an urban Indian population. *J Hum Hypertens* 2003;17:535-40
- [16] Reddy KS, Prabhakaran D, Chaturvedi V, Jeemon P, Thankappan KR, Ramakrishnan L, *et al.* Methods for establishing a surveillance system for cardiovascular diseases in Indian industrial populations. *Bull World Health Organ* 2006;84:461-9
- [17] Hypertension Study Group. Prevalence, awareness, treatment and control of hypertension among the elderly in Bangladesh and India: A multicentre study. *Bull World Health Organ* 2001;79:490-500
- [18] Gupta R. Meta analysis of prevalence of hypertension in India. *Indian Heart J* 1997;49:43-8
- [19] S. Sidhu, K. Kumari and Prabhjot, Socio-demographic Variables of Hypertension among Adult Punjabi Females, *J. Hum. Ecol.*, 17(3): 211-215 (2005)
- [20] Padmavati S. A meta-analysis-National Heart Institute, New Delhi. *Ind Heart J* 2002; 54:99-102
- [21] Bhardwaj R, Kandoria A, Marwah R, Vaidya P, Singh B, Dhiman P, Sharma A. Prevalence, Awareness and Control of Hypertension in Rural Communities of Himachal Pradesh JAPI 2010;58:423-25.
- [22] Kokiwar PR, Gupta SS, Durge PM. Prevalence of Hypertension in a Rural Community of Central India. *J A P I* 2012;60:26-29
- [23] Thankappan KR, Sivasankaran S, Khader SA. Prevalence, awareness, treatment and control of in Hypertension, Kumarakom, Kerala. *Indian Heart Journal* 2006: 58:28-33

- [24] Midha T, Idris MZ, Saran RK, Srivastav AK, Singh SK. Prevalence and determinants of hypertension in the urban and rural population of a north Indian district. *East Afr J Public Health*. 2009 Dec;6(3):268-73
- [25] Hypertension control. Technical Report Series: World Health Organization; 1996. Report No. 862.
- [26] Yuvaraj BY, Nagendra Gowda MR, 1 and Umakantha AG. Prevalence, Awareness, Treatment, and Control of Hypertension in Rural Areas of Davanagere. *Indian J Community Med*. 2010 January; 35(1): 138–141
- [27] Priscilla Igho Pemu, and Elizabeth Ofili, Hypertension Women: Part I, *J Clin Hypertens (Greenwich)*. May 2008; 10(5): 406–410.
- [28] Kumar P, Desai VK, Kosambia JK. Prevalence of hypertension amongst the employees of a mega-industry of South Gujarat. *Indian J Community Med* 1995;27:19-25.
- [29] Wannamethee SG et al. (2005) Metabolic syndrome vs Framingham Risk Score for prediction of coronary heart disease, stroke, and type 2 diabetes mellitus. *Arch Intern Med* 165: 2644–2650
- [30] Adler AI et al. (2000) Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ* 321: 412–419
- [31] Najarian RM et al. (2006) Metabolic syndrome compared with type 2 diabetes mellitus as a risk factor for stroke: the Framingham Offspring Study. *Arch Intern Med* 166: 106–111