

Chronic Tonsillitis and its Relation with Childhood Asthma

Rameez Shah¹, Md. Bahauddin², Md.Rafiqul Islam³, Md.Shahriar Islam⁴, S.M.Sadlee⁵, Mostafa Kamal Arefin⁶

^{1,4,6}Department of Otolaryngology & Head-Neck Surgery, Dhaka Medical College Hospital, Dhaka, Bangladesh

²ENT & Head-Neck Cancer Hospital Institute, Dhaka, Bangladesh

³Diabetic Association Medical College Hospital, Faridpur, Bangladesh

⁵Department of Internal Medicine, Dhaka Medical College Hospital, Dhaka, Bangladesh

Abstract: ***Objective:** Chronic tonsillitis and childhood asthma, both are common childhood diseases, and it's not surprising to think of an association between the two by otolaryngologists treating chronic tonsillitis in children. **Methods:** A total of 500 children, aged 4–17 years, either reporting to OPD, Dept of ENT, or admitted in the Dept of Otolaryngology, during the period January'2011 and January'2013 were surveyed using the ISAAC questionnaire; and tonsils were evaluated by physical examination. **Results:** A total of 446 questionnaires were completed giving an overall response rate of 89.2%. The prevalence of lifetime wheezing, 12-month wheezing and lifetime doctor-diagnosed asthma were 16.4%, 6.2% and 1.7%, respectively. The prevalence of frequent tonsillitis and tonsillar hypertrophy were 19.9% and 3.4%, respectively. Tonsillar hypertrophy was positively correlated with a history of frequent tonsillitis. **Conclusion:** - Frequent tonsillitis and tonsillar hypertrophy could be evaluated as risk factors for asthma due to their significant association with asthma-related symptoms. And thus, it is expected that this study would be a suitable baseline source for future trends on the prevalence, diagnoses and severity of asthma among 4–17 year old children.*

Keywords: Chronic tonsillitis; childhood asthma;

1. Introduction

Chronic tonsillitis is one of the most common disease entities encountered by otolaryngologists and is frequently associated with upper airway infections [1] including nasopharyngitis, pharyngitis, tonsillitis and otitis media constituting 87.5% of the total episodes of respiratory infections. Also the majority of acute upper respiratory tract infections are caused by viruses. [1], [2]

Asthma is also one of the most common chronic diseases in the pediatric population. [3] An estimated 5- 15% of children have asthma, and there are worldwide variations in the prevalence of the different symptoms of asthma.[4] The prevalence of asthma is increasing in western and developing countries.[5], [6]

The majority of acute asthma episodes are triggered by respiratory viral infections in which recruitment of Th2-type cells into the lungs might be responsible. [7] The frequency of upper respiratory tract infections was reported to be higher in asthmatic children than in controls. [8]

The tonsils are part of Waldeyer's ring, the basic function of which is antibody formation and defense against antigens. Allergic sensitization of the airways occurs partly in the mucosa of the afflicted organ, but also partly in the lymphatic stations draining these structures. Adenoids and tonsils are the relevant lymphatic stations of the nasal mucosa in humans. Allergy and sensitivity to different kinds of allergens are thought to be risk factors for adenoid hypertrophy (AH) in children. [9]

There are only a few studies on the relationship between

asthma symptoms and tonsillar tissue in the literature. In this study, we aimed to evaluate the relationship between chronic tonsillitis and childhood asthma among 4–17 year old children.

2. Material and Methods

A total of 500 children, aged 4–17 years, either reporting to OPD, Dept of Otolaryngology, or admitted in the *Dept of Otolaryngology*, during the period *January'2011 and January'2013* were surveyed using the ISAAC (International Study of Asthma and Allergies in Childhood) questionnaire [4], [10] Children from each class level were included to provide an equal distribution of children according to age.

The questionnaire included questions on the history of frequent tonsillitis, tonsillectomy and/or adenoidectomy and symptoms of asthma. A detailed form was completed for each child by the parents/attendants. Thereafter, the tonsils were evaluated by otolaryngologists in all children and scored on a four-point scale as follows:

Grade 1, tonsils were in the tonsillar fossa, barely seen behind the anterior pillars;

Grade 2, tonsils were visible behind the anterior pillars;

Grade 3, tonsils extended three-quarters of the way to the midline;

Grade 4, tonsils were completely obstructing the airway.[11-13]

Grades 3 and 4 were accepted as tonsillar hypertrophy. The interrelationship between tonsillar hypertrophy and asthma-

related symptoms was examined by the chi-squared test, and the 95% confidence interval (CI) was calculated for the differences where appropriate. Those with tonsillectomy and/or adenoidectomy were excluded. Frequent tonsillitis was defined as more than two episodes of tonsillitis per year. The interrelationships between a history of frequent tonsillitis, tonsillectomy and/or adenoidectomy and asthma-related symptoms were examined by the chi-squared test. A significant difference was indicated by a p -value < 0.05. Odds ratios and 95% confidence limits for risk factors were calculated with coefficients and standard errors.

3. Results

Of the 500 children, 54 were excluded due to refusal. A total of 446 questionnaires were completed with an overall 89.2% response rate. Of 446 children, 201 (45%) were boys and 245 (55%) were girls. The male to female ratio was 0.82. Demo-graphic data of the study population are given in Table 1.

Table 1: Demographic data of the study population

Number of the study population	446
Sex, n (%)	
Male	201 (45)
Female	245 (55)
Age, median years (range)	11(4-17)
Children with frequent tonsillitis, n (%)	89(19.9)
Children with tonsillar hypertrophy, n (%)	15 (3.4)
Children with tonsillectomy and/or adenoidectomy, n (%)	24 (5.5)
Children with adenoidectomy, n (%)	10 (2.2)

In this study, the prevalence of lifetime wheezing, wheezing in the previous year, and life time doctor diagnosed asthma were 16.4%, 6.2%, and 1.7%, respectively. The prevalence of other asthma-related symptoms is given in Table 2.

Table 2: Prevalence of asthma and other symptoms

Question	N (%)
Lifetime wheezing	73 (16.4)
Wheezing in last year	28 (6.2)
Attacks of wheezing in last year	
None	2 (0.5)
1-3	20 (4.5)
4-12	3 (0.6)
More than 12	2 (0.4)
Sleep disturbance by wheezing in last year	
Never woken with wheezing	13 (2.9)
Less than one night per week	10 (2.2)
One or more nights per week	3 (0.8)
Severe attacks of wheezing limiting speech in last year	6 (1.3)
Lifetime doctor diagnosed asthma	8 (1.7)
Wheezing after exercise in last year	20 (4.5)
Walking with cough in last year	123 (27.6)
Total	446 (100)

The prevalence of frequent tonsillitis was 19.9%. The results of tonsil scoring were as follows: Grade 1, 62.7%; Grade 2, 28.4%, Grade 3, 3.3%; Grade 4, 0.1%; tonsillectomy, 5.5%. Twenty four patients were found to have had a tonsillectomy and/or adenoidectomy. The prevalence of tonsillar hypertrophy in the school children was 3.4%.

A history of frequent tonsillitis was found to be significantly

associated with tonsillar hypertrophy (Odds ratio [OR] = 3.45; 95% confidence interval [CI] = 2.03-5.88). The history of frequent tonsillitis and tonsillar hypertrophy decreased with increasing age (Fig. 1).




Figure 1: The relationship between tonsillar hypertrophy and frequent tonsillitis according to age.

The number of children with frequent tonsillitis and tonsillar hypertrophy who showed a high level of asthma-related symptoms was significant. Frequent tonsillitis was found to be significantly associated with lifetime wheezing (OR = 2.84; CI = 2.15-3.74), 12-month wheezing (OR = 1.56; CI = 0.95-2.56), exercise wheezing (OR = 2.32; CI = 1.44-3.74) and waking with a cough (OR = 2.31; CI = 1.81-2.96). Frequent tonsillitis and asthma-related symptoms were found to decrease with increasing age except for 12-month wheezing which showed two peaks at 5 and 16 years, respectively (Fig. 2). Tonsillar hypertrophy was found to be significantly associated with lifetime wheezing (OR = 1.95; CI = 1.08-3.52).




Figure 2: The relationship between frequent tonsillitis and asthma-related symptoms according to age.

4. Discussion

This study aimed at establishing the relation between chronic tonsillitis and childhood asthma among 4-17 year old children at Dhaka Medical College Hospital, where the reporting children came from all over the Dhaka city and even from far off places of Bangladesh, where air pollution is a major problem. The prevalence of childhood asthma in Bangladesh is not well documented; however related studies prove it to be high. As reported to be 16.1% in one of the studies.^[14] The prevalence of asthma using the ISAAC questionnaire among school children has been reported to vary between 4.9% and 16.4% in Turkey^[15-22] and between 2.1% and 30.2% in some other countries.^[23] Bangladesh is a country with different geographic regions, where plant cover, society structure, socioeconomic status, climate, and level of air pollution vary. These factors can explain the

different prevalence ratios of allergic diseases obtained in different cities and countries. High rates for asthma and asthma-related symptoms are usually reported in English-speaking countries.^[24] In Bangladesh the discrepancy in the prevalence of symptoms of allergic diseases may be due to the selection of different age groups and regions and the fact that the parents completed the questionnaires. The low rates in Dhaka may be associated with a lower awareness of allergic diseases.

There was a positive correlation between frequent tonsillitis and tonsillar hypertrophy; thus frequent tonsillar infection may cause tonsillar hypertrophy. Patients with a history of recurrent tonsillitis had larger tonsils than those without tonsillitis.^[24] In the present study, children with tonsillar hypertrophy had significantly higher frequency of lifetime wheezing than children with a normal tonsil size. Other asthma-related symptoms were also common in children with tonsillar hypertrophy, but they were not statistically significant. These findings suggest that tonsillar hypertrophy might be a risk factor for asthma-related symptoms.

We also observed that the number of children with frequent tonsillitis who showed a high level of asthma-related symptoms was significant. Allergic children may have an increased tendency to upper airway diseases since the airways of asthmatics are more vulnerable due to allergic inflammation. Most asthmatic children might also have allergic rhinitis.^[26] Children with allergic rhinitis usually had nasal congestion causing them to breathe through the mouth. This might lead to frequent tonsillo-pharyngitis and enlarged tonsils resulting in tonsillectomy. Since respiratory viruses were the most common cause of upper airway infections¹ and these were well recognized as major triggers of acute exacerbation of asthma in children,^[7] frequent tonsillitis could trigger acute asthma episodes and asthma-related symptoms.

Exercise-induced wheezing in the previous year was statistically low in children with tonsillectomy/and or adenoidectomy. This might be explained by improvement in the course of the disease due to the removal of a triggering factor, the tonsils as the focal source of infection precipitating frequent upper airway infections in asthmatic children. The lack of a statistical relationship between other symptoms and tonsillectomy and/or adenoidectomy might be due to the small number of children with tonsillectomy/and or adenoidectomy.

5. Future Implications

Thus, it is expected that this study would be a suitable baseline source for future trends on the prevalence, diagnoses and severity of asthma among 4–17 year old children. Frequent tonsillitis and tonsillar hypertrophy could be evaluated as risk factors for asthma due to their significant association with asthma-related symptoms.

References

[1] Roncevic N, Popadic J, Stojadinovic A. Treatment of acute upper respiratory tract infections in children. *Med Pregl* 2002; 55: 397-400.

[2] Jain N, Lodha R, Kabra SK. Upper respiratory tract infections. *Indian J Pediatr* 2001; 68: 1135-8.

[3] Mannino DM, Homa DM, Akinbami LJ, Moorman JE, Gwynn C, Redd SC. Surveillance for asthma - United States, 1980–1999. *MMWR Surveill Summ* 2002; 51: 1-13.

[4] The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. *Lancet* 1998; 351: 1225-32.

[5] Anderson HR, Butland BK, Strachan DP. Trends in prevalence and severity of childhood asthma. *BMJ* 1994; 308: 1600-4.

[6] Burr ML, Butland BK, King S, Vaughan-Williams E. Changes in asthma prevalence: two surveys 15 years apart. *Arch Dis Child* 1989; 64: 1452-6.

[7] Tan WC. Viruses in asthma exacerbations. *Curr Opin Pulm Med* 2005; 11: 21-6.

[8] Hak E, Rovers MM, Sachs AP, Stalman WA, Verheij TJ. Is asthma in 2-12 year-old children associated with physician-attended recurrent upper respiratory tract infections? *Eur J Epidemiol* 2003; 18: 899-902.

[9] Modrzyński M, Mazurek H, Zawisza E. Allergic tonsillitis: myth or reality. *Postepy Hig Med Dosw (online)* 2005; 59: 450-6.

[10] The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variations in the prevalence of asthma symptoms: the International Study of Asthma and Allergies in Childhood (ISAAC). *Eur Respir J* 1998; 12: 315-35.

[11] Brodsky L, Moore L, Stanievich JF. A comparison of tonsillar size and oropharyngeal dimensions in children with obstructive adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 1987; 13: 149-56.

[12] Friedman M, Tanyeri H, La Rosa M, et al. Clinical predictors of obstructive sleep apnea. *Laryngoscope* 1999; 109: 1901-7.

[13] Walker RP. *Snoring and obstructive sleep apnea*. In: Bailey JB, Calhoun KH, Deskin RW, editors. *Head & Neck Surgery - Otolaryngology*. 2nd edition, Philadelphia, Lippincott-Raven Press, 1988; pp. 707-29.

[14] Khalequz zaman, et al. Asthma in rural Bangladeshi children, *Indian Journal of Paediatrics*, Vol 74- June 2007

[15] Selcuk ZT, Caglar T, Enunlu T, Topal T. The prevalence of allergic diseases in primary school children in Edirne, Turkey. *Clin Exp Allergy* 1997; 27: 262-9.

[16] Ones U, Sapan N, Somer A, et al. Prevalence of childhood asthma in Istanbul, Turkey. *Allergy* 1997; 52: 570-5.

[17] Kucukoduk S, Aydin M, Cetinkaya F, Dinc H, Gurses N, Saraclar Y. The prevalence of asthma and other allergic diseases in a province of Turkey. *Turk J Pediatr* 1996; 38: 149-53.

[18] Saraclar Y, Sekerel BE, Kalayci O, Cetinkaya, et al. Prevalence of asthma symptoms in school children in Ankara, Turkey. *Respir Med* 1998; 92: 203-7.

[19] Karaman O, Turkmen M, Uzuner N. Allergic disease prevalence in Izmir. *Allergy* 1997; 52: 689-90.

[20] Canitez Y, Sapan N. The prevalences of asthma, allergic

- rhinitis, and eczema in Bursa, Turkey: an ISAAC study. *J Allergy Clin Immunol* 2000; 105: S318.
- [21] Ece A, Ceylan A, Saraclar Y, Saka G, Gurkan F, Haspolat K. Prevalence of asthma and other allergic disorders among schoolchildren in Diyarbakir, Turkey. *Turk J Pediatr* 2001; 43: 286-92.
- [22] Bayram I, Guneser-Kendirli S, Yilmaz M, Altintas DU, Al-parслан N, Bingol-Karakoc G. The prevalence of asthma and allergic diseases in children of school age in Adana in south-ern Turkey. *Turk J Pediatr* 2004; 46: 221-5.
- [23] Williams H, Robertson C, Stewart A, *et al.* Worldwide varia-tions in the prevalence of symptoms of atopic eczema in the International Study of Asthma and Allergies in Childhood. *J Allergy Clin Immunol* 1999; 103: 125-38.
- [24] Asher MI, Weiland SK. The International Study of Asthma and Allergies in Childhood (ISAAC). ISAAC Steering Committee. *Clin Exp Allergy* 1998; Suppl 5: 52-66 (discus-sion 90).
- [25] Webb CJ, Osman E, Ghosh SK, Hone S. Tonsillar size is an important indicator of recurrent acute tonsillitis. *Clin Oto-laryngol Allied Sci* 2004; 29: 369-71.
- [26] Stone KD. Atopic diseases of childhood. *Curr Opin Pediatr* 2002; 5: 634-46.

Author Profile

Dr. Rameez Shah (Srinagar, Kashmir) MBBS, MS, MIMA (India) Dept of Otolaryngology and Head Neck Surgery, Dhaka Medical College, Dhaka Bangladesh

Dr. Md. Bahauddin (MBBS, DLO) ENT & Head-Neck Cancer Hospital Institute, Dhaka, Bangladesh

Dr. Md.Rafiqul Islam (MBBS, DLO) Diabetic Association Medical College Hospital, Faridpur, Bangladesh

Dr. Md.Shahriar Islam Dept of Otolaryngology and Head Neck Surgery, Dhaka Medical College, Dhaka Bangladesh

Dr. S.M.Sadlee Dept of Internal Medicine, Dhaka Medical College, Dhaka Bangladesh

Dr. Mostafa Kamal Arefin Dept of Otolaryngology and Head Neck Surgery, Dhaka Medical College, Dhaka Bangladesh