

Studies on Zooplanktonic Diversity of River Yamuna in Yamunanagar (Haryana) India

Priyanka Malhotra

Department of Zoology, Kurukshetra University, Kurukshetra – 136119

Abstract: Zooplankton contribute significant role in aquatic ecosystem. Any change in the quality of water is reflected in its community structure. The present paper deals with the qualitative and quantitative analysis of zooplankton in Yamuna river which receives industrial effluents and domestic sewage from point and nonpoint sources. The abundance, distribution, total population, group percentage and species diversity were studied and correlated with pollution indicating water quality characteristics. Species diversity values indicated a decrease from pre effluent point to effluent discharge channel and post effluent discharge point. Cladocera was recorded as the dominant group followed by Rotifera, Copepoda, Protozoa and Ostracoda.

Keywords: Industrial effluents, Species diversity, Sewage, Yamuna River, Zooplankton

1. Introduction

Aquatic life depends upon Planktonic populations. Any change in the quality of water directly affects its planktonic population *i.e* they serve as good indicators of health of aquatic ecosystem. Zooplankton community structure has significant potential for assessing aquatic ecosystem health. Their dominance and seasonality are highly variable in different water bodies according to nutrient status, age, morphometry and other locational factors (Jose and Sanal kumar, 2012). They are sensitive indicators of pollution in comparison with phytoplankton (Umadevi, 2013).

Yamuna is the largest tributary river of Ganges in north India (Negiet *al.*, 1991). Yamunanagar (30° 6' N latitude and 77° 17' E longitude) is an important industrial city of Haryana. The river Yamuna meanders through the district Yamunanagar (Haryana), India, and forms the eastern boundary with the neighbouring Saharanpur district. Along its path river is getting effluents from the maskaranala from

Saharanpur (Uttar Pradesh), India which is affecting the ecology of the river. Hence the present study was undertaken in order to relate the effect of industrial effluents on zooplankton diversity of river Yamuna.

2. Materials and Methods

Keeping in view the point of influx of discharges into river, three stations have been selected. Station-Y1 lies in village Kalanaur at upstream of the river before the influx of discharges, Station-Y2 lies 4-5 Kms downstream from station Y1 at middle reach of the river where the industrial effluents join the river, Station-Y3 at 5-6 kms downstream from station-Y2 after the influx of discharges (Fig. 1). Zooplankton samples were collected by filtering 25 L of water through plankton net of mesh size 50µm with demarcating collecting tube. The abundance of zooplankton was expressed as organisms L⁻¹. The organisms counted by drop count method were expressed per litre using formula:

$$\text{Total Planktons L}^{-1} = \frac{\text{Number of Organisms per drop} \times \text{Vol. of conc. sample in ml}}{\text{Volume of original sample in litres} \times \text{Vol. of one drop}}$$

Species Diversity of zooplankton was determined using Shannon and Weaver diversity index method (Shannon and Weaver, 1963; Washington, 1984).

$$D = - \sum \frac{n_i}{N} \log \frac{n_i}{N}$$

Where, D = Species Diversity, n_i = Number of individuals of ith species, N = Total number of individuals in the sample. The coefficient of correlation "r" was calculated on computer using SPSS package.

3. Results and Discussion

Thirteen taxa of zooplankton were recorded from different stations of river Yamuna including 4 of Cladocera, 3 of Rotifera, 2 of each Copepoda and Protozoa and 1 each of Ostracoda and Hymenoptera. The maximum density was recorded at station Y1 followed by station Y3 and station Y2. The mean zooplankton population was 300 L⁻¹ at station Y1, 193 L⁻¹ at station Y2 and 248 L⁻¹ at station

Y3. Maximum numbers of total zooplankton were found in July at all the stations (Table 1). With increase in warmth, the animals become more active, grow more rapidly and breed more quickly (Mellanby, 1963). Cladocera was the dominant group followed by Rotifera, Copepoda, Protozoa and Ostracoda. One genera of Hymenoptera, Polynema spp. was also observed along with zooplankton. Dominant group Cladocera with 33.5% was represented by Moina spp., Sida spp., Bosmina spp. and Ceriodaphnia spp. (Fig. 3). According to Szerocry n' Ska (2002) and Abranteset al. (2006) Cladocera indicated the eutrophic conditions resulted from pollution. Michael (1985) also designated Cladocerans as bio-indicators. High number of Cladocera in the present studies supports the view. Moina spp. was recorded as tolerant taxa common to all stations. Bilgramiet al. (1985) and Bulusuetal. (1967) have reported that Moina spp. is tolerant to heavy pollution. Mageed (2007) has also designated Moina spp. as dominant and tolerant taxa in lake Manzala of Egypt. The mean values for Sida spp. was high

at station Y1 but showed a decline at Y2 and Y3 depicting it as sensitive taxa. Rotifera the second dominant group was represented by *Brachionus* spp., *Keratella* spp. and *Monostylas* spp.. The role of Rotifera as bioindicators has been emphasized by Arora (1966). In the present studies, *Keratella* spp. and *Brachionus* spp. were common rotifers with a wide range of tolerance to different physicochemical conditions. Copepoda was represented by *Cyclops* spp. and Nauplius larva. Bhatti and Rana (1987) and Wolfram et al. (2002) regarded *Cyclops* spp. and Ostracods as strictly pollution sensitive taxa. In the present studies, it was although present at all stations but number was low. Similarly, *Cypris* spp. the only member of Ostracoda was also low in number. Protozoans were represented by two genera *Trinema* spp. and *Physarum* spp.. Total population of zooplankton showed a significant ($P < 0.05$) decrease from station Y1 ($300 \pm 41 L^{-1}$) to station Y2 ($193 \pm 28 L^{-1}$) and then increase at station Y3 ($248 \pm 31 L^{-1}$). This may be due to mixing of effluents through 'maskaranala' at station Y2. Statistically also, zooplankton showed a significant positive correlation with DO ($r = 0.604$, $P < 0.05$) and significant negative correlation with ammonia ($r = -0.498$, $P < 0.05$) advocating that ammonia was the important pollution indicating parameter causing a decline of plankton population. BOD also showed a significant negative correlation with zooplankton but the values were not statistically significant. Species diversity of zooplankton depicts a decreasing trend from station Y1 to Y2 and further increased at station Y3 indicating the effect of industrial pollution and sewage waste (Fig. 2). Trivedi (1981) and Malhotra et al. (2014) have emphasized the role of species diversity index in pollution and stated that a decrease in species diversity values point to polluted waters. These results depict station Y2 as stressed area with reference to species diversity.

4. Conclusion

The low population density of the Zooplanktons in river water depicted that river Yamuna is in a very poor tropic status. Total number and Species diversity of zooplankton decreased at station Y2 where industrial and sewage channel joins the river depicts the altered overall ecology of the stream thus reducing the capture fishery statistics. Therefore, the proper and efficient treatment of sewage should be carried out before discharging them into the river system.

References

- [1] Abrantes, N., Antunes, S.C., Pereira, M.J. and Goncalves, F. 2006. Seasonal succession of Cladocerans and phytoplanktons and their interactions in a shallow eutrophic lake (Lake Vela, Portugal). *Acta Oecologica*. 29(1): 54-64.
- [2] Arora, H.C. 1966. Rotifera as indicators of trophic nature of environments. *Hydrobiologia*. 27(1-2): 146-149.
- [3] Bhatti, D.S. and Rana, K.S. 1987. Zooplankton in relation to abiotic components in the fort moat of Bharatpur. Proceedings of National Academy of Sciences, India. 57(B): 111.
- [4] Bilgrami, K., Datta, S., Munshi, J.S. and Bhowmick, B.N. 1985. Biomonitoring of river Ganga at polluted sites in Bihar. In: Proceedings of Symposium on Biomonitoring state Environment. India: 141-145.
- [5] Bulusu, K.R., Arora, H.C. and Adoo, K.M. 1967. Certain observations on self-purification on Khan river and its effects on Kshipra river. *Environmental Health*. 9: 275-295.
- [6] Jose, R. and Sanalkumar, M.G. 2012. Seasonal Variations in the Zooplankton Diversity of River Achencovil. *International Journal of Scientific and Research Publications*. 2(11): 1-5.
- [7] Mageed, Adel A.A. 2007. Distribution and long term historical changes of zooplankton assemblages in lake Manzala (south Mediterranean sea, Egypt). *Egyptian Journal of Aquatic Research*. 33(1): 183-192.
- [8] Malhotra, P., Bhatnagar, A. and Chopra, G. 2014. Phytoplankton Diversity of Western Yamuna Canal and River Yamuna in Yamunanagar, Haryana, India. *International Research Journal of Environment Sciences*. 3(2): 1-7.
- [9] Mellanby, H. 1963. Animal life in fresh water. 6th Ed. Chapman and Hall Ltd., London.
- [10] Michael, R.G. 1985. Use of rotifers and cladocerans as potential bioindicators of Indian fresh water Ecosystem. In: Proceedings of Symposium on Biomonitoring state Environment. India, 1985:82-83.
- [11] Negi, S. S. (1991). Himalayan Rivers, Lakes and Glaciers, Indus Publishing Co., New Delhi, 182.
- [12] Shannon, E.E. and Weaver, W. 1963. The mathematical theory of communication, University of Illinois, Press, Urhanna, 117.
- [13] Szerocry, N Ska. Krystyna. 2002. Human impact of lakes recorded in the remains of Cladocera. *Quaternary International*. 95-96:165-174.
- [14] Trivedi, R.C. 1981. Use of diversity index in evaluation of water quality. WHO workshop on biological indicator and indices of environmental pollution. Cent. Bd. Prev. Cont. Water Poll/ Osm. Univ. Hyderabad, India: 175-188.
- [15] Umadevi, T. 2013. Limnological investigation and zooplankton diversity of Karanjriver, Karnataka. *International journal of Science and Research*. 2(3): 133-136.
- [16] Washington, H.G. 1984. Diversity, biotic and similarity indices: A review with special relevance to aquatic ecosystems. *Water Res.* 18: 653-694.
- [17] Wolfram, G., Kowarc, V.A., Humpesch, U.H. and Siegi, W. 2002. Distribution pattern of benthic invertebrate's communities in relation to industrial tailings and Trophy. *Water, Air and Soil Pollution*. 24(4): 63-91.

Legends to Figures

- [1] Fig. 1: Map showing location of selected stations on river Yamuna.
- [2] Fig. 2: Species diversity of zooplankton of river Yamuna at various stations.
- [3] Fig. 3: Percentage distribution of zooplankton of river Yamuna at various stations.



Figure 1: Map showing location of selected stations on river Yamuna




Figure 2: Species diversity of zooplankton of river Yamuna at various stations




Figure 3: Percentage distribution of zooplankton of river Yamuna at various stations

Table 1: Total number of zooplankton (L-1) (mean± S.E of mean) of river Yamuna at various stations

	Station Y1	Station Y2	Station Y3
August	178±9.94	107±11.3	172±21.6
September	192±5.66	134±11.5	175±3.56
October	173±11.2	136±23.5	166±10.1
November	192±13.0	128±3.78	154±12.9
December	135±14.7	110±11.4	148±11.7
January	218±12.7	127±2.12	184±19.1
February	207±5.3	95±0.47	156±3.77
March	348±34.6	248±3.6	316±5.44
April	433±4.64	210±2.88	365±10.8
May	500±15.5	274±14.1	321±7.64
June	492±41.6	336±29.6	350±17.9
July	532±32.6	412±33.9	473±27.5

