









- Cladophoraglomerata and Enteromorphaahlnneriana,” J Exp Mar BiolEcol, 298: pp. 111-123, 2004.
- [18] M.B. Arnao, “some methodological problems in the determination of antioxidant activity using chromogen radicals: a practical case,” Trends Food Sci Technol, 1:419, 2000.
- [19] V. Horincar, G. Parfene, G. Bahrim, “Evaluation of bioactive compounds in extracts obtained from three romanian marine algae species,” Romani BiotechnolLett, 6: pp. 71-78, 2011.
- [20] V. Lagouri, D. Boskou, “Nutrient antioxidants in oregano,” Int J Food Sci Nutr, 47: pp. 493-497, 1996.
- [21] M. Rajesh, A. Nagarajan, S. Perumal, M. Sellamuthu, “The antioxidant activity and free radical scavenging potential of two different solvent extracts of Camellia sinensis, Ficusbengalensisand Ficusracemosa,” Food Chem, 107: pp. 1000–1007, 2008.
- [22] V. Alexandru, M. Balan, A.Gaspar, O. Craciunescu, I. Moldovan, “Studies on the antioxidant activity, phenol and flavonoid contents of some selected Romanian medicinal plants used for wound healing,” Biotech Lett, 12(6): pp. 3467-3472, 2007.
- [23] K. M. Yoo, C.H. Lee, H. Lee, B. Moon, C. Y. Lee, “Relative antioxidant and cytoprotective activities of common herbs,” Food Chemistry, 106: pp. 929–936, 2008.
- [24] R.L. Prior, X. Wu, K. Schaich, “Standardized methods for the determination of Antioxidant capacity and phenolics in foods and dietary supplements,” J Agri Food Chem, 53: pp. 290-302, 2005.
- [25] Y. V Yuan, N.A. Walsh, “Antioxidative and ant proliferative activities of extracts from a variety of edible seaweeds,” Food Chem. Toxicol, 44: pp. 1144-1150, 2006