

A Study of Some Physicochemical Parameters and Heavy Metals in the Diwaniyah River / Euphrates, Iraq

Foad M. Alkam¹, Mohammed J. S. Al-Haidarey², Kasim K. Alasedi³

¹Department of Biology, College of Education, University of Qadisia, Diwaniyah, Iraq

²Department of Ecology, Faculty of Science, University of Kufa, Kufa, Najaf, Iraq

³Head of Ecology Dep., Research field: Analytical Chemistry and Environmental Pollution, Department of Ecology, Faculty of Science, University of Kufa, Kufa, Najaf, Iraq

Abstract: Metals tend to accumulate in water and move up through the food chain. So, studies to ascertain the level of heavy metals in environment and determine potentially hazardous levels for human are necessary. Samples were monthly collected from three sites along the Diwaniyah River, during Autumn 2012 to Summer 2013. In this investigation, concentration of four heavy metals copper (Cu), zinc (Zn), cadmium (Cd) and lead (pb) in water and sediments have been determined. In addition, some physiochemical parameters of water were investigated. Heavy metal concentrations were measured by flame atomic absorption spectrometry. The results showed that the highest concentrations of heavy metals in water and sediment samples were related to pb and Zn. However, the observed heavy metals concentrations in water and sediments were below the recommended limits for WHO. The investigation showed elevating levels of heavy metals in environment. Thus, a serious notification to industrial and manmade pollution, which can lead to ecosystem and food chain contamination, is necessary.

Keywords: Diwaniyah River, Limenological, Heavy metals, Water, Sediment.

1. Introduction

Environmental pollution is a worldwide problem, heavy metals belonging to the most important pollutants. The progress of industries has led to increased emission of pollutants into ecosystems. In natural aquatic ecosystem, heavy metals occur in low concentrations mainly due to weathering of soils and their associated bedrocks. In recent times however there has been an unprecedented increase in the level of these metals due to human activities. The occurrence of such metals in excess of natural load has become a problem of increasing concern not only to environmentalists but also to health practitioners. The concern about heavy metals stems from their persistence in the environment as they are not easily degraded either through biological or chemical means unlike most organic pollutants. The discharge of various metal pollutants in to the aquatic environment as pointed out by [1,2] could damage the quality of the ecosystem thereby rendering water body unsuitable for its intended uses posing a serious health threat to the immediate population. Heavy metals do not exist in soluble forms for a long time in water; they are present mainly as suspended colloids are fixed by organic and mineral substances [3].

longitude coordinates are from 32° 02' 48.38" N, 44° 46' 42.27" E to 31° 39' 15.48" N, 45° 01' 53.42" E respectively). For investigation of study, 3 sites were chosen along the Diwaniyah River as show in figure 1. The 1st site is before entrance the river to city, the 2nd site is in the mid of the city, and 3rd site is after the river passing the domestic wastewater treatment plant and factories (textile, and rubber factories).



Figure 1: The study sites

2. Materials and Methods

2.1. Study Area

Diwaniyah river (Shatt Al-Diwaniyah) is an extension of the Hilla river in the area stretching from the village of Sunnia which is north of the city, even the southern city of Diwaniyah, about 15 km inside the city (latitude and

2.2. Sampling Strategy

Samples of water and sediment were collected from three different sampling sites along the river (study site). The samples were collected monthly (started with November 2012 and finished at August 2013). At each of the sampling sites triplicate samples were collected. Water samples were collected at the depth of 30cm below the surface using 1liter

polyethylene bottles with screw caps which were acid washed and rinsed with distilled water prior to the sampling.

Samples were acidified with 1.5ml concentrated Nitric acid (HNO₃)/liter of sample, after air temperature (AT), water temperature (WT), water flow (WT), transparencies, dissolved O₂ (DO), pH (Digital Inolab 720, WTW-Germany) and conductivity (Model 340i/SET, WTW-Germany) (EC) measurements in the field and subsequently transported to the laboratory for analysis. Total dissolved solids (TDS) were determined by WTW multiple meter model 2005(origin Germany), total alkalinity (TA) and oxygen dioxide (CO₂) was determined according to the procedure [4], EDTA titrimetric method was used to determine total hardness as recommended by WHO [5]. Sediment samples (top 10cm) were collected using soil auger, kept on ice in polythene bag and transported to the laboratory [6]. Heavy metals (HMs) measured by flame atomic absorption spectroscopy type Shimadzu AA-6300 Japan according to method that described by APHA [7].

3. Results and Discussion

The mean values ± standard deviation of physiochemical analysis of water samples obtained from the study sites are represented in table 1. Air and water temperatures were in the range 12-35 °C to 14-40.12 °C, respectively, Electrical conductivity was observed 940 μs/cm at site 1 during Jan 2013 and 1193 μs/cm at site 3 in August 2013 respectively, TDS ranged from 113 -1375 mg/L in sites 1 (during July) and 3 (during Aug.) respectively, these values increased in winter months and decreased in summer months [8, 9].

During the study period, the pH was in the ranges from 6.7 to 8.1, during September and April respectively, water flow ranged 21- 44Cm/min in site 3 and 1 during march and October respectively, turbidity ranged from 16 NUT – 52 NUT in site 2 and 3 during September and October respectively, and transparency ranged between 26 Cm -71 Cm in site 1 during July and March respectively.

The dissolved oxygen, and BOD₅ were 4.2 mg/L - 8.9 mg/L (in site 3 and 1 during Aug. and March), and 1.2 mg/L – 5.9 mg/L (during Feb. and Aug. in site 1 and 3) respectively, the values of BOD₅ and Dissolved oxygen that recorded were in limits recommended by EPA [10] for the rivers water. The carbon dioxide, Alkalinity, and total hardness values ranged between 121mg/L – 269 mg/L, 119 mg/L – 271 mg/L, and 275 mg/L – 546 mg/L during April and August in site 1 and 3 respectively.

As show in Table 2, the correlations among TDS, EC, air and water temperature was positive, Positive correlations were recorded between total alkalinity, dissolved oxygen and pH, Many studies have found that Iraqi water was slightly alkalinity [11-14]. The present study found high concentrations of total hardness in all sites during different months, with no significant difference among study sites while there were a negative correlation with dissolved oxygen and positive correlation with water and air temperature. The solubility of oxygen in an aquatic system is influenced the temperature and dissolved salts [15], so the

correlation among dissolved oxygen and air temperature, water temperature, turbidity, total hardness, electrical conductivity, carbon dioxide and BOD₅ were negative (Table 2).

Table 3; show the concentration of heavy metals in water and sediments in Euphrates River. The results of this study showed that the positive correlation between heavy metals in dissolved phases with air and water temperature, dissolved oxygen, BOD₅, turbidity, EC, CO₂, pH, Total hardness, alkalinity, and TDS while they have a negative correlations with other physiochemical parameters (Figure:3), that was agree with [16]. The concentrations of some heavy metals in water of Euphrates River were low according to WHO and USEPA guidelines, but there were high compared to other Iraqi aquatic systems [17, 18]. This difference with other Iraqi studies may be related to the difficulty of investigating heavy metals in toxic aquatic system, where heavy metal concentrations will be low. Moreover the many factors affect the concentrations, such as: the flow of the dredged materials from upper regions of the river, dilution and increase of water flow, direct drainage from farmlands, factories, sewage disposal plants, dissolution of sediments, increases in the numbers of phytoplankton in water, bioaccumulation, chemical adsorption on sediments and complexes with organic matter [19-21] 57-60. The concentration of heavy metals in all phases were Cd < pb < Cu < Zn, that means the source and fate of these metals are one [22, 23] and that were very clear in table 4, and figure 2.

Table 1: Physiochemical parameters of water in Euphrates River / Diwanayah City, first line :(mode), second line: mean (±S.D).

	Site.1	Site.2	Site.3
Water temperature °C	12-34 21.8±7.5	12-35 21.98±7.496	12.3-34 21.783±7.049
Air temperature °C	14-39 25±7.7	14-39.5 25.375±7.652	15.5-40.12 26.55±7.519
Water flow	25.5-44 34±6.1	25-41 33.28±5.343	21-37 28.74±5.44
Turbidity NUT	25-49 37±7.8	16-50 36.42±9.77	31-52 41.548±7.45
Transparency Cm	20-71 46.8±17.7	28-66 44.85±13.45	24-60 40.83±13.30
EC μs/cm	940-1133 1028±62.8	1009-1178 1111±60.13	1018-1198 1119. ±60.53
TDS mg/L	113-1043 880±247.9	901-1365 1089±153.47	900-1375 1099±167.6
pH	7.5-8.1 7.8±0.24	7.5-8.1 7.8±0.21	6.7-8.1 25.8±32.88
Total hardness mg(CaCO ₃)/L	275-546 411±79.5	331-516 409.92±53.86	305-522 401.41±68.08
CO ₂ mg/L	121-201 163.3±25.3	126-245 176.42±38.22	135-269 188.08±42.59
Total alkalinity mg(CaCO ₃)/L	119-185 141.2±25.2	121-179 144.92±17.56	119-271 175.58±52.6
O ₂ mg/L	5.2-8.9 7.4±1.11	4.9-7.9 7±0.881	4.2-8 6.5167±1.01
BOD ₅ mg/L	1.2-4.9 2.04±1.06	1.6-5.2 ±1.041	2-5.9 2.742±1.13

Table 2: The correlation among physiochemical parameters of water during the period of study

WT	0.98	1.00																	
WF	0.05	0.12	1.00																
TUR	0.57	0.57	0.32	1.00															
TRANC.	-0.68	-0.66	-	-	1.00														
pH	0.43	0.39	-	0.33	-0.29	1.00													
TH	0.65	0.64	0.02	0.36	-0.65	0.35	1.00												
EC	0.64	0.62	0.11	0.38	-0.39	0.38	0.27	1.00											
CO2	0.62	0.58	-	0.38	-0.54	0.64	0.70	0.39	1.00										
TALKA	0.15	0.10	-	-	0.14	0.81	0.05	0.25	0.45	1.00									
O2	-0.75	-0.70	-	-	0.84	-	-	-	-0.72	-0.12	1.00								
BOD5	0.82	0.79	-	0.55	-0.62	0.54	0.71	0.56	0.81	0.33	-0.87	1.00							
TDS	0.39	0.36	-	0.26	-0.04	0.41	0.05	0.57	0.41	0.43	-0.30	0.43	1.00						
AT	WT	WF	TUR	TRANC.	pH	TH	EC	CO2	TALKA	O2	BOD5	TDS							

Table 3: Mode (first line), Average ± S.D (second line) of different phases of heavy metals in Euphrates River / Diwaniyah City

Element	Water		Sediment		Water		Sediment	
	Dissolved	Particulate	Exchangeable	Residual	Dissolved	Particulate	Exchangeable	Residual
Cd	ND-1.05	2.94-9.15	0.09-2.7	0.68-4.56	ND-1.9	3-10.9	0.94-2.6	0.94-2.6
	0.4675±0.37	4.9582±1.841	1.36±1.07	2.39±1.66	0.7092±0.6153	5.9992±2.5725	1.73±0.89	1.73±0.89
Zn	2.01-10.51	35.21-400	6.55-12.65	20.05-36.25	5.62-15.98	40.19-130.3	9.2-18.06	9.2-18.06
	7.7658±2.2027	75.369±102.43	9.52±2.83	27.39±6.77	11.465±3.1279	72.9142±34.1678	15.04±3.9	15.04±3.9
Pb	0.5-1.32	4.12-9.55	0.46-1.75	13.1-23.53	ND-2.9	5.99-13.11	0.81-2.08	0.81-2.08
	0.9583±0.2501	6.7317±1.976	0.92±0.58	18.73±4.52	1.0708±0.7774	8.8558±2.0053	1.42±0.54	1.42±0.54
Cu	2.45-4.01	27.13-41.12	4.9-1.75	17.52-22.37	2.71-5	28.5-44.41	5.75-9.78	5.75-9.78
	3.3458±0.584	32.2017±3.5069	6.91±1.34	20.02±2.11	3.6942±0.8148	35.6583±4.4763	7.74±1.73	7.74±1.73




Figure 2: The mean of Seasonal variation of Dissolved (A), Particulate (B), Exchangeable (C), and Residual (D) heavy metals in study sites.




Figure 3: The correlation between dissolved (A), Particulate (B) heavy metals and physiochemical properties in water during the period of study

Table 4: The correlation between particulate and dissolved heavy metals in period of study

Cd Diss	1							
Cd Parti	0.566284	1						
Zn Diss	0.682515	0.66217	1					
Zn Parti	0.36451	0.193174	0.326444	1				
Pb Diss	0.624618	0.336094	0.503229	0.459793	1			
Pb Parti	0.459987	0.241264	0.426779	0.426568	0.961602	1		
Cu Diss	0.607253	0.299806	0.491973	0.456734	0.970438	0.956628	1	
Cu Partic	0.461781	0.297996	0.422281	0.50834	0.91414	0.949575	0.920813	1
	Cd Diss	Cd Parti	Zn Diss	Zn Parti	Pb Diss	Pb Parti	Cu Diss	Cu Partic

Author Profile



Prof. Dr. Foad M. Alkam Major: Ecophysiology, Interest: Water pollution, Biology department, college of education, University of Al-Qadisiya, Iraq.



Mohammed J. S. Al-Haidarey is senior lecturer in Department of Ecology, Faculty of Science, Uni. of Kufa, Iraq. Area of research is ecotoxicology and wetland biogeochemistry



Kasim Kadhim Alasedi is Assistant professor in Analytical Chemistry. During 2000-2007 he has worked in Chemistry field and translation in USA. Currently, he is Head of Ecology Department, Faculty of Science, University of Kufa, Iraq.

References

[1] J.F. Abowei and F.D. Sikoki, Water Pollution Management and Control, Double trust. Publication. Company, Port Harcourt. pp.236, 2005.

[2] S.M. Saeed and I.M. Shaker 8th International Symposium on Tilapia in Aquaculture, pp.475-488, 2008.

[3] Ebenezer and Y. Eremasi, Scientia Africana 11(1), pp. 44-52, June 2012.

[4] G.T. Lind, "Handbook of Common Methods in Limnology, 2nd Ed., London, U.K, 1979.

[5] WHO "Global Water Supply and Sanitation Assessment" report, 2000.

[6] O. R. Awofolu, Z.Mbolekwa, V.Mtshemla and O. S. Fatoki, Levels of trace metals in water and sediment from Tyumen River and its effects on an irrigated farmland. Water SA, 31(1) 2005.

[7] APHA "Standard Methods for Examination of Water and wastewater" 20th Ed., Washington DC, USA, 2007.

[8] R. G. Wetzel, Limnology, lake and river ecosystems. 3th Ed. Academic Press, An Elsevier Science imprint, San Francisco, New York, London. 729, 2001.

[9] H. Al-Mousawi, H. A. Al-Saadi, and F. M. Hassan, spatial and seasonal variations of phytoplankton populations and related environments in Al-Hammer marsh, Iraq. Bas J. Sci. 12(1) pp. 9-20, 1994.

[10] RPI (River pollution index), RPI classification standard is adjusted with reference to monitoring report decimal rules published by Environmental Analysis Laboratory, starting from 2013.

[11] F.M. Hassan, and H.A. Al-Saadi, On the seasonal variation of phytoplankton population in Hilla River Iraq. J. Col. Edu. for Woman Univ. of Baghdad, 6 (2), pp.55, 1995.

[12] F.M. Hassan, A limnological study on Hilla River. Al-Mustansiryah J., 8 (1), 22: pp. 211-232, 1997.

[13] H. A. Al-Saadi, T. I. Kassim; A. A. Al-Lami and S. K. Salman, Spatial and seasonal variations of phytoplankton population in the upper region of Euphrates River, Iraq. Limnologica 30:pp.83-90, 2000.

[14] Al-Lami, A. W. Sabri, T. I. Kassim, and K. A. Rasheed, Phytoplankton of Samarra Reservoir (Iraq). Acta Hydrobiol. 38(3/4), pp.77-86, 1996.

[15] R. G. Wetzel and G. E. Likens, Limnological analyses 3rd Ed Springer-Verlag New York, 432, 2000.

[16] J. K. Abaychi and A. A. Z. DouAbul, Trace metals in Shatt Al-Arab river, Iraq, Water Res. 19(4), pp.457-462, 1985.

M.M. Al-Taee, A.N. Al-Khateeb, F.H. Hussein, F.M. Abid, Evaluation of soluble non-essential trace metals in Shatt Al-Hilla, Iraq, Asian J. Chem. 19 : pp.741-750, 2007.

[17] F. M. Hassan, M.M. Saleh, J. M. Salman, A Study of Physicochemical Parameters and Nine Heavy Metals in the Euphrates River, Iraq. E-Journal of Chemistry, 7(3), pp. 685-692, 2010.

[18] H. T. AL-Saad, Y. Z. Mustafa, and A. AL-Timeri, Concentration of trace metals in aquatic plants of the AL- Hammer marsh. Iraqi Marina Mesopotamia. 9(2): pp.323-328, 1994.

[19] E. Kaiser, D. B. Arscott, K. Tockner, B. Sulzberger, Sources and distribution of organic carbon and nitrogen in the Tagliamento River, Italy. Aquat. Sci. 66:pp. 103-116, 2004.

[20] M. J. S. Al-Haidarey, F. M. Hassan, A. A. Z DouAbul, A. A. Al-Kubaisy, The Geoaccumulation Index of Some Heavy Metals in Al-Hawizeh Marsh, Iraq, E-Journal of Chemistry. 7(5), pp. S157-S162, 2010.

[21] M. J. S. Al-Haidarey, Using the Site Specific Water Quality Index 1.0 for evaluation of water Quality Index In Shatt Al-Arab River. 2010, Marina Mesopotamia proceeding of the marine science conference, pp.217-228, 2008.

[22] M. A. H. Al-Kenzawi, M. J.S. Al-Haidarey, A.H. Talib, M.F. Karomi, Environmental Study of Some Water Characteristics at Um-Al-Naaj Marsh, South of Iraq. Baghdad Science Journal, 8(1):pp. 531-538, 2011.