

Internal Functions of Matter (m) of Einstein Equation, $E = mc^2$ and Emission of Energy (E) after finding the Mass of a Photon, Graviton and Equation of Unification Physics

Nirmalendu Das

Assistant Engineer, All India Radio, Siliguri (Retd)

Life Member: THE VON KARMAN SOCIETY for Advanced Study and Research in Mathematical Sciences (VKS),
Old Police line, Jalpaiguri, West Bengal (India)

Abstract: In 1905 Einstein introduced his famous equation $E = mc^2$ throughout the scientific world and scientists experimentally tested different ways even in the outer space that the speed of light is $2.99792458 \times 10^{10}$ cm/sec this is very unique equation for the mass equivalent energy system. This applicable to all fields for energy emission. Here the question is that matter is not alone, inside of it other particles are there. Which type of particles is responsible to create energy? Again, energy is composed of bunch or coagulation of photons, how many photons are responsible to create energy? We know the equation $E = hv$ and h is the Planck constant and according to De-Broglie, $v = c/\lambda$. But same question arises that one Planck is $6.6260755 \times 10^{-27}$ erg-sec, how many photons creates this energy? To solve all these facts, we need to think on the photon's mass. Because, energy is composed of number of photons which will depend the mass m . Again, in what way gravitational constant and the particle acts on it that internal mechanism is very important throughout the universe.

Keywords: Mass of a photon, graviton, unification of physics, Planck constant, Einstein's equation.

1. Introduction

The electromagnetic energies are composed of photons as $E = hv$, so photon is the agent of energy. Like the graviton particle is the agent gravitational force in the field of gravitation. The mass of photon and graviton is treated as zero in our modern physics. The zero value is most important in the field of mathematical terms. But in the case of matter, we can't agree because it has mass whatever it may be negligible. So, we need to know the mass of a photon, graviton. Therefore we need to determine the mass of these particles first.

1) Basic Equation And Method To Determine The Mass Of A Photon

According to well known Compton's Theory, when scattering takes place from an electron tightly bound to the atom, collision is, in fact with the atom to which the electron is tightly bound. As the Compton wavelength $\lambda_c = h / m_e c$ is inversely proportional to the mass of the scattering particle. Its magnitude for atom is nearly 10^{-4} times the value of an electron [1]. (The symbols have their usual meanings).

From the above mentioned relation between an electron and atom is seen that electron is tightly bound to the atom. Therefore, according to Compton wavelength the atom (λ_u) and electron (λ_e) must also have close relations between them. Let the ratio of wavelength of atom and electron is directly proportional with the total energy of the system. From this view the following relation will produce energy of the total system as:

$$E_1 = m_e c^2 \times \frac{\lambda_u}{\lambda_e} = m_e c^2 \times 5.485798962 \times 10^{-4} = 4.491284593 \times 10^{-20} \text{ erg} \dots\dots\dots (1)$$

And $E_1/[e] = 4.491284593 \times 10^{-10} \text{ erg} / [e] = 0.2803 \text{ keV}$,
where, $[e] = 1.60217733 \times 10^{-12} \text{ erg}$

a) This energy 0.2803 KeV is showing the average energy of Eigen value of electron [2] between the first excited state and generating state as,

(0.2256 KeV, when, $n_x = 2, n_y = n_z = 1$) + (0.3354 KeV, when, $n_x = n_y = 2, n_z = 1$) = $0.561/2 = 0.2805 \text{ KeV}$.

b) Again, if we multiply 0.2803 keV by Avogadro number (6.0221467×10^{23}), then we get, $N_A \times E_1 = 1.68 \times 10^{17} \text{ GeV}$, this energy is the energy range for unified theory [3] (10^{17} GeV).

From the above two evidence (a) & (b), we can assume that when electron and atom tightly bound, then it has a good function in the microscopic field and justified energy of total system of the equation: $E_1 = m_e c^2 \times \lambda_u / \lambda_e$.

The mass of an atom is constant; there are many particles inside of it. Besides these particles there is also some energy inside the atom which is nothing but the bunch of photons. According to the mass energy equivalence, energy is coagulated or bunched of photons or quanta. So, it is essential to know how many photons are present in an atom having constant mass of a photon (mass of a photon is symbolized by σ). Because, the symbol of a PHOTON giving in many books as γ (gamma, mass = 0), but for different types of gamma rays or energies (e.g. 1.022 etc), the symbol makes complexity. So, for exceedingly small mass of a photon, the symbol σ (small sigma) is used. Avogadro's number is used in the different field of science as Avogadro number of atoms or molecules. If we consider, Avogadro number of photon are present in an atom having photon's mass σ and just before the scattering of energy, if atom is irritated, then it will follow gas law — $E = 1/2 N m \bar{c}^2$ (where,

E = kinetic energy of gas atom, N = number of atom in a volume V, \bar{c}^2 = mean square velocity of the gas atom), then the particles inside the atom will follow the gas law. We can follow this gas law for the photon inside the atom and the energy considered as E_1 . So,

$$E_1 = \frac{1}{2} N_A \sigma c^2 \dots\dots\dots (2)$$

Where, E_1 = Kinetic Energy of Avogadro number of photons, c = velocity of a photon or light, N_A = Avogadro number = 6.0221367×10^{23} .

If this energy, E_1 incorporates with the obtained energy from the equation (1) for balancing the mass energy equivalence, then we get mass of Avogadro number of photons of a particle (m_0) of wavelength (λ') in the form of:

$$N_A \sigma = \frac{2h^2}{m_0 \{\lambda'\}^2 c^2} = 9.994456186 \times 10^{-31} \text{ gm} \dots\dots\dots (3)$$

Here, $m_0 = m_u$ = mass of atom, $\lambda' = \lambda_c$ = Compton wavelength of electron. So mass of a photon is,

$$\sigma = \frac{2h^2}{N_A m_u \lambda_c^2 c^2} = 1.659619614 \times 10^{-54} \text{ gm} \dots\dots\dots (4)$$


Is the mass of a photon? Need to prove it through an example. We know the Stark – Einstein equation as $E\lambda = N_A h c = 0.11962658$ Joule meter per mole. Where, λ = wavelength of the absorbed radiation. We can get this energy from the equation (4) by arranging as:

$$N_A h c = \frac{2h^3}{m_u \lambda_c^2 \sigma c} \dots\dots\dots (5)$$

The formation of L. H. S. of this equation (5), $N_A h c$ is showing the energy of $E\lambda$, so we can write,

$$E\lambda = \frac{2h^3}{m_u \lambda_c^2 \sigma c} = 0.11962658 \text{ Joule} \dots\dots\dots (6)$$

Therefore, the mass of a photon is true and we cannot treat as zero and its application is very wide from the particle to the universe. When the mass of a photon will be treated through the equation of theory of relativity of Einstein ($m = m_0 / \sqrt{1 - v^2/c^2}$), then we will get another equation which is applicable to all fields from particle to the universe. This equation is termed as the equation of UNIPICATON OF PHYSICS. The equation is: (when, $m_1 = 4.033$ amu, mass of α)



(THE EQUATION OF UNIFICATION OF PHYSICS)

The equation (7) is applicable to particle, the difference of energy from rest to excited state is very important. But the value of Pi changes slightly from its normal value which is not shown here. the equation (9) for the heavy mass and applicable to universe.

2. Basic Equation and Method to Determine the Mass of a Graviton

We know the Newton Law of Gravitation is,

$$F = G \frac{m \times M}{r^2} \text{ N} \dots\dots\dots (12)$$

When atom and electron is tightly bounded each other, we can ignore the distance between this two particles and if we consider the difference of wavelength ($\lambda' = 2.424979555 \times 10^{-12}$ m) of an atom ($\lambda_u = 1.331025216 \times 10^{-15}$ m) and an electron ($\lambda_c = 2.42631058 \times 10^{-12}$ m) as a function of r (distance) and if we put the mass of atom and electron in place of M as $m_u = 1.6605402 \times 10^{-27}$ kg & m represents as $m_e = 9.1093897 \times 10^{-31}$ kg respectively in the equation of Newton law of gravitation, then we get $1.716393863 \times 10^{-44}$ N and in terms of energy is $1.7163938 \times 10^{-58}$ Joule and then, mass = $m_0 = E/c^2 = 1.909745658 \times 10^{-75}$ kg.

$$E = G \frac{m_e \times m_u}{\{\lambda'\}^2} \times 10^{-14} = 1.7163938 \times 10^{-58} \text{ J} \dots\dots\dots (13)$$

Where, $\lambda' = (\lambda_c - \lambda_u) = 2.429796 \times 10^{-12}$ m, $G = 6.67259 \times 10^{-11}$ m³ kg⁻¹ sec⁻².

This energy is almost near to

$$\hbar/N_A = 1.751160297 \times 10^{-58} \text{ J} \dots\dots\dots (14)$$

And then, mass = $m_0 = E/c^2 = 1.948428595 \times 10^{-75}$ kg. This mass is 8.5177338×10^{17} times smaller than a photon's mass. Again, there is small difference of energy obtained from the equation (13) & (14) and the ratio is,

Ratio = $1.7163938 \times 10^{-58}$ Joule / $1.751160297 \times 10^{-58}$ J = 0.980146593 = $(0.990023531)^2$. Let, $\beta = 0.99$, a deformation of shape of the particle during reaction. Because at excited state matter will change its shape and as a result it may define as deformed shapes or the deformation parameter of the particle. Then we can arrange the equation (ii) in the form of:

$$E = G \frac{m_e \times m_u}{(\lambda' \beta)^2} \times 10^{-14} = 1.751160297 \times 10^{-58} \text{ J} \dots\dots\dots (15)$$

The equation (14) & (15) be treated as the energy of a graviton (E_g) and mass, g . Therefore, we can write the equation (14),

Energy of a graviton,
 $E_g = \hbar/N_A = 1.751160297 \times 10^{-58} \text{ J} \dots\dots\dots (16)$

And mass a graviton,
 $g = \hbar/N_A c^2 = 1.948428595 \times 10^{-75} \text{ kg} \dots\dots\dots (17)$

And for equation (15):

$$E_g = G \frac{m_e \times m_u}{(\lambda' \beta)^2} \times 10^{-14} = 1.751160297 \times 10^{-58} \text{ J} \dots\dots\dots (18)$$

$$g = G \frac{(m_e \times m_u)}{(\lambda' \beta)^2 \times c^2} \times 10^{-14} \text{ kg} = 1.948428595 \times 10^{-75} \text{ kg} \dots\dots\dots (19)$$

Now, from this equation (17) & (19) we get,

$$\frac{\hbar}{N_A g^2} = G \frac{(m_e \times m_u)}{(\lambda' \beta)^2 \times c^2} \times 10^{-14}$$

$$\text{Or, } \hbar = G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \times N_A \times 10^{-14} \text{ J-s}$$

$$\text{Or, } \hbar = G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \times (6.0221367 \times 10^{23}) \text{ J-s} \dots\dots\dots (20)$$

$$\text{Or, } \hbar = G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \times g_{\eta} \text{ J-s [Let, } (6.0221367 \times 10^{23}) = g_{\eta}]$$

$$\text{Or, } \hbar = G g_{\eta} \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \text{ J-s} \dots\dots\dots (21)$$

The R. H. S. of the equation (21) is expressing that in what way the gravitational constant (G), number of gravitons ($g_{\eta} = 6.0221367 \times 10^9$ gravitons) in a field of atom – electron interaction are functioning and showing the energy of Planck constant (\hbar) in L. H. S.

Non-observable or silent effect of the equation (21):

$$\hbar = g_{\eta} G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \text{ J-s}$$

During emission of energy from the equation, $E = mc^2$ or $E = h\nu$, the effect of g_{η} and G to be neutralized. We have the equation,

$$E = h\nu = 2\pi\hbar\nu = 2\pi g_{\eta} G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} \nu \dots\dots\dots (22), \text{ but } \nu = m_0 c^2 / \hbar,$$

$$\text{so, } E = 2\pi g_{\eta} G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} m_0 c^2 = 1.000000002 \times m_0 c^2 \dots\dots\dots (23)$$

Because, $g_{\eta} = 6,0221367 \times 10^9$ gravitons. Due to non-variation of the equation (23), we may use $E = mc^2$ normally. For this reason, we cannot observe the effect of graviton and gravitational constant during emission of energy. But, we need to know the internal functions of m in Einstein's equation when c^2 will act on it. To reach that point we should think about matter on the subject determination of unknown weight of radioactive element.

3. Determination of Unknown Weight of Radioactive Elements

The intensity of radioactive elements has so far been considered in terms of atoms which disintegrate per unit time. In radioactivity, the slandered unit is the "Curie" which is 3.7×10^{10} disintegration / sec. we can calculate the unknown weight (W) of radioactive element by using Curie equation in which one Curie unit and disintegration constant has taken important part.

$$W = \frac{m_0 Ci}{N_A \lambda} \dots\dots\dots (24)$$

For example [4]:

Mass of radioactive element RaB (Pb^{214}) = $m_0 = 214$. Disintegration constant, $\lambda = 4.31 \times 10^{-4}$ / sec, Avogadro number = $N_A = 6.0221357 \times 10^{23}$, one Curie = $Ci = 3.7 \times 10^{10}$ disintegration / sec. then we get unknown weight of radioactive element as,

$$W = \frac{m_0 Ci}{N_A \lambda} = \frac{214 \times 3.7 \times 10^{10} \text{ dis./sec}}{6.0221367 \times 10^{23} \times 4.31 \times 10^{-4} / \text{sec}} = 3.05616519 \times 10^{-8} \text{ gm} \dots\dots\dots (25)$$

How we can reach to this result to use the equation (7) or (8) for unification?

We can calculate the above amount of unknown weight (25) by taking the mass (σ') into the following imaginative equation.

If W is directly proportional to the square root of mass (σ') emitted from the particle m_0 and inversely proportional to the disintegration constant λ , then

$$W \propto \frac{\sqrt{\sigma'}}{\lambda} \text{ or } W = \frac{N_A}{c} \times \frac{\sqrt{\sigma'}}{\lambda} \dots\dots\dots (26)$$

Here, $N_A / c =$ Avogadro number / velocity of light = Constant. And where,

This equation (26) based on assumption only, but gives result same to Curie equation (25).

$$\sigma' = \frac{m_e (m_0^2 - m_1^2)}{\pi^2 N_A m_1^2} \text{ } (\sigma' = \text{Mass of populated photons}) \dots\dots\dots (27)$$

Placing the mass of radioactive element RaB (Pb^{214}) = $m_0 = 214$ in the equation (7), we get,

$\sigma' = 4.31375778 \times 10^{-49}$ gm, where, $m_e =$ mass of electron, $m_1 =$ mass of Alfa particle, $\pi = 3.141592654$. Again on putting the mass of populated photons (σ') and $\lambda = 4.31 \times 10^{-4}$ / sec, value of N_A / c in the equation (81), we get,

$$W = \frac{N_A \sqrt{\sigma'}}{c \lambda} = 3.061121126 \times 10^{-8} \text{ gm} \dots\dots\dots (27)$$

Result obtained from the equation (25) and (27) is almost same, for the small difference in decimal fraction observed at excited state due to deformation of particle, in this state, the value of π will change according to mass of particle, which not included here in this stapes. Before discussing the actual value of deforming π with respect to mass of particle,

we will classify the matter and the accurate values of these two equations (24) & (27) mentioned in Table – 1. When two results are same, then we can define matter (m_0) in better way.

$$\text{When, } W = \frac{m_0 c_i}{N_A \lambda} \text{ \& } W = \frac{N_A \sqrt{\sigma'}}{c \lambda} \text{ then, } \frac{m_0 c_i}{N_A \lambda} = \frac{N_A \sqrt{\sigma'}}{c \lambda} \text{ or } m_0 = \frac{N_A^2 \sqrt{\sigma'}}{C_i c} \dots\dots\dots (28)$$

The equation (28) is indicating that m_0 directly proportional to $\sqrt{\sigma'}$ (root of populated photon's mass). Because, $N_A^2/C_i c$ is constant. This equation (28) is expressing the real definition of matter. We know the definition of matter which has mass, volume, occupying space, momentum etc. But the equation (28) is showing different picture for actual reason of matter. Therefore, we can put this mass (m_0) in the Einstein equation, $E = m_0 c^2$ and the new energy equation will:

$$m_0 c^2 = \frac{N_A^2 \sqrt{\sigma'}}{C_i c} \times c^2 \text{ or, } E = \frac{N_A^2 \sqrt{\sigma'} c}{C_i} \dots\dots\dots (29)$$

$$\text{When, } m_0 c^2 = \frac{N_A^2 \sqrt{\sigma'} c}{C_i} \text{ then, } m_0 c = \frac{N_A^2 \sqrt{\sigma'}}{C_i} \dots\dots\dots (30)$$

The equation (29) expressing the internal functions of Einstein equation $E = m_0 c^2$ with velocity of light & divided by one Curie unit and the equation (30) indicating the momentum of particle.

To determine the accurate value of particle, we need to find the value of Pi at excited state. If it is possible to know the exact value of σ' , then on putting this value in equation (7), the value of Pi at excited state to be obtained. To determine this result, we can arrange the equation (30),

$$\sigma' = \frac{m_0^2 C_i^2 c^2}{N_A^4} \dots\dots\dots (31) \text{ The equation (7) is, } \sigma' = \frac{m_e (m_0^2 - m_i^2)}{\pi^2 N_A m_i^2}$$

Equating both the equations (31) & (7), we will get the value of Pi at excited state is:

$$\pi_e = \frac{\sqrt{N_A^3 m_e (m_0^2 - m_i^2)}}{m_0 m_i C_i c} \text{ or, } \pi_e = 3.152970491 \times \frac{\sqrt{(m_0^2 - m_i^2)}}{m_0} \dots\dots\dots (32)$$

In this regards we need to discuss about the Pi that what it is and why we call the value of Pi at excited state? Normally, the value of $\pi = 3.141592654$ used if any field of science and measurement. But for deformation of particle at excited state, the circumference and radius will not stay which the particle was at rest position.

Archimedes Syracus, the greatest mathematician of antiquity, rigorously established the equivalence of the two ratios in his treatise "Measurement of a Circle". He also calculate a value for Pi based on mathematical principles rather than on direct measurement of a circle's circumference, area and diameter. What Archimedes did was to inscribe and circumscribe regular polygons (polygons whose sides are all the same length) on a circle assumed to have a diameter of one unit and to consider the polygons respective perimeters as lower and upper bounds for possible values of the circumference of the circle which is numerically equal to Pi. According to his method for estimating Pi relied on inscribed and circumscribed regular polygons. The perimeter of the inscribed and circumscribed polygons served respectively as lower and upper bounds for the value of Pi. The sine and tangent functions can be used to calculate the polygons perimeters, as is shown here, but

Archimedes had to develop equivalent relations based on geometric constructions using 96-sided polygons [5].

On the basis of this, we observed that perimeter of circumscribed polygon or perimeter of inscribed polygon, in both the cases, the value of P_i and $P_c (= n \tan 180^\circ / n)$ has been changed (due to number, n) and showing that the numerical value are near to the value of Pi. When the shape of the circle changes (whatever by P_c or P_i), the value of π changes, this deformation of circle is important to the field of science. Because, the shape of the particle changes at excited state of it. As a result, energy absorbed or emits during that moment the normal value of π to be changed into its working value, the change of which will depend on the mass of the particle and shape of the particle to be changed accordingly on the value of deviated π . In our traditional theories, the value of π has used as 3.141592654 as a constant to the equations $\hbar = h/2\pi$, $K = 2\pi/\lambda$, $F = m^1 \times m^2 / 4\pi r^2$, etc for examples. The value of π must be constant in the case of uniform shape that is when particle will take place at rest and then, \hbar , K , F etc will more appropriate definitions for particle. so, we need to deviated value of π at excited state of matter.

Therefore, the equation (32).

$$\pi_e = 3.152970491 \times \frac{\sqrt{(m_0^2 - m_i^2)}}{m_0}$$

the equation (32) has important role at excited state of matter. Now we can symbolised π at excited state as π_e and π in normal or at rest of the particle, may denoted by π_0 in all equations in this field. If the mass of Alfa particle (m_i) become constant, then the value of Pi will be change viz with respect to the mass of particle (m_0), the emission of coagulation of photons (σ') or populated photon's mass remain same. But due to effect of one Curie and velocity of light, the equation

$$\pi_e = \frac{\sqrt{N_A^3 m_e (m_0^2 - m_i^2)}}{m_0 m_i C_i c}$$

will express the deviated Pi value at excited state of matter, the difference of this state (π_e) and at rest ($\pi_0 = 3.141592654$) is important. Because, "on the stability of rotating nuclei against fission through crevices shapes" [6]. G.Royer and F.Haddad shows that during fission process even at very high spins, an intermediate angular moment, the deformation energy is relatively constant up to $\beta = 0.99$ ($\beta = 0, 0.2, 0.4, 0.6, 0.8, 0.99$). where, β is the deformation parameter. In the first path, shape keeps almost spherical ends and the formation of a deep neck occurs before the elongation of the nuclear system. To get the result of $\beta = 0.99$, we can consider the value of Pi as, $\beta = 1 - (\pi_e - \pi_0)$ for heavy nucleus like uranium, thorium etc. to consider the value of 3.152970491 (refer equation – 32).

If we put $m_0 =$ mass of uranium = 235 in the equation (32), then we will get the value of π as $\pi_e = 3.152506608$, therefore,

$$\beta = 1 - (3.152506608 - 3.141592654) = 0.989086045 = 0.99.$$

in this regards we can compare β with π as,

$\beta = 0$ or $\pi_0 = 3.141592654$, (The shape of particle will take place almost spherical ends)

$\beta = 0.99$ or $\pi_e = 1 - (\pi_e - \pi_0)$, (the maximum deformation of particle (U^{235}) will be obtained).

So, it can be said that other deviated value of Pi will take place between 0.2 to 0.8 during excitation of matter. Therefore, we can write the equation of “**Unification of Physics**” in two forms as at rest and excited state of the particle sited here. the difference of energy is important in the case of liberating of energy. For example, if we putting the mass of uranium 235.1175 amu in this equation at rest and excited state, we will get the **energy of fast neutron as 1.215933 Mev**. The energy greater than 1.2 Mev is called the fast neutron.

$$\sigma'_0 = \frac{m_e (m_0^2 - m_1^2)}{\pi_0^2 N_A m_1^2} \dots\dots\dots (33) \quad \text{or,} \quad N_A \sigma'_0 = \frac{m_e (m_0^2 - m_1^2)}{\pi_0^2 N_A m_1^2}$$

(Mass of populated photons at rest) (Mass of Avogadro number)

$$\sigma'_e = \frac{m_e (m_0^2 - m_1^2)}{\pi_e^2 N_A m_1^2} \dots\dots\dots (35) \quad \text{or,} \quad N_A \sigma'_e = \frac{m_e (m_0^2 - m_1^2)}{\pi_e^2 N_A m_1^2}$$

(Mass of populated photons at excited state) (Mass of Avogadro number)

On putting the value of Pi in the equations (33) & (34) at rest and (35) & (36) at excited state, the difference between two energy, say ($N_A \sigma'_0 - N_A \sigma'_e$) must give result and we will get a new idea of internat functions of matter from the view of particle side that in what way energy is yielding from the rest after reactions. Application of the equation (35) is given here for example to determine the unknown weight of radioactive elements. We can compare the results of the Curie equation (24, group -4 in table-1) and new equation

(26, group -5 in table-1), there on variation in decimal fractions. This evidence is supporting that the equation of unification of physics is applicable to microscopic fields and the equation (26) is perfectly true. So, this (26) equation is the parallel equation of (24).

In this regards following constants considered to find the unknown weight of radioactive elements.

1. Fundamental constants:

Mass of electron (m_e) = $9.1093897 \times 10^{-28}$ gm, Avogadro number = 6.0221367×10^{23} , One Curie (Ci) = 3.7×10^{10} dis./sec, c = Velocity of light = $2.99792458 \times 10^{10}$ cm/sec, Value of Pi at rest ($\pi = \pi_0$) = 3.141592654, Mass of Alfa particle (m_1) = 4.033 amu, ϕ = Curie Photon = 2.94×10^5 photons.

2. Calculated results taken as constants:

Mass of a photon (σ) = $1.659619614 \times 10^{-54}$ gm, Energy of a photon [\bar{e}] = $9.309779229 \times 10^{-22}$ ev, $\Sigma = \sigma'/\sigma$ = Populated Photons yields from the mass m_0 to follow the equation (35), this is constant per particle based on mass. Value of Pi at excited state (π_e) = $3.152970491 \times \sqrt{(m_0^2 - m_1^2)}/m_0$. (Change of Pi depend on the mass of m_0).

The results of group - 8 in this table-1 is showing between 1/2 and 1/3 of energy of Planck constant ($\hbar/[e]$) = $6.5821220 \times 10^{-16}$ ev and value of Pi at excited state shown in group (9).

DETERMINATION OF UNKNOWN WEIGHT OF RADIOACTIVE ELEMENTS BY NEW PROCESS (TABLE - 1)

Radio-active ELEMENTS AND NUCLIDE	MASS m_0	DEINTEGRATION CONSTANT λ /sec. TYPE OF DIS.	$W = \frac{m_0 Ci}{N_A \lambda}$ gm	$W = \frac{N_A \sigma'_0}{c \lambda}$ gm Eq ^l , based on assumption	$\sigma'_e = \frac{m_e (m_0^2 - m_1^2)}{\pi_e^2 N_A m_1^2}$ gm	EMITTED PHOTONS FROM THE MASS m_0 $\Sigma m_0 = \sigma'/\sigma =$ mass of populated photons	$E = \sigma' \bar{e} \times \bar{e}$ \bar{e} = energy of a photon ev	VALUE OF π_e AT EXCITED STATE
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Ra ${}_{88}\text{Ra}^{226}$	226	1.374×10^{-11} α	1.0105849	1.0105849	4.7781×10^{-49}	2.87906×10^3	2.6803×10^{-16}	3.152468421
Ac ${}_{88}\text{Ac}^{227}$	227	1.08×10^{-9} α, β	0.0129137	0.0129137	4.8205×10^{-49}	2.90459×10^3	2.7041×10^{-16}	3.152472853
MsTh ₁ ${}_{88}\text{Ra}^{228}$	228	3.81×10^{-9} β	3.6767×10^{-3}	3.6767×10^{-3}	4.8630×10^{-49}	2.93024×10^3	2.7279×10^{-16}	3.152477192
Th ${}_{90}\text{Th}^{232}$	232	1.56×10^{-18} α	9.1372×10^6	9.1372×10^6	5.0352×10^{-49}	3.03395×10^3	2.8245×10^{-16}	3.152494057
$\Phi = \frac{3.7 \times 10^{10}}{4\pi (100)^2} = 2.94436644 \times 10^3$ (Reference value).						2.7411×10^{-16}	3.141592654	

We discussed about the equation (28) that how the mass of a particle acts and defined Einstein's equation $E = mc^2$. But we do not know the effect of the particle graviton and gravitational constant. The equation (23) is expressing the side effect of mc^2 , here we observed that the action of graviton and gravitational constant works as silent or non - observable mode. We did not include σ' , the mass of populated photons (7) or in the form of σ'_0 or σ'_e when the particle m at rest or at excited state which comes out after interacting with the mass of electron, alpha particle showing in the equation (33) & (35). The Avogadro number of σ' as $N_A \sigma'_0$ or $N_A \sigma'_e$ is expressing in the mass of Avogadro

number of populated photons in the equation (34) & (36) at rest and excited state respectively. So, we need to find the real formation of matter (m) in Einstein equation $E = mc^2$ by the action of σ' both in rest and excited state with graviton and gravitational constant.

We formed the equation,

$$E = 2\pi g_{\eta} G \frac{(m_e \times m_u)}{(\lambda' \beta)^2} v \dots\dots\dots (22)$$

From the equation De-Broglie, $v = c/\lambda$, on putting the value of v in the equation (22), we get,

$$E = 2\pi g_{\eta} G \frac{(m_e \times m_u) c}{\lambda (\lambda' \beta)^2} \dots\dots\dots (37) \text{ but, } E = \frac{N_A^2 \sqrt{\sigma'} c}{Ci} \dots\dots\dots (29)$$

For actual internal functions of m_0 is given here with effect of gravitons, gravitational constant, Planck constant, one Curie, velocity of light, with deformed condition of electron & atom. For this environment need two equations (37) and (29) while come in contact.

$$\frac{N_A^2 \sqrt{\sigma'} c}{Ci} = 2\pi g_{\eta} G \frac{(m_e \times m_u) c}{\lambda (\lambda' \beta)^2}$$

or $\lambda = 2\pi g_{\eta} G Ci \frac{(m_e \times m_u)}{N_A^2 \sqrt{\sigma'} (\lambda' \beta)^2}$ Again, $\lambda = h / m_0 c$, therefore,

$$m_0 = \frac{h N_A^2 \sqrt{\sigma'} (\lambda' \beta)^2}{2\pi g_{\eta} G (m_e \times m_u) Ci c} \dots\dots\dots (38)$$

The equation (38) is explaining the internal mechanism of m_0 perfectly interactions in the denominator with the effect of gravitons (g_{η}), G , m_e , m_u , Ci , c , and in numerator, populated photons (σ') obtained from the equation (7), difference of wavelength of electron & atom (λ') with deformation parameter (β) as a functions of $(\lambda' \beta)^2$ and Planck constant with Avogadro number² as $h \times N_A^2$. If we put this

$$\text{(A) } E = mc^2, \text{(B) } E = \frac{N_A^2 \sqrt{\sigma'} c}{Ci}, \text{(C) } E = 2\pi g_{\eta} G \frac{(m_e \times m_u) c}{\lambda (\lambda' \beta)^2}, \text{(D) } E_e = \frac{h N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2 c}{2\pi g_{\eta} G (m_e \times m_u) Ci}$$

Applications of Pt in the form of (a) $m_0 = 195.084$ gm. and (b) $m_0 = 195.084$ amu $\times 1.6605402 \times 10^{-27}$ kg = $3.239448244 \times 10^{-25}$ kg or 195.084 amu $\times 1.6605402 \times 10^{-24}$ gm = $3.239448244 \times 10^{-22}$ gm when required.

For equation (A): when, $E = mc^2$, then, $E = 2.91147 \times 10^{-8}$ Joule in M.K.S. system or $E = 0.291147$ erg in C.G.S. system.

And when, when, $m_0 = 195.084$ gm, then, $E = 1.75332755 \times 10^{23}$ erg, (when, c in centimeter/sec) or $E = 1.75332755 \times 10^{19}$ Joule (when, c in meter/sec), $c = 2.99792458 \times 10^{10}$ cm/sec in C.G.S. system.

For equation, (B):

Before determining the mass of populated photons (σ') at excited state, we need to calculate the value of π at excited state. The equation (32) is,

$$\pi_e = 3.152970491x \frac{\sqrt{(m_0^2 - m_1^2)}}{m_0}$$

When, $m_0 = 195.084$ amu and $m_1 = 4.033$ amu, then, $\pi_e = 3.15296661$ and so, to follow the equation (35), we get, $\sigma'_e = 3.560292829 \times 10^{-49}$ gm and on putting the value of σ'_e in the equation (B) when $\sigma' = \sigma'_e$, then, $E = 1.75332755 \times 10^{23}$ erg for mass 195.084 gm & in terms of atomic mass, $E = (195.084 \text{ amu} \times 1.6605402 \times 10^{-24} \text{ gm}) \times c^2 = 0.291147$ erg or 2.91147×10^{-8} Joule. There is no variation of result between the equation (A) and (B).

For equation (C):

To determine the energy, E , we need to find the value of λ for the mass $m_0 = 195.084$ amu through the equation $\lambda = h / m_0 c = 6.82245 \times 10^{-18}$ m. on putting the value of λ in the equation (C), we get, $E = 2.91147 \times 10^{-8}$ Joule, when, $G =$

$$9 \times 6.67259 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ sec}^{-2}, \lambda' = \text{difference of wavelength of electron and atom} = 2.424979555 \times 10^{-12} \text{ m}, g_{\eta} =$$

m_0 in the equation $E = mc^2$, then we get internal functions of m_0 by acting c^2 and finally we can explain the energy E as:

$$E = \frac{h N_A^2 \sqrt{\sigma'} (\lambda' \beta)^2 c}{2\pi g_{\eta} G (m_e \times m_u) Ci} \dots\dots\dots (39), \text{ where, } \sigma' = \frac{m_e (m_0^2 - m_1^2)}{\pi^2 N_A m_1^2}$$

The equation (39) will function at rest and at excited state both changing π as π_0 at rest and π as π_e at excited state in the equation (33) & (35) respectively. In this condition, the mass of populated photons will change from σ' to σ'_0 and σ'_e at rest and at excited state respectively. Therefore E will change accordingly.

So, we can arrange the equation (39) for two states as:

$$E_0 = \frac{h N_A^2 \sqrt{\sigma'_0} (\lambda' \beta)^2 c}{2\pi g_{\eta} G (m_e \times m_u) Ci} \dots\dots\dots (40) \text{ and } E_e = \frac{h N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2 c}{2\pi g_{\eta} G (m_e \times m_u) Ci} \dots\dots\dots (41)$$

The value of graviton (g_{η}) in M.K.S. system, $g_{\eta} = 6.0221367 \times 10^9$ gravitons and in C.G.S. system, $g_{\eta} = 6.0221367 \times 10^{11}$ gravitons is applicable. For example, we can take an element, Platinum (At. No. - 78) of mass (m_0) is 195.084 amu for verification of the following equations:

$$\text{(A) } E = mc^2, \text{(B) } E = \frac{N_A^2 \sqrt{\sigma'} c}{Ci}, \text{(C) } E = 2\pi g_{\eta} G \frac{(m_e \times m_u) c}{\lambda (\lambda' \beta)^2}, \text{(D) } E_e = \frac{h N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2 c}{2\pi g_{\eta} G (m_e \times m_u) Ci}$$

6.0221367×10^9 gravitons, $m_e = 9.1093897 \times 10^{-31}$ kg, $m_u = 1.6605402 \times 10^{-27}$ kg, $\beta = 0.990023531$, $c =$ velocity of light = 2.99792458×10^8 m and $\pi = 3.141592654$.

For equation (D):

We find the mass of $\sigma'_e = 3.560292829 \times 10^{-49}$ gm and in C.G.S. system, $G = 6.67259 \times 10^{-8}$ cm³ gm⁻¹ sec⁻², accordingly, we should use all values of the equation (D) in C.G.S. unit and then we will get, $E = 1.75332755 \times 10^{23}$ erg for mass 195.084 gm and then for Pt - atom, 0.291147 erg.

i) Planck constant ($6.6257055 \times 10^{-27}$ erg-s):

This unit is very active and applicable to all energy fields. We can classify the Planck constant from the equation (41) that how graviton & gravitational constant played role with the σ'_e , Ci , and other particles.

$$h N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2 c = 2\pi g_{\eta} G (m_e \times m_u) Ci E_e$$

$$h = \frac{2\pi g_{\eta} G (m_e \times m_u) Ci (m_{excited} \times c^2)}{N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2 c}$$

(When, $E_e = m_{excited} \times c^2$, $m_0 \rightarrow m_{excited} = m$)

$$h = \frac{2\pi g_{\eta} G (m_e \times m_u) Ci (m c)}{N_A^2 \sqrt{\sigma'_e} (\lambda' \beta)^2} \dots\dots\dots (42)$$

$$\text{When } \sigma'_e = \frac{m_e (m_0^2 - m_1^2)}{\pi_e^2 N_A m_1^2} \text{ and } \pi_e = 3.152970491x \frac{\sqrt{(m_0^2 - m_1^2)}}{m_0}$$

For example: Mass of Thorium, Th = $232.03806 = m = m_0$, on putting this mass in the equation of finding the value of π at excited state, we get, $\pi_e = 3.152494213$. Again, on placing the value of π_e and m_0 in the equation of unification, get, $\sigma'_e = 5.036871453 \times 10^{-49}$ gm. On putting all values of the equation (42), we get, $h = 6.6246586 \times 10^{-27}$ erg. Hence, the equation (42) is showing the internal functions of Planck

constant. Where, $g_{\eta} = 6.0221367 \times 10^{11}$ gravitons and all values are in C.G.S. system.

ii) Determination of Planck constant by using one million photons:

We calculated the mass of a photon $\sigma = 1.659619614 \times 10^{-54}$ gm, accordingly the energy, $\bar{\epsilon} = 9.309779229 \times 10^{-22}$ ev.

If 10^6 photons are able to form Planck constant, then we can come to a conclusion that the equation mass of a photon and the equation of unification of physics is true. Because, this equation obtained from photon's equation through the equation of relativity of Einstein and already mentioned in a chart of equations stage by stage of formation of the equation of unification physics. According to Sommerfeld, the expression for the energy and electron moving in a region of uniform potential is given by,

$$E = \frac{h^2 k^2}{8\pi^2 m} \dots\dots\dots (43) \text{ Where, } k = \text{wave number of particle} = \frac{2\pi}{\lambda}$$

For an electron, let $m = m_e$, $\lambda = \lambda_c = 2.42631058 \times 10^{-10}$ cm (Compton wavelength of electron). Then, $K = 2.589604711 \times 10^{10}$. So the above equation brings, $E = 0.2554995364 \times 10^6$ ev. It is $1/2$ of the total energy of electron. So, for total energy of electron is:

$$E = \frac{2h^2 k^2}{8\pi^2 m} = \sum_e (\text{Total photons of electron}) \text{ or } E = \frac{h^2}{4\pi^2 m_e} = \frac{\sum_e}{k^2} \dots\dots\dots (44)$$

But Σ'_e , the residual photons emits 1000 photons at excited state and acts with quantum number ($j = l \pm 1/2$) as a function of $\sqrt{j} = \sqrt{(l \pm 1/2)}$, $l=1$ that is if we put $\sqrt{3/2} \Sigma'_e$ in place of Σ_e in R.H.S of the equation (42), then we will get,

$$\frac{\sqrt{\frac{3}{2}} \times \Sigma_e}{k^2} = \sqrt{\frac{3}{2}} \times \frac{547.8841 \times 10^{24}}{(2.589604 \times 10^{10})^2} = 1.000110 \times 10^6 \text{ photons} \dots\dots\dots (45)$$

L.H.S of this equation (45) is showing $h^2 / 4\pi^2 m_e$, in it $h/2\pi$ factor is known as the spinning of photon. Therefore, for escape of photons from an electron, the equation (45) will turn to

$$\left[\frac{\sqrt{\frac{3}{2}} \times \Sigma_e}{k^2} \right]^{1/2} = 1000.307956 \text{ photons} \dots\dots\dots (46)$$

We can omit the decimal fraction, and then we can come to the point that an electron can emit or absorb 1000 photons at excited state of it. Where we can use these 1000 photons which may come out an electron? Let us try to solve this problem to give some examples.

If 1000 electrons functioned at excited state and each electron liberates 1000 photons, then we get 10^6 photons or we will get same value from the equation – 45 which linked to Sommerfeld's equation. If we divided 10^6 photons by $\sqrt{2}$ as quantum number ($l = 0$), then we get 6.5830×10^{-16} ev which will almost equal to Planck constant ($h [e] = 6.5821220 \times 10^{-16}$ ev-s). In case of emission of photons, quantum number functioned as the $\sqrt{j} = \sqrt{(l \pm 1/2)}$, when, $l = 0$, then, we get, $\sqrt{j} = \sqrt{1/2}$, and Planck energy $h [e]$:

$$\text{Planck constant} = \frac{10^6 \text{ photons}}{\sqrt{2}} \times \bar{\epsilon} = 6.5830 \times 10^{-16} \text{ ev} \dots\dots\dots (47)$$

Where, the value of Planck constant = 6.5821×10^{-16} ev – sec. So, the mass of a photon is very important in the microscopic field of particle.

4. Conclusion

After finding of mass of a photon and graviton, it brings a new knowledge of internal functions of matter that energy liberates from matter using Einstein's equation. Again Planck constant also related to energy equation, as a result this simple theory knocked us as a lesson that all matter is made by the photons thus the universe. The equation of unification is vital one as it can explain many known and unknown phenomena from the particle to the universe. The populated photons are responsible to describe all facts. For example, we can determine the unknown weight of radioactive elements, Planck constant is composed of one million photons and electron liberates 1000 photons, if we keep this photons in a box of \AA^2 in meter with effect of quantum numbers, then this energy will almost same to Eigen's value of electron. This proves that electron emits energy from its outermost shell. Photon and graviton is interlinked in our surroundings in every reaction. All of us are floating in the ocean of photons and gravitons. The application of this theory has no end. Actually this theory is "Endless Theory of the Universe (Complete Unified Theory)". So, lot of scope of research in this single subject from particle to the universe.

References

[1] [2] C.L. Arora, Modern physics & electronics, S.Chand & Company.Ltd, Ram Nagar, New Delhi-110055, 1983, Page-52, 111.
 [3] Paul Davies, THE NEW PHYSICS, The Pitt Building, Trumpington Street, Cambridge CB2-IRP, 1989, P 71
 [4] Irving Kaplan, Nuclear Physics, 2nd edition, 1987, Narosa Publishing House, New Delhi- Madras, Bombay, Page- 253
 [5] Scientific American, V – 258, No. 2, February, 1988, P- 112.
 [6] G. Royer & F. Haddad, Journal of Physics, G: Nuclear & Particle Physics, ISSN: 0950 – 3899, Vol – 21, No. – 3, P – 21 (1995) 339-349 March 1995.