

# *In vitro* Assessment of Antibiotic Efficacy of Streptomycin Isolated from *S. griseus* Against Selected Bacteria

Manish Kumar<sup>1</sup>, Sushma<sup>2</sup>

<sup>1</sup>Ph.D. Research Scholar, Department of Biochemistry & Biochemical Engineering  
Sam Higginbottom Institute of Agriculture, Technology & Sciences  
(Deemed-to-be-University) Allahabad-211007 (U.P.) India

<sup>2</sup>Assistant Professor, Department of Biochemistry & Biochemical Engineering  
Sam Higginbottom Institute of Agriculture, Technology & Sciences  
(Deemed-to-be-University) Allahabad-211007 (U.P.) India

**Abstract:** *S. griseus* isolated from soil samples was screened for production of streptomycin by plate assay. Identification of streptomycin antibiotic was done by paper chromatography using selected strains of bacteria by measuring the zone of inhibition. Among all the bacterial strains studied, the highest zone of inhibition was measured against *E. coli* whereas lowest was measured against *S. aureus* after 24 hrs in nutrient agar media.

**Keywords:** *S. griseus*, *E. coli*, *S. aureus*, Streptomycin, Paper chromatography.

## 1. Introduction

Actinomycetes are found in fresh water, seawater, and cold and warm-blooded animals and composts. Actinomycetes produce over 6,000 chemically different antibiotics of microbial origin and continue to be an excellent source of novel compounds. Many of these natural products are commercially important medicinal compounds with a variety of therapeutic uses (Butler *et al.*, 2002). Actinomycetes are now commonly referred to as actinobacteria. Actinobacteria are neat because they tend to produce cool secondary metabolites. Many of which have been successfully isolated and turned into useful drugs and other organic chemicals. The genus *Streptomyces* is a particularly fruitful source of these compounds, a number of which have been developed as antifungals, antibiotics (antibacterials) and chemotherapeutic (anticancer) drugs. Actinomycetes, mainly *Streptomyces* species, produce tetracyclines, amino glycosides (streptomycin and its relatives), macrolides (erythromycin and its relatives), chloramphenicol, ivermectin, rifamycins and most other clinically useful antibiotics that are not beta-lactams. (Raja and Prabakarana, 2011).

*Streptomyces* is the largest genus of actinobacteria and the genus of the family streptomycetaceae (Kampfer, 2006). Over 500 species of *Streptomyces* bacteria have been described by Euzéby (2008). As with the other actinobacteria, streptomycetes are gram-positive and have genomes with high GC-content (Madigan and Martinko, 2005), which are found predominantly in soil and decaying vegetation. Most *Streptomyces* produce spores and are noted for their distinct "earthy" odor which results from production of a volatile metabolite, geosmin. *Streptomyces* are characterized by a complex secondary metabolism. They produce over two-thirds of the clinically

useful antibiotics of natural origin; e.g., neomycin (Kieser *et al.*, 2000). Actinomycetes are mostly distributed wide range of environment like terrestrial and marine. The presence of extreme environment especially at cryophilic region also reported by Raja *et al.* (2010b).

Obviously various actinomycetes, first of all the *Streptomyces* species and filamentous fungi are the most producers in respect of numbers, versatility and diversity of structures of the produced metabolites. The significance and frequency of these main types of microbes as producers of bioactive metabolites had varied significantly during the last decades. In the beginning of the antibiotic era, the fungal (*Penicillin*, *Griseofulvin*) and bacterial (*Gramicidin*) species were in the foreground of the interest, but after the discovery of streptomycin and later chloramphenicol, tetracyclines and macrolides the attention turned to the *Streptomyces* species. In the fifties and sixties, the majority (70%) of antibiotics were discovered from these species. In the next two decades the significance of the non-*Streptomyces* actinomycetales species (rare actinos) were increased, up to a 25-30% share of all antibiotics. From the early nineties the number of bioactive compounds isolated from various filamentous and other microscopic and higher fungal species had continuously increased up to more than 50% by the turn of the millennium (Kieser *et al.*, 2000). Actinomycetes are Gram-positive bacteria frequently filamentous and sporulating with DNA rich in guanine and cytosine from 57-75%. Some of their secondary metabolites have employed as useful microbial compounds (Prescott *et al.*, 2002). Example- streptomycin from *S. griseus* for treatment of tuberculosis caused by *M. tuberculosis* and the immunosuppressant drug, tacrolimus (FK506) produced by *S. tsukubaensis*. About 100 genera of actinomycetes exist in soil (Yokota, 1997) but none of them are able to produce streptomycin. The objective of the study is to evaluate

antibacterial efficacy of streptomycin secreted from *S. aureus* against selected bacteria.

## 2. Materials and Methods

Soil samples collected by sterile method from various locations nearby Mallathahalli, Bangalore, were air-dried under room temperature for about 5 days before isolation of microorganism. For this, 0.5 gm of soil samples was suspended in 9.5 ml of sterile distilled water and 0.1 ml of this was spread on four Petri plates with nutrient agar medium using  $10^{-1}$ ,  $10^{-2}$ ,  $10^{-3}$  and  $10^{-4}$  fold dilution, respectively, which were then incubated for bacteria, incubation was carried out at  $37^{\circ}\text{C}\pm 1^{\circ}\text{C}$  for 24 hrs whereas for actinomycetes, incubation was carried out at  $28\pm 1^{\circ}\text{C}$  for 3 days using starch casein nitrate agar media. The isolated bacteria were characterized on the basis of morphological and cultural characters by inoculating the selected strain onto nutrient agar media and starch casein nitrate agar media. Antimicrobial activities of isolates were tested preliminarily by cross streak method of Lemos *et al.* (1985).

### 2.1 Streptomycin production by *Streptomyces griseus*

Inoculum was prepared in starch casein nitrate broth containing soluble starch - 10, potassium phosphate dibasic - 2, potassium nitrate - 2, sodium chloride -2, casein -0.3, magnesium sulphate (7 hydrated) - 0.05, calcium carbonate - 0.02 and ferrous sulphate (7 hydrated) - 0.01 g/l ; inoculated *S. griseus*. The broth was incubated at  $28^{\circ}\text{C}\pm 1^{\circ}\text{C}$  for 7 days in an orbital shaker at 150 rpm for streptomycin production. On 8<sup>th</sup> days, production media was centrifuged at 6000 rpm for 10 minutes and supernatant was taken. Added 2 gm of activated charcoal followed by shaking the contents for 10 minutes. The content was filtered using Whatman no. 1 filter paper. The antibiotic was eluted from charcoal by spraying 100ml of methanol: HCl (9:1). The content was filtered and charcoal was removed after neutralizing the solution (pH-7) with NaOH. Equal volume of ethyl acetate was added and allowed to precipitate out followed by its centrifugation at 1000 rpm for 10 minutes and drying the organic layer in desiccator. Finally, eluted antibiotic was dissolved in phosphate buffer (0.25M and pH 7.2).

### 2.2 Identification of streptomycin

Identification of streptomycin produced was done by paper chromatography (Snell *et al.*, 1956). Ascending chromatograms on Whatman No.1 were developed containing 1 cm of solvent mixture (Acetone: Acetic acid: Water, 50:3:47). Samples were spotted 2 cm above the base of the paper, and dried thoroughly before placing in the solvent. After the solvent had migrated to the top of the paper (usually 12 cm) the chromatograms were air dried and exposed to steam to ensure adequate removal of acetic acid.

### 2.3 Measurement of zone of inhibition by agar diffusion method

Agar well diffusion method was used to check the cultures for the production of antimicrobial metabolites by Sen *et al.* (1995). 24hrs fresh cultures were diluted with pre-sterilized

normal saline and were spread by spreader on solidified nutrient agar media. Wells were prepared over the nutrient agar plates. About 50 $\mu$ l cell free supernatant of streptomycin antibiotic was added in the wells and the plates were incubated at  $37^{\circ}\text{C}$  for 24hrs. After 24 hrs, the zones of inhibition were recorded.

### 2.4 Statistical analysis

21 Petri plates were measured for statistical analysis from which 3 Petri plates for each culture were taken as triplicate. All data from this experiment were calculated as mean $\pm$ SD. Set your page as A4, width 210, height 297 and margins as follows:

## 3. Results and Discussion

Mixed culture was observed in the nutrient agar media plate of seven strains of bacteria (*B. subtilis*, *B. megaterium*, *B. macerans*, *B. brevis*, *M. luteus*, *S. aureus* and *E. coli* respectively) and one Actinomycetes (*S. griseus*) were isolated from soil samples by starch casein nitrate agar selective media. The microorganisms isolated were identified based on their morphological, culture characterization and biochemical tests. On paper chromatography, the  $R_f$  value of streptomycin was found 0.5 conforming the compound as streptomycin.





Figure 1: (a) Paper chromatography of produced streptomycin sample showing  $R_f$  value compare with marketed antibiotic sample (b) Zone of inhibition of streptomycin against *E. coli* and *S. aureus*

The produced antibiotic was checked for antibacterial activity against 7 selected pathogens. Among all the bacterial strains studied, the highest zone of inhibition ( $24.53 \pm 0.80$  mm) was measured against *E. coli* whereas lowest ( $11.51 \pm 0.6$  mm) was measured against *S. aureus* after 24 hrs in nutrient agar media (Table 1, Figure 1) followed by *B. subtilis* ( $18.10 \pm 0.82$ ), *B. megaterium* ( $19.00 \pm 1.14$ ), *B. macerans* ( $20.43 \pm 2.46$ ), *B. brevis* ( $14.66 \pm 1.50$ ) and *M. luteus* ( $16.00 \pm 5.30$ ) respectively, for streptomycin production from *S. griseus*. Streptomycin is a bactericidal antibiotic (Sharma *et al.*, 2007). It is a protein synthesis inhibitor which binds to small 16S rRNA of the 30S subunit of the bacterial ribosome, interfering with the binding of formyl-methionyl-tRNA to the 30S subunit. This leads to codon misreading, eventual inhibition of protein synthesis and ultimately death of microbial cells through a mechanism, which is still not very clear (Voet and Voet, 2004).

**Table 1:** Diameter of zone of inhibition against isolated strains\*

| Test microorganisms  | Diameter of zone of inhibition (mm) |
|----------------------|-------------------------------------|
| <i>B. subtilis</i>   | 18.10±0.82                          |
| <i>B. megaterium</i> | 19.00±1.14                          |
| <i>B. macerans</i>   | 20.43±2.46                          |
| <i>B. brevis</i>     | 14.66±1.50                          |
| <i>M. luteus</i>     | 16.00±5.30                          |
| <i>S. aureus</i>     | 11.51±0.69                          |
| <i>E. coli</i>       | 24.53±0.80                          |

\*Diameter of zone of inhibition of cultures were measured in above table as mean±SD

#### 4. Conclusion

In the present study, streptomycin was found to be most effective against *E. coli* and moderately effective against *B. macerans*, *B. megaterium*, *B. subtilis* and *M. luteus* and least against *B. brevis* and *S. aureus*.

#### 5. Acknowledgement

The authors are thankful to Best Biotek Research Labs Pvt. Ltd., Bangalore for kind help and support to carry out the piece of work.

#### References

- [1] Butler M.J., Broheim P., Jovetic S., Marinelli F., Postma P.W., Bibb M.J., 2002, Engineering of primary carbon metabolism for improved antibiotic production in *Streptomyces lividans*. *Appl. Envir. Microbiol.*, 68(10): 4731-4739.
- [2] Euzéby, J.P., 2008. Genus streptomyces. List of Prokaryotic names with Standing in Nomenclature. <http://www.bacterio.cict.fr/s/streptomyces.html>.
- [3] Kampfer, P., 2006. The Family Streptomycetaceae, Part I: Taxonomy. In: *The Prokaryotes: A Handbook on the Biology of Bacteria*, Dworkin, M. (Eds.). Springer, Berlin, pp: 538-604.
- [4] Kieser T., Bibb M.J., Buttner M.J., Chater K.F.,

Hopwood D.A., 2000, *Practical Streptomyces Genetics*. John Innes Foundation, Norwich, England, ISBN 0-7084-0623-8.

- [5] Lemos M.L., Toranzo A.E., Barja J.L., 1985. Antibiotic activity of epiphytic bacteria isolated from intertidal seaweeds. *Microbial Ecology*, 11, 149-163.
- [6] Madigan, M. and J. Martinko, 2005. *Brock Biology of Microorganisms*. 11th Edn., Prentice Hall, New Jersey, USA.
- [7] Prescott L.M., Harley J.P., Klein D.A., 2002. *Microbiology* (5<sup>th</sup> edition) W. C. Brown Publishers, 546.
- [8] Raja A., Prabakaran P., Gajalakshmi P., Rahman A.H., 2010b, A population study of psychrophilic actinomycetes isolated from rothang hill-manali soil sample. *J. Pure Applied Microbiol.*, 4: 847-851.
- [9] Raja A. and Prabakaran P., 2011. Actinomycetes and drug-an overview. *American Journal of Drug discovery and Development*, 1(2): 75-84.
- [10] Sen K.S., Haque F.S., Pal C.S., 1995. Nutrient optimization for production of broad-spectrum antibiotics by *Streptomyces antibioticus* Str. 15.4. *Acta Microbiologica Hungarica*, 42, 155-162.
- [11] Sharma D., Cukras A.R., Rogers E.J., Southworth, D.R., Green R., 2007. Mutational analysis of S12 protein and implications for the accuracy of decoding by the ribosome. *Journal of Molecular Biology*, 374(4), 1065-1076.
- [12] Snell N., Ijichi K., Lewis J.C., 1956. Paper chromatographic identification of polypeptide Gram-positive inhibiting antibiotics. *Applied Microbiology*, 4, 13.
- [13] Schantz J.T., Ng K.W., 2004. *A manual for primary human cell culture*. World Scientific Publisher, 89.
- [14] Voet D., Voet J.G., 2004. *Biochemistry* (3<sup>rd</sup> edition). *John Wiley & Sons*, 1341.
- [15] Yokota A., 1997. Phylogenetic relationship of actinomycetes, atlas of actinomycetes. The Society for actinomycetes, 194.

#### Author Profile

**Manish Kumar**, Ph.D. research scholar in the Department of Biochemistry & Biochemical Engineering, JSBB, SHIATS, Allahabad and received M.Phil degree from Singhania University, Rajasthan.

**Dr. (Mrs.) Sushma**, presently working as faculty in the Department of Biochemistry & Biochemical Engineering, JSBB, SHIATS, Allahabad, received her M.Sc. degree from GBPUA&T udham singh nagar and Ph.D. (Biochemistry) from Indian Veterinary Research Institute (IVRI), Bareilly in 1987 and 1992 respectively.