Development and Validation New Analytical Methods of Levofloxacin in IV Infusions by UV-Visible spectrophotometric Methods and Determine Assay, % Purity and Stability in Three Marketed Brands

Swapna .G¹, Deepika .G²

^{1, 2}Assistant Professor, Department of pharmaceutical Analysis & Quality Assurance, Nirmala College of Pharmacy, Atmakuru, Mangalagiri, Guntur-522503, India

Abstract: The present study was undertaken to develop and validate a simple, accurate, precise, reproducible and cost effective UV-Visible spectrophotometric method for the estimation of levofloxacin in bulk and pharmaceutical formulation in three different Brands of cipla lkem, pdpl throught the experiment the solvent used for uv method was water and the absorption spectra was carried out at 288nm and for visible method reagent used is 2,4 DNP in methanol and determination carried out at 510 nm. The concentration 'range is 2-10µg/ml. The method was shown linear in the mentioned concentrations with correlation coefficient of $\mathbb{R}^20.9999$. The %purity of levofloxacin is 97.4%-115.2%. The percent relative standard deviation (RSD %) of method precision and intermediate precision is<2. The recovery values forlevofloxacin is 99.92-100.33%. The relative standard deviation of six replicates of assay was less than 2%. The limit of detection and limit of quantification was $0.087\mu g/mL$ and $0.164\mu g/mL$. The percent relative standard deviation of robustness and ruggedness of the method was 0.136 - 0.213%. The Assay values of levofloxacin are 405.9mg-495.5mg. The stability is found tobe good in refrigerator samples than bench top assay samples. Hence, proposed method was precise, accurate and cost effective. This method could be applicable for quantitative determination of the bulk drug as well as dosage formulation.

Keywords: Levofloxacin, UV, visible, Assay, Purity, Stability

1. Introduction

Levofloxacin is a broad spectrum second generation fluoroquinolone antibacterial agent, structurally related to nalidixic acid 1. Its spectrum of activity includes most strains of bacterial pathogens responsible for respiratory, urinary tract, gastrointestinal, and abdominal infections, including gram negative, gram positive and some atypical bacterial pathogens 2-9. Levofloxacin stops multiplication of bacteria by preventing the reproduction and repair of their genetic material 10. Literature survey revealed that few analytical methods are available for the individual estimation of levofloxacin by HPTLC 11-12, by HPLC 13-14 and by conductometry 15. Recently some UV spectrophotometric methods were reported for the estimation of levofloxacin using various solvent such as 0.1M HCl 16, 100% methanol 17, acetonitrile 18 and the mixer of water, methanol and acetonitrile 19. But single estimation of this drug with 0.1M sodium hydroxide as solvent has not been reported in bulk and in pharmaceutical formulation. Thus, the aim of the present work was to develop and validate a simple, reproducible and economic analytical method to estimate levofloxacin in routine analysis.

2. Materials and Method

Materials: Pure Standard of levofloxacin in the form of levofloxacin hemihydrates powder was received as a kind gift from cistron pharmaceuticals ltd. Which was used as reference standard. The commercial levofloxacin dosage

forms used were levofloxacin 500mg/ml purchased from local market, the brands were cipla, alkem, pdpl.

Instrument: An Elico double beam UV-visible spectrophotometer SL 164 with 1 cm matched quartz cells was used for all spectral and absorbance measurements.

Method Development

1.Determination of \lambdamax: The standard stock solution of 100 µg/mL of levofloxacin was prepared by weighing 100 mg of the drug, taken in 100 mL volumetric flask and diluted with water in uv method ,and 2,4 DNP in ethanol in visible method. By appropriate dilution of standard stock solutions with diluent, different solutions containing different concentration (2, 4, 6,8,10 µg/mL) of levofloxacin were scanned in the range of 200-800 nm to determine the wavelength of maximum absorbance. Levofloxacin has shown maximum absorption at 288 nm in UV region and 510nm in visible region.

Method Validation: The proposed method was validated for different parameters like linearity, precision, accuracy, specificity, robustness, LOD, LOQ and assay,%purity and stability.

2. Linearity: The linearity was determined by plotting concentration against corresponding absorbance. Standard stock solutions, 100μ g/mL were further diluted to obtain 2μ g/mL-10g/mL solutions. The calibration curves were

constructed by plotting absorbance versus concentration and the regression equations were calculated.

3. Precision: The system precision is a measure of the method variability. It was determined by performing three replicate analyses of the same working solution. Precision of the method was demonstrated by intraday and interday variation studies. method was determined by preparing the samples of the same batch in nine determinations with three concentrations (2, 4, 6 μ g/mL) and three replicate (n=3) each on same day i.e. zero hour, fourth hour and eighth hour. The percentage RSD of the results was used to evaluate the method precision. The interday precision was determined by assaying the samples in triplicate (n=3) per day for consecutive 3days.

4. Accuracy: Accuracy of the method was calculated by recovery studies at three levels (80%, 100% and 120%) by standard addition method. An accurately weighed tablet powder equivalent to 10mg of levofloxacin was diluted with water to make100 μ g/mL and was sonicated for 20 minutes.water in uv method and 2,4 DNP reagent in 0.1 N Hcl is added in visible method. The solution was then filtered through a Whatmann filter paper (No. 41). From this solution, 1mL was transferred to three 10 mL volumetric flasks and 0.8 mL (Flask 1), 1 mL (Flask 2), and 1.2 mL (Flask 3) of stock solution of API was added and then it was made up to the mark with diluent to make them 80%, 100% and 120% spiking .

5. Specificity in the presence of excipients: The specificity test was carried out using only excipients. Spectra for placebo granules, blank and sample were measured and compared. The sample solution was kept in the oven (600C) and under the UV lamp (254 nm) for 72h in order to verify that none of the degradation products interfered with the quantification of the drug.

6. Robustness: The robustness of an analytical procedure is the measure of its capacity to remain unaffected by small but deliberate variations in method parameters and provides an indication of its reliability during normal usage. It was determined by carrying out the analysis by two analysts at two different absorption maxima ± 2 nm of 288nm and 510 nm.

The absorbance was measured and assay was calculated for three times.

7. Limit of detection (LOD) and limit of quantitation (LOQ): LOD and LOQ were calculated from the data obtained from the linearity studies. The slope of the linearity plot was determined. For each of the replicate determinations standard deviation (SD) of the responses was calculated. From these values, the limit of detection and limit of quantitation were determined on the basis of standard deviation and slope of the regression equation.

Assay of levofloxacin IV Infusion:

To analyze the concentration of levofloxacin injection, 2mL of levofloxacin infusion (which contain 5 mg mL-1) was transferred in 100ml volumetric flask and was diluted with water in uv method and 2,4 DNP in methanol in 0.1 N Hcl in visible method. This solution was further diluted to get final concentration of $10\mu g/mL$ of levofloxacin. The % assay of the drug was calculated. All determinations were conducted by thrice time.

Statistical analysis: The results were expressed as mean \pm SD. Some results were expressed as %RSD.

3. Results and Discussion

The method discussed in the present work provides a convenient and accurate way for analysis of levofloxacin. The different concentrations of 2, 4, 6, 8, 10 μ g/mL were scanned and the wavelength of maximum absorption was found a 288 nm (**Figure 1**).



Figure 1: UV Spectrum of Levofloxacin (λmax)

The drug obeyed the Beer's law with the concentration range $2-10\mu$ g/mL with R2 value 0.999 and represented excellent linear relationship of the newly developed method. The LOD and LOQ of the developed method were determined by injecting progressively low concentrations of the standard solutions for 6 times and the values of LOD and LOQ were found to be as 0.087μ g/mL and 0.164μ g/mL respectively.

The precision of the proposed method was checked by intra-day and inter-day repeatability of responses which confirmed adequate sample stability and method reliability over 24 h periods where RSD% amongst responses was found as < 2% (**Table 1**).

Table 1: Intra-Day and In	ter-D	ay Precision	of Assay	UV-
Visible Sp	ectro	photometer		

Inter-day precision					
Concentration ($\mu g/mL$) A		Absorbance	SD	% RSD	
0 hour		4 hour		8 hour	
2	0.145	0.143	0.144	0.001	0.694
4	0.289	0.285	0.284	0.003	0.925
6	0.428	0.426	0.424	0.002	0.469
Intra-day precision					
2	0.147	0.146	0.145	0.001	0.685
4	0.292	0.291	0.288	0.002	0.711
6	0.431	0.429	0.425	0.003	0.713

The accuracy was evaluated at three different concentrations which were conducted in successive

analysis (n = 3) using the proposed method and the value was expressed as percentage of recovery between the mean concentrations of recovered and injected concentration of the drug. The average recoveries were found to be as 99.96%, 100.33% and 99.92% for the concentration levels of 80%, 100% and 120% respectively (**Table 2**).

% recover y	con c	Drug adde d	Dru g foun d	%recover y	Avg.recover y	% rsd
80	10	8	7.99	99.8	99.96	0.26 0
80	10	8	8.02	100.25	99.96	0.26 0
80	10	8	7.98	99.75	99.96	0.26 0
100	10	10	10.0 1	100.1	100.3	0.25 9
100	10	10	10.0 5	100.5	100.3	0.25 9
100	10	10	10.0 4	100.4	100.3	0.25 9
120	10	12	12.0 1	100.8	99.92	0.16 9
120	10	12	11.9 9	99.92	99.92	0.16 9
120	10	12	11.9 7	99.75	99.92	0.16 9

 Table 2: Determination of Accuracy of Levofloxacin by UV-Visible Spectrophotometer (N=3)

The assays were validated by means of the analysis of variance. Levofloxacin content in pharmaceutical dosage form in IV injection was determined by these proposed methods which were in good agreement with the label claims with RSD 0.01% for injectable dosage form (**Table 3, 4**). The %RSD was found in the range of 0.136 – 0.213% for robustness and ruggedness.

Table 3: UV Method Assay, % Purity

S. No	Brand Name	%Purity	Assay		
1	Alkem	99.42%	482.8mg		
2	Cipla	99.6%	495.5mg		
3	pdpl	100%	497.48mg		

Table 4: Visible Method Assay, % Purity

S. No	Brand Name	%Purity	Assay
1 2 3	Alkem Cipla pdpl	988% 97.4% 115.2%	405.9mg 403.5mg 409.75mg

The specificity of the analytical method was proved by comparing the spectra of placebo and degradation product of sample solution with that of accuracy sample.stability was found to be good for refregirator assay sample with % RSD<2. All experimental results were within the range of the acceptability which indicated that the developed method was sensitive enough and accurate for quantitative analysis of levofloxacin. Therefore, the method was applied for quantitative analysis of levofloxacin in bulk and pharmaceutical dosage form.

4. Conclusion

This UV-visible spectrophotometric technique was quite simple, accurate, precise, reproducible and sensitive. The UV-visible method has been developed for quantification of levofloxacin in pharmaceutical dosage forms. Levofloxacin of the brand PDPL showed good % purity and Assay, The validation procedure confirms that this is a workable method for their quantification in the raw material and also in the formulations.

References

- A.P. William: Antimicrobial agent: Goodman & Gilman's The Pharmacological Basis of Therapeutics, New York, Edition 10th 2001:1183
- [2] Bertino, J.; Fish, D. Clinical Therapeutics 2000; 22: 798.
- [3] Gomes GC, Salgado HRN: Validation of UV Spectrophotometric method for determination of Lomefloxacin in Pharmaceutical Dosage Form. Acta Farm Bonaerense 2005; 24:406-408.
- [4] Ross DL, Riley CM: Antibacterial effect of nalidixic acid derivatives. International Journal of Pharmacy 1990; 63: 237-240.
- [5] British Pharmacopoeia. Vol.1, 2009.
- [6] United States Pharmacopoeia. Edition 26th, 2009: 2439.
- [7] Indian Pharmacopoeia. Govt. Of India, Ministry of Health and Family Welfare, The Indian Pharmacopoiea Commission, Ghaziabad, Vol. 2, 2010:1008.
- [8] RK Maheshwari, SC Chaturvedi, NK Jain: Novel spectrophotometric estimation of some poorly water soluble drugs using hydrotropic solubilizing agents. Indian Journal of Pharmaceutical Science 2006; 68: 195-198.
- [9] MA Mohammad, NH Zawilla, FM El-Anwar, Aly SM El-Moghazy .Chem Pharm Bull (Tokyo) 2007;55:1-6.
- [10] F. Belal, A.A. Al-Majed, A.M. Al-Obaid: Method of analysis of 4-quinole antibacterials: Elsevier Science 1999; 50: 765–786.
- [11] S.N. Meyyanathan, G.V.S. Ramasarma, B. Suresh: Analysis of levofloxacin in pharmaceutical preparations by high performance thin layer chromatography. Journal of Separation Science 2003; 26:1698–1700.
- [12] Chepurwar SB, Shirkhedkar AA, Bari SB, Fursule RA, Surana SJ:Validated HPTLC method for simultaneous estimation of levofloxacin hemihydrate and ornidazole in pharmaceutical dosage form. Journal of chromatographic science 2007; 45: 531.
- [13] Ines, M.; Santoro, R. M.; Singh, A. K.; Erika, R. M.;Hackmam, K. Journal of Pharmaceutical and Biomedical Analysis 2006; 40:179.
- [14] Neckle, U.; Joukhadar, C.; Mayer, B. X., Anal. Chim.Acta. 2002, 463, 199.
- [15] G. Altiokka, Z. Atkosar, N.O. Can: The determination of levofloxacin by flow injection analysis using UV
- [16] V.N. Desai, O.E. Afieroho, B.O.Dagunduro: A simple UV spectrophotometric method for the determination of levofloxacin in dosage formulations .Tropical Journal of Pharmaceutical Research 2011; 10:75–79.

- [17] N.M. Kassab, M.S.D. Amaral, A.K. Singh: Development and validation of UV spectrophotometric method for determination of levofloxacin in pharmaceutical dosage forms. Quim.Nova 2010; 33: 968–971.
- [18] A.A. Shirkhedkar, S.J. Surana: Quantitative determination of levofloxacin hemihydrates in bulk and tablets by UV-Spectro- photometry and first order derivative methods. Pakistan Journal of Pharmaceutical Science 2009; 22:301–302.
- [19] Mahfuza Maleque, Md. Raquibul Hasan, Farhad Hossen: Development and validation of a simple UV spectrophotometric method for determination of levofloxacin both in bulk and marketed dosage formulations. Journal of Pharmaceutical Analysis 2012;2(6)