

References

- [1] Shilpa S,Somashekar et al (2011), Prevalence of Anemia among adolescent girls; a one year cross sectional study. Journal of clinical and diagnostic research, may; vol 6[3], 372-377
- [2] A.K.Sinha,G.M.Singh Karki prevalence of anemia among adolescents in Biratnagar, Morang, Dist.Nepal. A International journal of pharmaceutical and biological archives 2012;3[5];1077-1081
- [3] Siddaram S M,Venkatesh G. M. International journal of biological and medical research .2011; 2[4]; 22-924
- [4] Kaur S, Deshmukh PR, Garg BS. (2006), Epidemiological correlates of nutritional anemia in adolescent girls of rural Wardha. *Indian J Community Med*; 31:255-58.
- [5] SHINDE, M., & ANJUM, S. (2007). Introduction to Research In Nursing.*Sneha Publication India (Dombivili)*.
- [6] Kishore J.(2006), National Health Programs of India. 6th ed.New Delhi: Century Publications; 82-84.
- [7] Bhudhagaonkar, J., & Shinde, M. (2014). Impact of Structured Education Regarding Menstrual Hygiene Practices among Adolescent Girls. *International Journal of Science and Research (IJSR)*, 3(5), 244-252.
- [8] Kaur S, P.R. Deshmukh, B.S.Garg. (2006), Epidemiological correlates of Nutritional Anemia in Adolescent girls of rural Wardha. *Indian Journal of Community Medicine*.;31:4.
- [9] Agrawal KN,Agrawal DK,Sarma A,Sarma K,Prasad K,Kalita MC.et al . (2006)Prevalence of anemia in pregnant and lactating women in India *Indian J MedRes*;124:173-184
- [10] Bulliy G, Mallic G, Sethy GS, Kar SK. (2007)Hemoglobin status of non-school going adolescent girls in three districts of Orissa, India. *Int J Adolesc Med Health*; 19: 395-406.
- [11]Rajaratnam J, Abel R, Asokan JS, Jonathan P. (2000) Prevalence of anaemia among the adolescent girls of rural Tamil Nadu. *Indian Pediatric*; 37:532-36.
- [12]Singh J, Singh JV, Srivastava AK, Suryakant. (2006)Health status of the adolescent girls in the slums of Lucknow. *Indian J Community Med*; 31(2):102-03.
- [13]Gupta N, Kochar G. [2009 Feb 13]; Pervasiveness of anaemia in adolescent girls of the lower socio-economic groups of the district of Kurukshetra (Haryana). *The Internet Journal of Nutrition and Wellness* ; 7(1): URL:<http://www.ispub.com/journal>.
- [14]Verma A, Rawal VS, Kedia G, Kumar D, Chauhan J. (2004)Factors which influenced anaemia among girls of the school going age (6-18 years) from the slum of Ahmedabad city. *Indian J Community Med*; 29(1):25-26.
- [15]WHO. (1989) Preventing and controlling iron deficiency anemia through primary health care, Geneva,
- [16]Shobha S, Sharada . (2003) Efficacy of twice weekly iron supplementation in anemic adolescent girls. *Indian Pediatric* ; 40: 1186-90.
- [17]Pathak P, Singh P, Kapil U, Raghuvanshi RS, (2003), Prevalence of iron,vitamin A and iodine deficiencies amongst adolescent pregnant mothers. *Indian J Paediatr*; 70: 299-301.
- [18]World Health Organization. Programming for adolescent health and development. WHO Tech Rep Ser No. 886.

Author Profile

Professor and HoD, Department of Community Health Nursing
B.L.D.E.A'S Shri. B.M.Patil Institute of Nursing Sciences Bijapur-586103

Associate Professor, Department of Medical Surgical Nursing
B.L.D.E.A'S Shri. B.M.Patil Institute of Nursing Sciences Bijapur-586103