

Effect of Four Herbal Seeds on Blood Parameters in Turkey Poults

Darshana B. Bhaisare¹, D. Thyagarajan²

¹Division of Poultry Science, Central Avian Research Institute, Izatnagar, Bareilly-243122, Uttar Pradesh, India

²Director of Distance Education, Madhavaram Milk Colony, Tamil Nadu Veterinary and Animal Sciences, University, Chennai-600 051, India

Abstract: A biological study using Nandanam Turkey poults (*Meleagris gallapavo*) for eight weeks duration was carried out to evaluate the effect of four herbal seeds as phytobiotics on haematological and serum biochemical parameters in turkey poults. One hundred and fifty poults were randomly subjected to five dietary treatments in a completely randomized design with basal diet (T1), 0.5% (5g/kg) level of each seeds thyme (T2), fenugreek (T3), fennel (T4) and cumin (T5) in cages. Haemoglobin (Hb) content and serum biochemical parameters such as serum total proteins, calcium, cholesterol and triglyceride were estimated using a Biosystem autoanalyzer. Haemoglobin content in overall at eighth week of age revealed significant ($P<0.05$) difference between treatments while highly significant ($P<0.01$) difference was found in total proteins with increased level recorded in fenugreek and lower level in cumin supplemented groups at eighth week of age. No significant difference in serum calcium was noticed at eighth week of age. But the overall serum cholesterol values were highly significant ($P<0.01$) with highest values recorded in fennel and lowest in cumin supplemented groups. Whereas highly significant ($P<0.01$) difference was observed in overall at eighth week of age with high levels noticed in control group then followed by cumin supplemented group compared to all other treatments.

Keywords: turkey poults, phytobiotics, thyme, fenugreek, fennel, cumin

1. Introduction

Bacterial resistance to antimicrobial drugs has become an issue of increased public concern and scientific interest during the last couple of decades. This resulted from a growing concern that the use of antimicrobial drugs in veterinary medicine and animal husbandry may compromise human health if resistant bacteria develop in animals and are transferred to humans via the food chain or the environment. Antimicrobials are being used in poultry production to treat or prevent diseases and also to promote growth in livestock and poultry grown for meat. Antibiotic as growth promoter in poultry industry has been seriously criticized by governmental policy makers and consumers as a potential harm to human health (Williams and Losa, 2001). It has been reported that the use of antibiotics as a growth promoter in chicken has caused some unwanted results in humans (Botsoglou and Fletouris, 2001). Therefore, most antibacterial performance promoters have been banned due not only to cross-resistance but also to multiple resistances (Neu, 1992). Recognizing this, the European Union has totally banned the use of growth promoting antibiotics in animal feed. On the other hand, many antibiotics are being used widely in food animals as growth promoters and to prevent and treat infectious diseases as there are no regulatory provisions are in place in many developing countries to monitor their use. The ban of antibiotics in animal feeds has consequences in the international trade of poultry meat as well because many countries adopt stringent Sanitary and Phytosanitary measures on imports of animal and plant products. It is also very important that the ways and means for the economy of the poultry production cannot be overlooked. During the last two decades, considerable research has been done in exploring suitable alternative to antibiotics as growth promoters. Herbs or products including plant extracts, essential oils or the components of the essential oils hold promise as alternatives to

antibiotics (Ocak et al., 2008). Substantial attention has been paid to medicinal herbs as replacements for antibiotic growth promoters (Ibrahim et al., 2005). There is evidence suggesting that herbs, spices, and various plant extracts have appetizing, digestion-stimulating and antimicrobial properties (Sharifiet al., 2013).

Phytobiotics are the preparations of vegetative origin containing chemical constituents, which favourably affect the microflora of the digestive system. Some of the beneficial effects of phytobiotics include increment in feed intake, stimulation of digestive secretions, stimulation of immune system, antimicrobial, antioxidant and coccidiostatic activity, pharmacological actions such as laxative, spasmolytic and astringent effects as well as prevention of flatulence and denaturizing. Under the intensive management systems, herbal extracts are already being tried as feed supplements to improve growth performance (William and Losa, 2001). Several studies indicated that these feed additives could be used in poultry ration as antifungal, antibacterial and antioxidant compounds (Khosravi et al., 2008). Even though phytobiotics provides potential alternative to the antibiotic usage, more research is needed to understand their mechanism of action. The aim of this study is to shed light on the growth stimulating effects by antimicrobial properties of four herbal seeds namely, thyme, fenugreek, fennel and cumin by conducting in-vitro and in-vivo studies.

2. Materials and Method

Biological trial was conducted at the Institute of Poultry Production and Management, Madhavaram Milk Colony, Chennai-51, located in Chennai metropolitan area of Tamil Nadu. A biological experiment of eight weeks duration was conducted in turkey poults to find out the effect of four herbal seeds namely thyme (*Thymus vulgaris*), fenugreek

(*Trigonella foenum graecum*), fennel (*Foeniculum vulgare*) and cumin (*Cuminum cyminum*) as phytobiotics on haemoglobin content and serum biochemical parameters in turkey poults. A total number of 150 straight run day old turkey poults of Nandanam turkey variety belonging to single hatch were utilised for the study. The poults were weighed individually and wing banded. Poults were randomly divided into five treatment groups with three replicates of 10 poults each. The experimental birds were reared under identical management conditions up to 8 weeks of age. Mean body weight was recorded between 0 and 8 weeks at biweekly intervals. The dietary treatments included the basal diet as control (T₁), supplemented experimental groups each receiving 5g/kg (0.5%) thyme seeds (T₂) or fenugreek seeds (T₃) or fennel seeds (T₄) or cumin seeds (T₅) added to the basal diet. These experimental feeds were fed to the respective treatment groups ad libitum from 0 to 4 and 5 to 8 weeks of age. Two milliliters of blood was collected from jugular vein of four poults in each treatment at fourth, sixth and eighth weeks of age using a 2ml syringe for estimation of haematological and serum biochemical parameters. A sample of 0.5ml of blood was taken in vial containing 0.5mg of EDTA (Ethylene Diamine Tetra Acetic Acid) as an anticoagulant. The remaining blood was kept undisturbed for two hours and the serum was separated.

2.1 Estimation of haemoglobin

Haemoglobin (Hb) content was estimated as per the Sahli's acid hematin method (Sahli, 1909).

2.2 Total Serum Protein

The total proteins level in the serum samples were estimated in A15 Biosystem auto analyzer by using commercially available AGAPPE kit based on Direct Biuret method (Gornall et al., 1949) and BCG method (Dumas et al., 1970).

2.3 Serum Calcium

The calcium in the serum samples was estimated in the A15 Biosystem auto analyser using commercially available AGAPPE kit based on the OCPC method (Kessler and Wolfman, 1964).

2.4 Serum Cholesterol and Triglycerides

Cholesterol and triglycerides in the serum were estimated using A15 Biosystem auto analyser using commercially available kits from AGAPPE diagnostics Ltd. based on CHOD-PAP methodology (Allain et al., 1974) and GPO-method (Buccolo and David, 1973) respectively.

2.5 Statistical Analysis

All the statistical analyses were performed by using SPSS software (version 7.5) as per Snedecor and Cochran (1994). The means were compared by one way ANOVA for significant differences among treatments. The percentage in data was converted into arc sin values before statistical analysis.

3. Results

The data recorded in biological experiment were grouped, analyzed statistically and presented.

3.1 Haemoglobin Content

Effect of four herbal seeds as phytobiotics feed supplementation on haemoglobin (g/dl) at fourth, sixth and eighth weeks of age in turkey poults are presented in table 1. There was a highly significant ($P < 0.01$) difference for haemoglobin content at fourth and sixth week of age with highest haemoglobin content in fenugreek and cumin supplemented group were evident. Haemoglobin content in overall and males at eighth week of age were significant ($P < 0.05$) with highest values recorded in fenugreek and cumin while insignificant difference was noted in females. Irrespective of treatment, consistent lower haemoglobin content was observed in all age groups in thyme groups.

Table 1: Effect of four herbal seeds as phytobiotics feed supplementation on mean (\pm SE) haemoglobin (g/dl) between 0 and 8 weeks of age in turkey poults (n=4)

Treatments	Age in weeks				
	4	6	8		
			Male	Female	Overall
T1	11.15 ^a \pm 0.36	10.70 ^{ab} \pm 0.31	11.40 ^{ab} \pm 0.31	11.60 \pm 0.23	11.50 ^a \pm 0.18
T2	10.31 ^b \pm 0.13	9.80 ^b \pm 0.18	10.27 ^b \pm 0.07	10.87 \pm 0.59	10.57 ^b \pm 0.30
T3	11.45 ^a \pm 0.13	11.65 ^a \pm 0.33	11.93 ^a \pm 0.47	11.47 \pm 0.27	11.70 ^a \pm 0.26
T4	10.25 ^b \pm 0.10	10.80 ^{ab} \pm 0.41	10.93 ^{ab} \pm 0.66	11.13 \pm 0.64	11.03 ^{ab} \pm 0.41
T5	11.25 ^a \pm 0.15	11.55 ^a \pm 0.33	12.17 ^a \pm 0.35	11.30 \pm 0.25	11.73 ^a \pm 0.27
F value	8.148**	5.449**	3.393*	0.437 ^{NS}	2.879*

a, b, c - means within column bearing different superscripts differ significantly ($P < 0.05$)

* - Significant ($P < 0.05$), ** - Significant ($P < 0.01$), NS - Non-significant ($P > 0.05$)

3.2 Serum Total Proteins

Influence of four herbal seeds as phytobiotics feed supplementation on serum protein (g/dl) in turkey poults is presented in table 2. Significant ($P < 0.05$) effect on serum total proteins was observed at fourth week of age with highest value spotted in thyme and lowest in fennel supplemented groups. Whereas statistically insignificant effect was noted in total proteins at sixth week of age. At eighth week of age significant ($P < 0.05$) effect was found in total proteins on overall mean and in males with compare to control. Cumin treated birds showed significantly ($P < 0.05$) lower total proteins compared to the other herbal seeds supplemented group. Control birds had intermediary values which is statistically comparable with all other treatments, in males cumin supplemented birds had significantly lower serum total proteins compared to other herbal seeds supplemented birds which had statistically similar serum total proteins. Except thyme fed birds all the other herbal seeds supplemented birds had similar serum total proteins to that of control.

Table 2: Effect of four herbal seeds as phytobiotics feed supplementation on mean (\pm SE) serum total proteins (g/dl) between 0 and 8 weeks of age in turkey poults (n=6)

Treatments	Total proteins				
	4 th week	6 th week	8 th		
			Male	Female	Overall
T1	3.47 ^{bc} \pm 0.05	3.67 \pm 0.05	3.27 ^{bc} \pm 0.03	3.41 \pm 0.24	3.34 ^{ab} \pm 0.11
T2	3.92 ^a \pm 0.05	3.59 \pm 0.05	3.69 ^a \pm 0.12	3.45 \pm 0.09	3.57 ^a \pm 0.08
T3	3.46 ^{bc} \pm 0.05	3.62 \pm 0.05	3.56 ^{ab} \pm 0.09	3.44 \pm 0.21	3.50 ^a \pm 0.11
T4	3.42 ^c \pm 0.02	3.57 \pm 0.05	3.53 ^{ab} \pm 0.09	3.54 \pm 0.13	3.53 ^a \pm 0.07
T5	3.69 ^{ab} \pm 0.11	3.63 \pm 0.05	3.18 ^c \pm 0.11	3.06 \pm 0.36	3.12 ^b \pm 0.17
F value	10.982*	0.592 ^{NS}	5.100*	0.668 ^{NS}	2.665*

a, b, c, d - means within column bearing different superscripts differ significantly (P<0.05),

* - Significant (P<0.05), ** - Significant (P<0.01), NS - Non-significant (P>0.05)

3.3 Serum Calcium

Serum calcium level variation due to feed supplementation of four herbal seeds as phytobiotics feed supplementation in turkey poults are presented in table 3. Inclusion of herbal seeds had significant (P<0.01) effect on the serum calcium level of turkey poults at fourth and sixth weeks of age. At eighth week, even though males recorded significant (P<0.05) effect for phytobiotic supplementation, the mean values of female and overall average was not altered by phytobiotic supplementation.

Table 3: Effect of four herbal seeds as phytobiotics feed supplementation on mean (\pm SE) serum calcium (mg/dl) between 0 and 8 weeks of age in turkey poults (n=6)

Treatments	Age in weeks				
	4 th	6 th	8 th		
			Male	Female	Overall
T1	5.96 ^c \pm 0.03	6.06 ^c \pm 0.02	5.18 ^b \pm 0.35	5.23 \pm 0.54	5.20 \pm 0.26
T2	7.19 ^a \pm 0.05	6.35 ^{bc} \pm 0.2	5.87 ^{ab} \pm 0.07	5.88 \pm 0.19	5.85 \pm 0.11
T3	6.87 ^b \pm 0.05	6.42 ^b \pm 0.01	6.02 ^{ab} \pm 0.16	6.31 \pm 0.04	6.16 \pm 0.10
T4	6.77 ^b \pm 0.05	6.06 ^c \pm 0.02	6.24 ^{ab} \pm 0.65	6.15 \pm 0.09	6.19 \pm 0.50
T5	7.18 ^a \pm 0.06	7.16 ^a \pm 0.05	6.94 ^a \pm 0.49	5.04 \pm 0.04	6.17 \pm 0.44
F value	104.971**	22.946**	2.100*	0.769 ^{NS}	1.198 ^{NS}

a, b, c - means within column bearing different superscripts differ significantly (P<0.05)

* - Significant (P<0.05), ** - Significant (P<0.01), NS - Non-significant (P>0.05)

3.4 Serum Cholesterol and Triglycerides

Mean serum cholesterol and triglyceride (mg/dl) in different herbal seeds supplemented groups in turkey poults is presented in table 4. Serum cholesterol level was significantly (P<0.05) affected at fourth week of age with hyper cholesterol in thyme while hypo cholesterol in fenugreek supplemented groups were seen, while no significant difference was observed at sixth weeks of age. Whereas at eighth week of age, significant (P<0.05) difference was observed both in males and in females leading to a significant (P<0.01) effect on overall values also. The serum cholesterol level was elevated in males due to fennel supplementation whereas cumin supplementation has resulted in significant (P<0.05) decrease in overall values. Apart from this, serum cholesterol level in males, females and both the sex together has not been affected by supplementation of herbal seeds. There was highly significant (P<0.01) difference in serum triglycerides level at fourth and sixth weeks of age. At fourth week except thyme supplementation, addition of other phytobiotics (P<0.05) depressed serum triglyceride level. The magnitude of depression was higher in fenugreek and fennel supplemented birds making it significantly (P<0.05) lower than cumin fed birds. At the sixth week also all the phytobiotic herbal seeds except thyme had significant (P<0.05) reducing effect of serum triglyceride in turkey poults. However at eighth week males, females and both the sex together registered significantly (P<0.05) lower triglyceride in turkey poults due to supplementation of herbal seeds. The decrease was strongly noticed in thyme, fenugreek and fennel supplemented birds making it significantly (P<0.05) lower than cumin fed birds.

Table 4: Effect four herbal seeds as phytobiotics feed supplementation on mean (\pm SE) serum cholesterol and triglycerides (mg/dl) between 0 and 8 weeks of age in turkey poults (n=6)

Treatment s	Cholesterol					Triglycerides				
	4 th week	6 th week	8 th week			4 th week	6 th week	8 th week		
			Male	Female	Overall			Male	Female	Overall
T1	126.5 ^a _b ± 3.5	120.5 ± 1.5	135.05 ^b ± 5.50	136.50 ^{ab} ± 0.5 0	136.00 ^{ab} ± 2.2 7	178.50 ^a ± 2.06	173 ^a ± 50	160.00 ^a ± 10	162.05 ^a ± 4.50	161.25 ^a ± 2.02
T2	138.0 ^a ± 50	134.0 ± 50	130.33 ^b ± 1.76	128.33 ^{ab} ± 7.5 4	129.33 ^{bc} ± 3.4 9	174.75 ^a ± 2.06	172 ^a ± 30	97.33 ^c ± 2.60	99.33 ^c ± 5.24	98.33 ^c ± 2.65
T3	114.0 ^b ± 50	124.0 ± 50	139.67 ^{ab} ± 6.3 3	141.67 ^a ± 2.73	140.67 ^{ab} ± 3.1 2	142.50 ^c ± 1.26	138 ^b ± 5	96.33 ^c ± 6.44	91.00 ^c ± 2.08	93.67 ^c ± 3.25
T4	128.0 ^a ± 50	128.0 ± 50	150.00 ^a ± 1.53	145.00 ^a ± 4.16	147.05 ^a ± 2.28	143.00 ^c ± 1.08	137 ^b ± 50	99.67 ^c ± 3.71	100.00 ^c ± 2.08	99.83 ^c ± 1.90
T5	123.0 ^a ± 50	117.0 ± 50	127.00 ^b ± 3.79	114.00 ^b ± 11.15	120.05 ^c ± 6.02	148.00 ^b ± 1.08	139 ^b ± 50	143.67 ^b ± 4.26	120.33 ^b ± 10.84	132.00 ^b ± 7.37
F value	3.354*	2.137 ^N _s	5.164*	3.332*	7.588**	127.457* *	16.408* *	42.751* *	17.630* *	39.797* *

a, b, c - means within column bearing different superscripts differ significantly ($P < 0.05$)

* - Significant ($P < 0.05$), ** - Significant ($P < 0.01$), NS - Non-significant ($P > 0.05$)

4. Discussion

4.1 Haemoglobin Content

Haemoglobin content was favourably changed at fourth, sixth and eighth week of age by supplementation of thyme seeds in feed of turkey poults. Haemoglobin content in males at eighth week of age was significantly low. These could be due to effect of some bioactive principles present in thyme seeds against haemopoietic process in the body. The results of present study was found different with Demir et al. (2008) who indicated that feeding thyme powders did not change the blood haemoglobin level but Rahimi et al. (2011) observed the improved hemoglobin concentration, but not significantly due to inclusion of 0.1% aqueous extracts of thyme in the drinking water however in broiler chickens. The statistical analysis revealed highly significant ($P < 0.01$) difference for haemoglobin content at fourth and sixth week of age with higher haemoglobin content in fenugreek supplemented group. Haemoglobin content in overall and males at eighth week of age were significantly influenced by fenugreek supplemented groups with increased level recorded in fenugreek. These results indicated that certain bioactive principles in fenugreek seeds have positive effect on haemoglobin level. Haemoglobin content at fourth, sixth and eighth week of age was positively influenced by addition of fennel seeds at 0.5% level in feed and also in males at eighth week of age. In fennel supplemented groups lesser values were observed compared to control. This result suggests that these plants may contain substances which involved in the derangement of the haemopoietic process. In contrary with present findings, Abdullah and Rabia (2009) who observed significant ($P < 0.05$) improvement in haemoglobin in broiler chickens fed fennel seeds as compared to control. Higher haemoglobin content at fourth, sixth and eighth week of age in cumin supplemented group was evident. The results was in disagreement with Al-Kassi (2010) who showed

that broiler chicks fed with 1.5% cumin powder developed a significant ($P < 0.05$) decrease in the levels of haemoglobin. Further AL-Kassie et al. (2011) revealed lower level of haemoglobin in broiler which fed with cumin powder at 0.25, 0.50, 0.75 and 1% level.

4.2 Serum Total Proteins

Thyme supplementation elevated serum total proteins level at fourth week and in males at eighth week. The overall serum total proteins of thyme fed birds were insignificantly higher than control at eighth week. In accordance with present results Toghyani et al. (2011) reported that 5g/kg thyme powder feed supplementation significantly ($P < 0.05$) increased serum total proteins and globulin concentration. Addition of fenugreek has not resulted in any change in total proteins of turkey poults at 4th, 6th, 8th weeks of age. Similar to the findings of this experiment, Abbas (2010) has also reported no significant effect on total proteins in diet of broiler containing 3g/kg fenugreek seeds. Addition of fennel seeds in the turkey feed has no influence on serum total proteins level up to eighth week of age. No previous literature could be cited to explain the findings of this experiment. Inclusion of cumin seeds has in no way connected with a change in serum total proteins of turkey poults up to eight weeks of age. However, Shabaan (2012) recorded lower values of total proteins in an experiment containing cumin seeds at 0.15 and 0.30% in diet of sbroilers. But in contrary with present findings Elagib et al. (2012) found that in broiler chick fed with diets containing cumin showed increase in total proteins fraction of the blood proteins.

4.3 Serum Calcium

Perusal of literature has indicated that the research on supplementation of phytobiotics on serum calcium level has not been studied adequately as evidenced by scanty literature. Inclusion of thyme seeds elevated serum

calcium level at fourth week of age in turkey poult. However during the rest of the period (sixth and eighth week of age), the numerical increase observed in this group was not statistically significant.

Feeding of turkey poult with fenugreek seeds has significantly ($P<0.05$) enhanced the serum calcium level at four and six weeks of age. However the increase at eighth week was statistically not significant. The turkey poult receiving fennel seeds in feed recorded significantly ($P<0.05$) higher serum calcium level at fourth week and same value at sixth week of age compared to control. Increase in serum calcium level at eighth week was not significantly higher than control. The addition of cumin seeds in the feed of turkey poult has resulted in increased level of serum calcium at four and six weeks of age. At eighth week even though males recorded significantly ($P<0.05$) higher serum calcium in males, females recorded insignificantly lower values. The overall serum calcium level at eighth week compared to control was numerically higher but statistically significant.

4.4 Serum Cholesterol and Triglycerides

It was observed from the study that supplementation of thyme seeds has no effect on serum cholesterol level. However earlier workers (Isa, (2011) at 0.02 and 0.04%; Navid and Nezhady, (2011) at 1.5% and Toghyani et al. (2011) at 0.5%) recorded significant decrease in serum cholesterol level due to thyme supplementation. High levels of fibers of these seeds can increase the excretion of bile and this can decrease the cholesterol level of blood (Zargari, 2001). Even though thyme supplementation had no significant effect on serum triglycerides during four and six weeks of age, significant decrease was observed at eighth week in overall and sex separate values. The results was supported by Isa (2011) who showed that broilers fed diet supplemented with thyme powder at 0.02 and 0.04% up to 6 weeks resulted in significant reduction ($P<0.05$) in the levels of serum triglycerides. The main reason is these substances have effect on cholesterol and triglycerides and decrease these harmful parameters in blood serum. The reduction in triglyceride and cholesterol level noticed by thyme has been attributed to the lowering effect of thymol or carvacrol on HMG-Co A reductase the rate limiting enzyme of cholesterol synthesis (Lee et al., 2003). Similar to thyme supplementation, the addition of fenugreek seeds in the diets of turkey poult also showed no significant change in serum cholesterol level during entire experiment period. These results often related to the mode of action of fenugreek in bird metabolism which includes competition with cholesterol at binding sites or interference with the cholesterol biosynthesis in the liver. Hypocholesterolaemic effects of fenugreek are owing to increased conversion of hepatic cholesterol to bile salts due to loss in the faeces of complexes of these substances with fenugreek fiber and saponins (Lee et al., 2003). However Abbas (2010) ascertained the effect of fenugreek seeds on broiler performance at the rate of 3g per kg of diet and showed significant ($P<0.05$) reduction in serum cholesterol level. Addition of fenugreek to the diets of turkey poult revealed the decreasing trend in serum triglyceride level compared to control throughout the study period. There are

no earlier research findings is available in the literature to corroborate the findings of this experiment. The dietary inclusion of fennel in the diet of turkey poult resulted in consistent numerical increase throughout the experiment period. However, the increase was high in magnitude in males at eighth week making it statistically significant ($P<0.05$). Addition of fennel in the diet of turkey poult revealed a decreasing trend in serum triglyceride level compared to control throughout the experimental period. No meaningful comparison could be possible as there is no traceable literature available on this parameter. Even though the dietary inclusion of cumin seeds had depressive effect on serum cholesterol during entire study period, the difference was statistically significant ($P<0.05$) only for overall mean at eighth week. The decrease in the level of cholesterol is expected to be due to the active compound that found in cumin which acts as inhibitors to the active enzyme hepatic 3-hydroxyl-3 methylglutaryl coenzyme A (HMG-CoA) that synthesized the cholesterol (Crowell, 1999). This reduction in blood cholesterol could be contributed in some cases to the reduction in some hormones secreted by the cortex of the adrenal glands which in turn causes the reduction in the secretion of fatty acids from the adipose tissues or reduction of fat oxidation which leads to the reduction of the level of fatty acids including blood cholesterol.

A similar trend of decreasing effect on serum triglyceride has been noticed in supplementation of cumin seeds throughout the study period similar to fenugreek and fennel seeds supplementation in turkey poult. Similar finding have already been obtained by Al-Kassi (2010) at 1.5% and Al-Kassie et al. (2011) at 0.25, 0.50, 0.75, and 1% observed a significant ($P<0.05$) decrease of serum cholesterol and triglyceride in broiler chicks fed with cumin powder in support of the current findings.

5. Conclusion

Haemoglobin content was constantly lower in thyme supplemented group while it was recorded higher in case of cumin supplemented group. The serum total proteins was significantly higher in thyme, fennel and fenugreek supplemented group compared to cumin supplemented group. The highly significant difference was observed in serum cholesterol level at eighth week of age in turkey poult with highest level in fennel supplemented group and lowest was in cumin supplemented group. The difference in serum triglyceride level was highly significant at eighth week of age with highest triglyceride level in control group while lower level was recorded in fenugreek, thyme, fennel and cumin supplemented groups. Among all treatments supplementation of cumin seeds in feed of turkey poult showed better performance in respect of haemoglobin content, serum calcium and cholesterol level. Further, it is necessary to carry out studies to validate the findings of these at different concentrations of seeds and at different duration.

References

- [1] A.G. Gornall, C.J. Bardawill and M.M. David, "Determination of serum proteins by means of the

- biuret reaction", Journal of Biological Chemistry, 177, pp. 51-66, 1949.
- [2] A.M. Abdullah, J.A. Rabia, "The effect of using fennel seeds (*Foeniculum vulgare*) on productive performance of broiler chickens", International Journal of Poultry Science, 8(7), pp. 642-644, 2009.
- [3] B.T. Dumas, W.A. Watson, H.G. Biggs, "Albumin standards and serum albumin with bromocresol green", Clinical Chemistry Acta, 258, pp. 21-30, 1970.
- [4] C.C. Allain, L.S. Poon, C.S.G. Chan, W. Richmond and P.C. Fu, "Enzymatic determination of total serum cholesterol", Clinical Chemistry, 20, pp. 470-475, 1974.
- [5] E. Demir, K. Kilinc, Y. Yildirim, F. Dincer and H. Eseceli, "Comparative effects of mint, sage, thyme and flavomycin in wheat-based broiler diets", Archiva Zootechnica, 11(3), pp. 54-63, 2008.
- [6] G. Buccolo, H. David, "Quantitative determination of triglycerides by the use of enzymes", Clinical Chemistry, 19, 476-482, 1973.
- [7] G. Kessler, M. Wolfman, "An automated procedure for the simultaneous determination of calcium and phosphorus", Clinical Chemistry, 10, pp. 686-703, 1964.
- [8] G.A.M. Al-Kassi, "Effect of feeding cumin (*Cuminum cyminum*) on the performance and some blood traits of broiler chicks", Pakistan Journal of Nutrition, 9(1), pp. 72-75, 2010.
- [9] G.A.M. Al-Kassie, A.M. Mohseen, A.A. Raghad, "Modification of productive performance and physiological aspects of broilers on the addition of a mixture of cumin and turmeric to the diet", Research Opinions in Animal Veterinary Science, 1(1), pp. 31-34, 2011.
- [10] G.W. Snedecor, W.G. Cochran, "Statistical Methods". 8th ed, Oxford and IBH Publishing Company, New Delhi, India, 1994.
- [11] H. Sahli, Lehrbuch d. klin. Untersuchungen Methode. 5th ed, 846, 1909.
- [12] H.A.A. Elagib, E.M. Nabiela, S.A. Abbass, T.A.N. Ginawi, "Effect of natural spices on plasma proteins in broiler chicks", Journal Nutrition and Food Science, 2(7), pp. 152, 2012.
- [13] H.C. Neu, "The crisis in antibiotic resistance", Science, 257, pp. 1064-1073, 1992.
- [14] H.M. Navid, M.M. Nezhady, "The effect of using Thyme, Garlic and Nettle on performance, carcass quality and blood parameters", Annals of Biological Research, 2(4), pp. 315-320, 2011.
- [15] J.A.A. Isa, "Performance and Lipid profile of broilers fed two medicinal plant", http://www.scholar.najah.edu/publication/thesis_2011
- [16] K.W. Lee, H. Everts, H.J. Kappert, M. Frehner, R. Losa and A.C. Beynen, "Dietary carvacrol lowers body weight gain but improves feed conversion in broiler chickens", Journal of Applied Poultry Research, 12, pp. 394-399, 2003.
- [17] M. Shabaan, 2012. Effect of using Thyme (*Thymus vulgaris*) and Cumin (*Cuminum cyminum*) seeds for improving the utilization of low energy broiler diet. Egyptian Journal of Poultry Science 32(3), 579-591.
- [18] M. Toghyani, M. Mohammadrezaei, A. Gheisari, S. Ali, Tabeidian and H. Ghalamkarix, "Effect of cocoa and thyme powder alone or in combination on humoral immunity and serum biochemical metabolites of broiler chicks", 2nd Intl. Conf. Agric. Animal Science, 22, pp. 114-118, 2011.
- [19] N.A. Botsoglou and D.J. Fletouris, "Drug Residues in Foods. Pharmacology, Food Safety and Analysis" New York, Marcel Dekker, Inc. pp. 541-548, 2001.
- [20] P. William, R. Losa, "The use of essential oils and their compounds in poultry nutrition", World Poultry-Elsevier, 17(4), pp. 14-15, 2001.
- [21] P.L. Crowell, "Prevention and therapy of cancer by dietary monoterpenes", Journal of Nutrition, 129, pp. 775-778, 1999.
- [22] R.J. Abbas, "Effects of using Fenugreek, Parsley and Sweet Basil seeds as feed additives on the performance of broiler chickens", International Journal of Poultry Science, 9(3), pp. 278-282, 2010.
- [23] S. Rahimi, Z.T. Zadeh, M.A.K. Torshizi, R. Omidbaigi and H. Rokni, "Effect of the three herbal extracts on growth performance, immune system, blood factors and intestinal selected bacterial population in broiler chickens", Journal of Agricultural Science Technology, 13, pp. 527-539, 2011.
- [24] S.D. Sharifi, S.H. Khorsandi, A.A. Khadem, A. Salehi and H. Moslehi, "The effect of four medicinal plants on the performance, blood biochemical traits and ileal microflora of broiler chicks", Veterinarski Arhiv, 83 (1), pp. 69-80, 2013.
- [25] Zargari, "Medical plants", Second edition, Tehran University Press, pp. 25-36, 2001