

# On Some New Results of a Subclass of Univalent Functions Defined by Ruscheweyh Derivative

Waggas Galib Atshan<sup>1</sup>, Ali Hussein Battor<sup>2</sup>, Amal Mohammed Dereush<sup>3</sup>

<sup>1</sup>Department of Mathematics, College of Computer Science and Mathematics, University of Al-Qadisiya, Diwaniya-Iraq

<sup>2,3</sup>Department of Mathematics, College of Education for Girls, University of Kufa, Najaf – Iraq

**Abstract:** In this paper, we introduce a new class of univalent functions defined by Ruscheweyh derivative in the open unit disk  $U = \{z \in \mathbb{C}: |z| < 1\}$ , we obtain basic properties, like, coefficient inequality, distortion and covering theorem, radii of starlikeness, convexity and close-to-convexity, extreme points, Hadamard product, closure theorems and convolution operator for functions belonging to the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Keywords:** Univalent function, Ruscheweyh derivative, Distortion theorem, Radius of starlikeness, Extreme points, Hadamard product.

**2014 Mathematics Subject Classification:** 30C45.

## 1. Introduction

Let  $\Sigma$  denote the class of functions of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1)$$

which are analytic and univalent in the open unit disk  $U$ .

If a function  $f$  is given by (1) and  $g$  is defined by

$$g(z) = z + \sum_{n=2}^{\infty} b_n z^n, \quad (2)$$

is in the class  $\Sigma$ , then the convolution (or Hadamard product) of  $f$  and  $g$  is defined by

$$(f * g)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n, \quad z \in U. \quad (3)$$

Let  $\Sigma^+$  denote the subclass of  $\Sigma$  consisting of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (a_n \geq 0, n \in \mathbb{N}). \quad (4)$$

We aim to study the subclass  $\Sigma^+(\sigma, c, \beta, \lambda)$  consisting of function  $f \in \Sigma^+$  and satisfying the condition:

$$\left| \frac{\sigma [z (D^\lambda f(z))'' - ((D^\lambda f(z))' - 1)]}{cz (D^\lambda f(z))'' + ((1 - \sigma)(D^\lambda f(z))' + 1)} \right| < \beta, \quad z \in U, \quad (5)$$

where  $0 \leq \sigma < 1, 0 \leq c < 1, 0 < \beta < 1$  and  $D^\lambda f(z)$  is the Ruscheweyh derivative [6], [7] of  $f$  of order  $\lambda$  defined as follow:

$$D^\lambda f(z) = z + \sum_{n=2}^{\infty} a_n A_n(\lambda) z^n,$$

where

$$A_n(\lambda) = \frac{(\lambda + 1)(\lambda + 2) \dots (\lambda + n - 1)}{(n - 1)!}, \quad \lambda > -1, \quad z \in U \quad (6)$$

Another classes studied by several authors, like, [2] and [4] consisting of functions of the form (4).

## 2. Coefficient Inequality

In the following theorem, we obtain necessary and sufficient condition to be the function in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 1:** Let the function  $f$  be defined by (4). Then  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$  if and only if

$$\sum_{n=2}^{\infty} [n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda) a_n \leq \beta(2 - \sigma), \quad (7)$$

where  $0 < \beta < 1, 0 < \sigma < 1, 0 \leq c < 1$ , and  $\lambda > -1$ . The result (7) is sharp for the function

$$f(z) = z + \frac{\beta(2 - \sigma)}{[n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda)} z^n, \quad (n \geq 2). \quad (8)$$

**Proof:** Suppose that the inequality (7) holds true and  $|z| = 1$ . Then we have

$$\left| \sigma [z (D^\lambda f(z))'' - ((D^\lambda f(z))' - 1)] - \beta [cz (D^\lambda f(z))'' + ((1 - \sigma)(D^\lambda f(z))' + 1)] \right|$$

$$= \left| \sum_{n=2}^{\infty} (\sigma n^2) A_n(\lambda) a_n z^{n-2} - \beta \left[ \sum_{n=2}^{\infty} (cn^2 - cn + n - \sigma n) A_n(\lambda) a_n z^{n-2} + (2 - \sigma) \right] \right|$$

$$\leq \sum_{n=2}^{\infty} [n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda) a_n - \beta(2 - \sigma) \leq 0,$$

by hypothesis, hence, by maximum modulus principle  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ .

Conversely, assume that  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ , so that

$$\left| \frac{\sigma [z (D^\lambda f(z))'' - ((D^\lambda f(z))' - 1)]}{cz (D^\lambda f(z))'' + ((1 - \sigma)(D^\lambda f(z))' + 1)} \right| < \beta, \quad z \in U,$$

hence

$$\begin{aligned} & \left| \sigma \left[ z \left( D^\lambda f(z) \right)'' - \left( \left( D^\lambda f(z) \right)' - 1 \right) \right] \right. \\ & \quad < \beta \left| cz \left( D^\lambda f(z) \right)'' \right. \\ & \quad \left. + \left( (1 - \sigma) \left( D^\lambda f(z) \right)' + 1 \right) \right|. \end{aligned}$$

Therefore, we get

$$\left| \sum_{n=2}^{\infty} (\sigma n^2) A_n(\lambda) a_n z^{n-2} < \beta \left| \sum_{n=2}^{\infty} (cn^2 - cn + n - \sigma n) A_n(\lambda) a_n z^{n-2} + (2 - \sigma) \right|,$$

thus

$$\sum_{n=2}^{\infty} [n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda) a_n \leq \beta(2 - \sigma).$$

**Corollary 1:** Let the function  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then

$$a_n \leq \frac{\beta(2 - \sigma)}{[n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda)} z^n, n \geq 2.$$

### 3. Distortion and Covering Theorems

We introduce the growth and distortion theorems for the function  $f$  in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 2:** Let the function  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then

$$\begin{aligned} & \left| z - \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z|^2 \right| \leq |f(z)| \\ & \leq |z| + \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z|^2, |z| < 1. \end{aligned}$$

The result is sharp and attained

$$f(z) = z + \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} z^2.$$

**Proof:**

$$\begin{aligned} |f(z)| &= |z + \sum_{n=2}^{\infty} a_n z^n| \\ &\leq |z| + \sum_{n=2}^{\infty} a_n |z|^n \leq |z| + |z|^2 \sum_{n=2}^{\infty} a_n. \end{aligned}$$

By Theorem (1), we get

$$\sum_{n=2}^{\infty} a_n \leq \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)}. \tag{9}$$

Thus

$$|f(z)| \leq |z| + \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z|^2.$$

Also

$$\begin{aligned} |f(z)| &\geq |z| - \sum_{n=2}^{\infty} a_n |z|^n \\ &\geq |z| \\ &\quad - |z|^2 \sum_{n=2}^{\infty} a_n \\ &\geq |z| \\ &\quad - \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z|^2, \end{aligned}$$

and this completed the proof.

**Theorem 3:** Let  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then

$$\begin{aligned} & 1 - \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z| \leq |f'(z)| \\ & \leq 1 \\ & \quad + \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} |z| \end{aligned}$$

with equality for

$$f(z) = z + \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)} z^2.$$

**Proof:** Notice that

$$\begin{aligned} & [2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1) \sum_{n=2}^{\infty} n a_n \\ & \leq \sum_{n=2}^{\infty} [n(\sigma(n - 2 + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda) a_n \\ & \leq \beta(2 - \sigma), \tag{10} \end{aligned}$$

from Theorem 1. Thus

$$\begin{aligned} |f'(z)| &= \left| 1 + \sum_{n=2}^{\infty} n a_n z^{n-1} \right| \\ &\geq 1 - \sum_{n=2}^{\infty} n a_n |z|^{n-1} \geq 1 - |z| \sum_{n=2}^{\infty} n a_n \\ &\geq 1 - |z| \frac{\beta(2 - \sigma)}{[2(\sigma(2 + \beta) - \beta(c + 1))] (\lambda + 1)}. \tag{11} \end{aligned}$$

Combining (10) and (11), we get the result.

### 4. Radii of starlikeness, convexity and close-to-convexity:

In the following theorems, we obtain the radii of starlikeness, convexity and close-to-convexity for the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 4:** Let  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then  $f$  is starlike in the disk  $|z| < R_1$ , of order  $\alpha$ ,  $0 \leq \alpha < 1$ , where

$$R_1 = \inf_n \left[ \frac{(1 - \alpha) [n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda)}{(n - \alpha) \beta(2 - \sigma)} \right]^{\frac{1}{n-1}}, n$$

**Proof:**  $f$  is starlike of order  $\alpha$ ,  $0 \leq \alpha < 1$ , if

$$\operatorname{Re} \left\{ \frac{z f'(z)}{f(z)} \right\} > \alpha.$$

We must show that

$$\left| \frac{z f'(z)}{f(z)} - 1 \right| < 1 - \alpha,$$

for  $|z| < R_1$ .

Indeed we have

$$\begin{aligned} \left| \frac{z f'(z)}{f(z)} - 1 \right| &\leq \frac{\sum_{n=2}^{\infty} (n - 1) a_n |z|^{n-1}}{1 - \sum_{n=2}^{\infty} a_n |z|^{n-1}} \\ &\leq 1 - \alpha, \quad (0 \leq \alpha < 1) \tag{13} \end{aligned}$$

hence by Theorem 1, (13) will be true if

$$\frac{n - \alpha}{1 - \alpha} |z|^{n-1} \leq \frac{[n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda)}{\beta(2 - \sigma)}$$

or if

$$\begin{aligned} & |z| \\ & \leq \left[ \frac{(1 - \alpha) [n(\sigma(n + \beta) - \beta(c(n - 1) + 1))] A_n(\lambda)}{(n - \alpha) \beta(2 - \sigma)} \right]^{\frac{1}{n-1}}, n \\ & \geq 2 \tag{14} \end{aligned}$$

the theorem follows easily from (14).

**Theorem 5:** Let  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then  $f$  is convex in  $|z| < R_2$  of order  $\alpha, 0 \leq \alpha < 1$ , where  $R_2$

$$= \inf_n \left[ \frac{(1-\alpha)[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{n(n-\alpha)\beta(2-\sigma)} \right]^{\frac{1}{n-2}}, n \geq 2 \quad (15)$$

**Proof:**  $f$  is convex of order  $\alpha, 0 \leq \alpha < 1$ , if

$$Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha.$$

Thus it is enough to show that

$$\left| \frac{zf''(z)}{f'(z)} \right| \leq 1 - \alpha,$$

for  $|z| < R_2$ .

Indeed we have

$$\left| \frac{zf''(z)}{f'(z)} \right| \leq \frac{\sum_{n=2}^{\infty} n(n-1)a_n|z|^{n-1}}{1 - \sum_{n=2}^{\infty} na_n|z|^{n-1}} \leq 1 - \alpha, (0 \leq \alpha < 1) \quad (16)$$

Hence by Theorem 1, (16) will be true if

$$\frac{n(n-\alpha)|z|^{n-1}}{(1-\alpha)} \leq \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{\beta(2-\sigma)}$$

or if

$$\left| z \right| \leq \left[ \frac{(1-\alpha)[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{n(n-\alpha)\beta(2-\sigma)} \right]^{\frac{1}{n-1}}, n \geq 2. \quad (17)$$

**Theorem 6:** Let  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then  $f$  is close-to-convex function in  $|z| < R_3$  of order  $\alpha, 0 \leq \alpha < 1$ , where  $R_3$

$$= \inf_n \left[ \frac{(1-\alpha)[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{n\beta(2-\sigma)} \right]^{\frac{1}{n-1}}. \quad (18)$$

**Proof:**  $f$  is close-to-convex function of order  $\alpha, 0 \leq \alpha < 1$ , if

$$Re\{f'(z)\} > \alpha.$$

Thus it is enough to show that

$$|f'(z) - 1| = \left| \sum_{n=2}^{\infty} na_n z^{n-1} \right| \leq \sum_{n=2}^{\infty} na_n |z|^{n-1}.$$

then

$$\sum_{n=2}^{\infty} \left( \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{\beta(2-\sigma)} \right) \mu_n \frac{\beta(2-\sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)} = \sum_{n=2}^{\infty} \mu_n = 1 - \mu_1 \leq 1.$$

Using Theorem 1, we easily get  $(z) \in \Sigma^+(\sigma, c, \beta, \lambda)$ .

Conversely, let  $f(z) \in \Sigma^+(\sigma, c, \beta, \lambda)$  is of the form (4). Then

$$a_n \leq \frac{\beta(2-\sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}, (n \geq 2).$$

Setting

$$\mu_n = \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{\beta(2-\sigma)} a_n, \text{ for } n \geq 2$$

and

$$\mu_1 = 1 - \sum_{n=2}^{\infty} \mu_n.$$

Then

Thus

$$|f'(z) - 1| \leq 1 - \alpha \text{ if } \sum_{n=2}^{\infty} \frac{na_n|z|^{n-1}}{1-\alpha} \leq 1, \quad (19)$$

hence by Theorem 1, (19) will be true if

$$\frac{n|z|^{n-1}}{1-\alpha} \leq \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{\beta(2-\sigma)}$$

or if

$$\left| z \right| \leq \left[ \frac{(1-\alpha)[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)}{n\beta(2-\sigma)} \right]^{\frac{1}{n-1}}, n \geq 2 \quad (20)$$

the result follows easily from (20).

### 5. Extreme Points:

In the following theorem, we obtain extreme points for the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 7:** Let  $f_1(z) = z$  and

$$f_n(z) = z + \frac{\beta(2-\sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)} z^n, \text{ for } n = 2, 3, \dots$$

Then  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$  if and only if it can be expressed in the form

$$f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z),$$

where

$$\left( \mu_n \geq 0 \text{ and } \sum_{n=1}^{\infty} \mu_n = 1 \text{ or } 1 = \mu_1 + \sum_{n=2}^{\infty} \mu_n \right).$$

**Proof:** Let

$$f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z) = z + \sum_{n=2}^{\infty} \frac{\beta(2-\sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)} \mu_n z^n,$$

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n = z + \sum_{n=2}^{\infty} \frac{\beta(2-\sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))]A_n(\lambda)} \mu_n z^n = \mu_1 z + \sum_{n=2}^{\infty} \mu_n f_n(z).$$

Thus

$$f(z) = \sum_{n=1}^{\infty} \mu_n f_n(z) = \mu_1 f_1(z) + \sum_{n=2}^{\infty} \mu_n f_n(z).$$

### 6. Hadamard Product

In the following theorem, we obtain the convolution result for function belong to the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 8:** Let  $f$  and  $g \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then  $f * g \in \Sigma^+(\sigma, c, \delta, \lambda)$  for

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, g(z) = z + \sum_{n=2}^{\infty} b_n z^n,$$

where

$$\delta \leq \frac{\beta^2(2 - \sigma)[n\sigma(n - 2)]}{A_n(\lambda)[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]^2 - \beta^2(2 - \sigma)[n(\sigma - (c(n - 1) + 1))]}.$$

**Proof:** Since  $f, g \in \Sigma^+(\sigma, c, \beta, \lambda)$ , then we have

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} a_n \leq 1 \quad (21)$$

and

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} b_n \leq 1. \quad (22)$$

We must find the smallest number  $\delta$  such that

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]A_n(\lambda)}{\delta(2 - \sigma)} a_n b_n \leq 1. \quad (23)$$

By Cauchy-Schwarz inequality, we have

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} \sqrt{a_n b_n} \leq 1. \quad (24)$$

Thus, it is enough to show that and

$$\delta \leq \frac{\beta^2(2 - \sigma)[n\sigma(n - 2)]}{A_n(\lambda)[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]^2 - \beta^2(2 - \sigma)[n(\sigma - (c(n - 1) + 1))]}.$$

This complete the proof.

**Theorem 9:** Let  $g \in \Sigma^+(\sigma, c, \beta, \lambda)$ . Then

$$h(z) = z + \sum_{n=2}^{\infty} (a_n^2 + b_n^2)z^n$$

belong to the class  $\Sigma^+(\sigma, c, \delta, \lambda)$ , where

$$\delta \geq \frac{2\beta^2(2 - \sigma)n\sigma}{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]^2 A_n(\lambda)}.$$

**Proof:** Since  $f, g \in \Sigma^+(\sigma, c, \beta, \lambda)$  so by Theorem 1, yields

$$\sum_{n=2}^{\infty} \left[ \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} a_n \right]^2 \leq 1$$

and

$$\sum_{n=2}^{\infty} \left[ \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} b_n \right]^2 \leq 1,$$

we obtain from the last two inequalities

$$\sum_{n=2}^{\infty} \frac{1}{2} \left[ \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} \right]^2 (a_n^2 + b_n^2) \leq 1, \quad (27)$$

but  $h(z) \in \Sigma^+(\sigma, c, \delta, \lambda)$  if and only if

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]A_n(\lambda)}{\delta(2 - \sigma)} (a_n^2 + b_n^2) \leq 1, \quad (28)$$

where  $0 < \delta < 1$ , however (27) implies (28) if

$$\frac{[n(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]A_n(\lambda)}{\delta(2 - \sigma)} \leq \left[ \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} b_n \right]^2.$$

$$\frac{[n(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]A_n(\lambda)}{\delta(2 - \sigma)} a_n b_n \leq \frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} \sqrt{a_n b_n},$$

that is

$$\sqrt{a_n b_n} \leq \frac{[(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]\delta}{[(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]\beta}, \quad (25)$$

from (24), we get

$$\sqrt{a_n b_n} \leq \frac{\beta(2 - \sigma)}{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}. \quad (26)$$

Therefore, in view of (25) and (26) it is enough to show that

$$\frac{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda)}{\beta(2 - \sigma)} \leq \frac{[(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]\delta}{[(\sigma(n + \delta)) - \delta(c(n - 1) + 1)]\beta}$$

Simplifying, we get

$$\delta \geq \frac{2\beta^2(2 - \sigma)n\sigma}{[n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]^2 A_n(\lambda)}.$$

### 7. Closure theorems

We shall prove the following closure theorems for the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ , let the function  $f_i(z)$  ( $i = 1, 2, \dots, m$ ) defined by

$$f_i(z) = z + \sum_{n=2}^{\infty} a_{n,i} z^n, (a_{n,i} \geq 0, n \in N, n \geq 2) \quad (29).$$

**Theorem 10:** Let the functions  $f_i(z)$  defined by (29) be in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$  for every  $i = 1, 2, \dots, m$ . Then the function  $h(z)$  defined by

$$h(z) = z + \sum_{n=2}^{\infty} c_n z^n, (c_n \geq 0, n \in N, n \geq 2)$$

also belongs to the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ , where

$$c_n = \frac{1}{m} \sum_{i=1}^m a_{n,i}.$$

**Proof:** Since  $f_i(z) \in \Sigma^+(\sigma, c, \beta, \lambda)$ , therefore from Theorem 1, we obtain

$$\sum_{n=2}^{\infty} [n(\sigma(n + \beta)) - \beta(c(n - 1) + 1)]A_n(\lambda) a_{n,i} \leq \beta(2 - \sigma), \quad (29)$$

then

$$\sum_{n=2}^{\infty} [n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda) c_n$$

$$= \sum_{n=2}^{\infty} [n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda) \left[ \frac{1}{m} \sum_{i=1}^m a_{n,i} \right]$$

$$\leq \beta(2 - \sigma).$$

Hence  $h(z) \in \Sigma^+(\sigma, c, \beta, \lambda)$ .

**Theorem 11:** Let the functions  $f_i(z)$  defined by (29) be in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ , for every  $i = 1, 2, \dots, m$ . Then the function  $h(z)$  defined by

$$h(z) = \sum_{i=1}^m d_i f_i(z) \text{ and } \sum_{i=1}^m d_i = 1, d_i \geq 0$$

in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ .

**Proof:** By definition of  $h(z)$ , we have

$$h(z) = \sum_{i=1}^m d_i z + \sum_{n=2}^{\infty} \left[ \sum_{i=1}^m d_i a_{n,i} \right] z^n,$$

since  $f_i(z)$  are in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ , for every  $i = 1, 2, \dots, m$ , we obtain

$$\sum_{n=2}^{\infty} [n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda) a_{n,i}$$

$$\leq \beta(2 - \sigma)$$

for every  $i = 1, 2, \dots, m$ , hence we can see that

$$\sum_{n=2}^{\infty} [n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda) \left[ \sum_{i=1}^m d_i a_{n,i} \right]$$

$$= \sum_{i=1}^m d_i \left[ \sum_{n=2}^{\infty} [n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda) a_{n,i} \right]$$

$$\leq \beta(2 - \sigma) \sum_{i=1}^m d_i = \beta(2 - \sigma).$$

Thus  $h(z) \in \Sigma^+(\sigma, c, \beta, \lambda)$ .

### 8. Convolution Operator

**Definition 1 [2,5]:** The Gaussian hypergeometric function denoted by

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \cdot \frac{z^n}{n!}, |z| < 1,$$

where  $c > b > 0, c > a + b$  and

$$(x)_n = \begin{cases} x(x+1)(x+2) \dots (x+n-1) & \text{for } n = 1, 2, 3, \dots \\ 1 & n = 0 \end{cases}$$

**Definition 2[3]:** For every  $f \in \Sigma^+$ , we defined the convolution operator  $W_{a,b,c}(f)(z)$  as below:

$$W_{a,b,c}(f)(z) = {}_2F_1(a, b; c; z) * f(z)$$

$$= z + \sum_{n=2}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} a_n z^n,$$

where  ${}_2F_1(a, b; c; z)$  is Gaussain hypergeometric function (see[2] and [5]) introduced in Definition 1.

**Theorem 12:** Let  $f$  is given by (4) be in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$ . Then the convolution operator  $W_{a,b,c}(f)$  is in the class  $\Sigma^+(\sigma, c, \beta, \lambda)$  for  $|z| \leq r(\beta, \delta)$ , where

$$r(\beta, \delta)$$

$$= \inf_n \left[ \frac{\delta [n(\sigma(n+\beta) - \beta(c(n-1) + 1))]}{\beta [n(\sigma(n+\delta) - \delta(c(n-1) + 1))]} \frac{(a)_n (b)_n}{(c)_n n!} \right]^{\frac{1}{n-1}}.$$

The result is sharp for the function

$$f_n(z) = z + \frac{\beta(2 - \sigma)}{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda)} z^n, n \geq 2.$$

**Proof:** Since  $f \in \Sigma^+(\sigma, c, \beta, \lambda)$ , we have

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda)}{\beta(2 - \sigma)} a_n \leq 1.$$

It is sufficient to show that

$$\sum_{n=2}^{\infty} \frac{[n(\sigma(n+\delta) - \delta(c(n-1) + 1))]}{\delta(2 - \sigma)} \frac{(a)_n (b)_n}{(c)_n n!} a_n \leq 1. (30)$$

Note that (30) is satisfied if

$$\frac{[n(\sigma(n+\delta) - \delta(c(n-1) + 1))]}{\delta(2 - \sigma)} \frac{(a)_n (b)_n}{(c)_n n!} a_n |z|^{n-1} \leq \frac{[n(\sigma(n+\beta) - \beta(c(n-1) + 1))] A_n(\lambda)}{\beta(2 - \sigma)} a_n,$$

solving for  $|z|$  we get the result.

### Reference

- [1] G. E. Andrews, R. Askey and R. Roy, Special functions, Cambridge University Press, Cambridge, U. K., (1999).
- [2] W. G. Atshan, and R. M. Abd, New class of univalent functions with negative coefficients defined by Ruscheweyh derivative, Gen. Math. Notes, 18(2)(2013), pp 77-91.
- [3] W. G. Atshan and S.R. Kulkarni, On a new class of analytic functions with negative coefficients, Analete Universitatii Oradea, Fasc. Mathematica, Tom, XVII(2009), 34-51.
- [4] M. Darus, Some subclasses of analytic functions, J. Math. Comp. Sci. Math. Ser., 16(3)(2003), 121-126.
- [5] Y. C. Kim and F. Ronning, Integral transforms of certain subclass of analytic functions, J. Math. Appl., 258(2001), 466-489.
- [6] S. Ruscheweyh, New criteria for univalent functions, Proc. Amer. Math. Soc., 49(1975), 109-115.
- [7] H. M. Srivastava and S. Owa(Eds), Current Topics in Analytic Function Theory, World Scientific, Singapore, (1992).

### Author Profile



**Waggas Galib Atshan**, Assist. Prof. Dr. in Mathematics (Complex Analysis), teacher at University of Al-adisiya, College of Computer Science & Mathematics, Depart. of Mathematics, he has 90 papers published in various journals in

Mathematics till now, he taught seventeen subjects in mathematics till now (undergraduate, graduate), he is supervisor on 20 students (Ph.D., M.Sc.) till now, he attended 23 international and national conferences.

**Ali Hussein Battor**, Prof. Dr. in Mathematics (Functional Analysis), at University of Kufa, College of Education for Girls, Department of Mathematics, he has many papers published in various journals in mathematics, he teached number of subjects in mathematics (undergraduate, graduate), he is supervisor on more than 25 students till now, he attended More than 30 international and national conferences.