

A Comparative Analysis of the Physico-chemical Properties and Heavy Metal Pollution in Three Major Rivers across India

Sunita Singh¹, Darshan Malik², JayitaThakur³, Amarjeet Kaur⁴, Raj Kishore Singh⁵, Shashi Nijhawan⁶

^{1, 2, 3, 6} Department of Biochemistry, Shivaji College, University of Delhi, New Delhi, India

^{4, 5} University School of Environment Management, GGSIP University, New Delhi, India

Abstract: *There is severe deterioration in the quality of water due to the discharge of municipal and industrial effluents into the rivers. The municipal pollutants majorly comprise of untreated domestic and sewage wastes, while the industrial pollutants constitute the discharge of heavy metals into the river which is responsible for the increase in metal load in water. In this study we have carried out a comparative analysis of the various physico-chemical parameters and heavy metal load in three major rivers of India. The water quality of the Ganges river at Kanpur and Varanasi, the Yamuna river at Delhi and the Sabarmati river at Ahmedabad were analyzed for the determination of metal load (Lead, Copper, Zinc, Chromium, Cadmium and Nickel) and biological load (physico-chemical parameters like pH, Total dissolved solids, Total suspended solids, Biological oxygen demand, Chemical oxygen demand and Total coliform). Our study indicates that, of the four stations, the levels of Chromium exceeded the acceptable levels in the Yamuna river (0.08mg/L). The levels of Cadmium and Lead were found to be within permissible limits, however, as they tend to persist in the system for a long time, they undergo bioaccumulation and biomagnification. This leads to severe nephrosis and liver damage over a long period of time.*

Keywords: Heavy metals, Physico-chemical parameters, Anthropogenic, Domestic and Industrial effluent, Human Health, Toxicity and Genotoxicity

1. Introduction

The rivers play an extremely important role in the lives of the people of India. The river systems are used in irrigation of crops, acquiring potable water, generation of electricity, as well as for the provision of livelihoods for a large number of people across the country. The rivers also play an important role in the Hindu mythology. Important cities like Delhi, Varanasi, Kanpur and Ahmedabad are situated on the banks of the Yamuna, Ganges, and Sabarmati respectively. These rivers during their meandering receive a wide array of chemical constituents including toxicants from a variety of natural and anthropogenic sources (Kar *et al.*, 2008). With the rapid growth in population and increased industrial activities, the preservation and maintenance of our natural water resources has become an increasingly difficult task. The quality of water resources is on a severe decline as there is a continuous influx of undesirable chemicals, while the demand of potable water is on a rise.

Industrial pollution has far reaching implications. A large number of industries like dyeing, tanning, printing and battery manufacturing units operate alongside the rivers and discharge their effluents into it, contaminating the groundwater and river water, thus affecting the quality of water. Heavy metals like Lead (Pb), Copper (Cu), Zinc (Zn), Chromium (Cr), Cadmium (Cd) and Nickel (Ni) are the metallic elements that have a relatively high density and are toxic when present above the permissible limits (Lenntech Water Treatment and Air Purification, 2004). These heavy metals mainly enter the human body through food and water and are known to have serious health implications including carcinogenesis (Schwartz, 1994). Fertilizers, insecticides and pesticides also add to the water pollution by leaching. Some of the major causes of heavy metal accumulation in the rivers

are flawed mining processes, discharge of industrial effluents containing metallic solutions, dumping of solid wastes which contain metal salts and some agricultural practices such as the use of metal-based biocides (Anand and Pandey, 2014).

Lead is ubiquitously present heavy metal with consequences which range from cognitive impairment in children to peripheral neuropathy in adults. Pb poisoning also causes neuromuscular and central nervous system disorders (Kaur, 2012). It is used in paints, storage batteries, and the oxide is used in producing fine crystal glass. It enters the human body either through inhalation of dust from lead paints or through various foods, notably fish and plants. The heavy metal copper, which is an essential nutrient, when present at low concentrations causes headache, nausea, vomiting and diarrhea. At higher levels of deposition, it leads to liver and kidney malfunctioning (USEPA, 1999). As it is slow in degradation, it leads to bioaccumulation and magnification over long periods of time. Electroplating industries situated on the banks of the rivers are the primary source of copper effluents in the water bodies (Boxall *et al.*, 2000).

Zinc is an essential element for all living beings. Zn is involved in various physiological and metabolic activities of many organisms (Pillai, 1983). Proteins and enzymes containing Zn are involved in metabolism, including the replication and translation of genetic material. The toxicity of Zinc causes vomiting, diarrhea, icterus, liver and kidney damage. The sources of Zn in the rivers are effluents from electroplating industries, sewage discharge and the immersion of painted idols (Boxall *et al.*, 2000; Dean *et al.*, 1972). Nickel is a well known neurotoxic, genotoxic and carcinogenic agent which may cause health problems like nickel dermatitis, giddiness, diarrhea, degeneration of the liver and various types of cancer (Das *et al.*, 2008). Industries like Stainless steel manufacturing units, electroplating

factory discharge Ni in their effluents. Other sources of contamination include electrical equipment, household appliances, batteries and the dumping of municipal sewage (Dean *et al.*, 1972). Occupational and environmental exposure to Cr and Cd has been previously linked to a wide array of inflammatory and degenerative conditions and cancer. Cadmium is a by-product of Zinc production and is one of the most toxic heavy metals. The potential for Cadmium to harm human health depends upon the form present and the amount taken (Bernard, 2008). The intake of Cd over a long period of time leads to its bio-accumulation in the kidney and liver for years and results in severe kidney and liver damage. It is employed in several industrial processes such as protective coatings (electroplating) for metals like iron, preparation of Cd-Ni batteries, control rods and shields within nuclear reactors and television phosphors. Chromium is a notorious environmental pollutant used in several anthropogenic activities. It is a transition metal which is present in the hexavalent Cr (VI) and trivalent Cr (III) forms (Dey and Paul, 2013). Hexavalent chromium toxicity is more severe due to its high permeability to bio-membranes. Ingestion of large amounts of chromium (VI) also has severe detrimental health effects. It can lead to severe respiratory, cardiovascular, gastrointestinal, hepatic and renal damage and potentially death (Bagchi *et al.*, 2002). The presence of Cr in the Yamuna river is attributed to electroplating industries situated near the banks of the river.

The values of pH, Total dissolved solids (TDS), Total suspended solids (TSS), Biological oxygen demand (BOD), Chemical oxygen demand (COD) and total coliform are indicators of the levels of microorganisms present and quality of the water body. BOD is a measure of the amount of oxygen that bacteria will consume while decomposing organic matter under aerobic conditions while COD does not differentiate between the biologically available and inert organic matter (Lenore *et al.*, 2005; Malik *et al.*, 2014).

In the present study, the physico-chemical parameters like pH, TDS, TSS, BOD, COD and total coliform have been examined from the samples collected from the four stations viz., i) Ganges at Kanpur and Varanasi, ii) Yamuna at Delhi and iii) Sabarmati at Ahmedabad. In view of the hazardous nature of metal ions, an attempt has also been made to study the pollution potential of Pb, Cu, Zn, Cr, Cd and Ni at the above mentioned stations.

2. Materials and Methods

2.1 Sampling of river water

The water samples for analysis were collected from midstream at 15 cm depth from the Ganges river at Kanpur (Nanarao Ghat) and Varanasi (Dashashwamedh Ghat), Yamuna river at Delhi (Nizamuddin Bridge) and Sabarmati

river at Ahmedabad (Sabarmati Ashram). The sample bottles were then soaked in 10 percent HNO₃ for 24 h and rinsed thoroughly with double distilled water prior to use. The water samples were then preserved with concentrated HNO₃ to bring down the pH to <2.0. They were then stored at 4°C and brought to the laboratory for metal analysis (Ali and Jain, 2001).

2.2 Physico-chemical analysis of river water

The water was processed immediately post collection, in order to ensure that the water profile is not affected by storage. The various physico-chemical parameters like pH, TDS, TSS, BOD and COD were estimated using pH meters and titrimetric methods (APHA, 2005) respectively. The total coliforms present in the water were also estimated by dilution method to estimate the most probable number (MPN) of coliform bacteria in 100 ml of culture (APHA, 20th edition, 1998).

2.3 Heavy metal analysis of river water

The analysis of the toxic metals present in the water samples were carried out using Perkin Elmer Atomic Absorption Spectrometer (Model 3110). The operational conditions were adjusted to obtain optimal determination. The quantification of metals was based upon calibration curves of standard solutions of metals (Jain and Ali, 2001).

3. Results

3.1. Analysis of physico-chemical parameters

Water samples were collected from four stations of the three major rivers, namely Ganges (Kanpur and Varanasi), Yamuna (Delhi) and Sabarmati (Ahmedabad), and were analyzed for various physico-chemical parameters (Table 1). The pH of the water samples were found to vary between the acceptable range of 7.38 (Sabarmati, Ahmedabad) and 7.72 (Yamuna, Delhi) (Fig. 1). The total TDS found in the samples ranged between 235 mg/L (Ganges, Kanpur) to 642 mg/L (Yamuna, Delhi) respectively, which though well within the permissible limits, is quite high. The total TSS was found to range between 12 mg/L (Ganges, Varanasi) and 230 mg/L (Yamuna, Delhi). The BOD levels were from 4.6 mg/L (Sabarmati at Ahmedabad) to 29 mg/L (Yamuna, Delhi), while the COD levels were found to vary from 19.7 mg/L (Ganges, Varanasi) and 110 mg/L (Yamuna, Delhi) (Fig. 2). The total coliform count was found to be the least at 62342 MPN/100 ml (Ganges, Varanasi) and the maximum of 11786556 MPN/100 ml (Yamuna, Delhi) (Fig. 3), which was much higher than the accepted levels.

Table 1: The Physicochemical parameters of water viz., pH, TDS, TSS, BOD, COD and Coliform organism were determined

LOCATION	pH	Total Dissolved Solids TDS (mg/L)	Total Suspended solids TSS (mg/L)	Biochemical Oxygen Demand (BOD) 20 °C (mg/L)	Chemical Oxygen Demand (COD) (mg/L)	Coliform Organism (MPN/100 ml)
Ganga (Kanpur- Nanarao Ghat)	7.4	235	22	6.7	19.7	131000
Ganga (Varanasi- Dashashwamedh Ghat)	7.56	320	12	7.4	21.5	62342
Yamuna (Delhi- Nizamuddin Bridge)	7.72	642	230	29	110	11786556
Sabarmati (Ahmedabad-Sabarmati)	7.38	245	16	4.6	31	590000




Figure 1: pH analysis of water samples collected from four different stations across India viz., Ganges at Kanpur and Varanasi, Yamuna River at Delhi and Sabarmati at Ahmedabad

3.2. Analysis of heavy metal levels

Water samples collected from the four stations located on the rivers Ganges, Yamuna and Sabarmati were analyzed for the levels of heavy metals present in them (Table 2). In the present study the heavy metals Pb, Cu, Zn, Cr, Cd and Ni were analyzed and observed for their toxicity.






Figure 2: Comparative analysis of the physico-chemical parameter soft water samples collected from four different stations. Biological Oxygen Demand (BOD) and Chemical Oxygen Demand (COD) were estimated using titrimetric methods (APHA, 2005). The values of BOD and COD are indicators of the levels of microorganisms present in a water body.



A



B

Figure 3 A & B: Analysis of the Coliform in the collected water samples. The maximum numbers of coliforms were detected in the Yamuna river in Delhi at Nizamuddin Bridge. Indicating a high microbial load. The lowest numbers of coliforms were detected in Ganga at Kanpur and Varanasi.

Our results suggest that the concentrations of metal ions vary significantly depending upon the nature of the effluent discharged into the rivers. The level of the toxic heavy metals (mg/L) reported by ICMR, CPCB, WHO and EPA have been graphically represented (Fig. 4). The levels of lead was found to range between 0.01 mg/L (Ganges, Kanpur) and 0.030 mg/L (Yamuna, Delhi), which is within the permissible limits (Table 2). Zinc levels were found to range between 0.15 mg/L (Ganges, Kanpur) and 0.133 mg/L (Yamuna, Delhi) and were well within the acceptable levels. The levels of copper which ranged between 0.02 mg/L (Ganges, Kanpur) and 0.06 mg/L (Yamuna, Delhi) were also within the permissible limits (Fig. 5). Of the four stations, the lowest level of Cadmium was found to be 0.043 mg/L (Ganges, Varanasi) and the highest was 0.061 mg/L (Yamuna, Delhi) (Fig. 5). Thus, the levels of these metals are in permissible limits except for Chromium ranging from 0.02 to 0.08 mg/L, which is above the toxicity level in Yamuna at Nizamuddin.




Figure 4: Prescribed limits of metals viz., Lead, Zinc, Cadmium, Copper, Chromium, Nickel reported by ICMR (Indian Council of Medical Research), CPCB (Central Pollution Control Board), WHO (World Health Organization) and EPA (Environmental Pollution Agency)

Table 2: The determination of toxic heavy metals was done in the collected samples from different places

Station	Lead (Pb) (mg/L)	Copper (Cu) (mg/L)	Zinc(Zn) (mg/L)	Chromium(Cr +6) (mg/L)	Cadmium (Cd) (mg/L)	Nickel (Ni) (mg/L)
Ganga (Kanpur- Nanarao Ghat)	0.01	0.02	0.15	0.02	0.075	0.70
Ganga (Varanasi-Dashashwamedh Ghat)	0.01	0.02	0.2	0.02	0.043	0.55
Yamuna (Delhi-Nizamuddin Bridge)	0.030	0.06	0.133	0.08	0.061	0.32
Sabarmati (Ahmedabad-Sabarmati Ashram)	0.02	0.05	1.2	0.02	0.059	0.46







Figure 5: Heavy metal concentrations in the water samples collected from the different stations were determined using Elmer Atomic Absorption Spectrometer (AAS). Significant levels of Zn, Cr and Ni were observed. The four stations included in study are: A- Ganga (Kanpur- Nanarao Ghat); B-Ganga (Varanasi-Dashashwamedh Ghat); C-Yamuna (Delhi-Nizamuddin Bridge); D- Sabarmati (Ahmedabad-Sabarmati Ashram).



4. Discussion

The exposure to toxic metals pose a serious human health hazard based on ubiquitous environmental presence, the extent of exposure, and the toxicity and disease states associated with it (Ray *et al*, 2014). It has been known that heavy metal ions such as Pb, Cu, Zn, Cd, Cr and Cd when present at an elevated level may enter our food chain through the vegetables grown on the banks of the river and accumulate in different parts of the body. This eventually causes reduced growth and impaired metabolism. Studies have shown that metal binding to the cell nucleus causes promutagenic damage by the production of reactive oxygen species. This leads to metal-mediated carcinogenicity and acute toxicity (Kasprzak, 1995; Cunningham, 1997).

Our study of water samples collected from the four different stations indicate high levels of Chromium at Yamuna River in Delhi (0.08mg/L). This value is much higher than the toxic levels recommended by the ICMR (0.05mg/L) and WHO (50µg/L). Chromium exposure has severe deleterious effect on chromosomal DNA and causes mutations in it. It is also responsible for the oxidative damage of proteins and the generation of free radicals. These free radicals act as carcinogens and lead to the formation of tumors (Dayan and Paine, 2001). Though the levels of Lead and Cadmium was found to be well within the acceptable range as per the health agencies, these metals are known to bio-accumulate in the system and attain toxic levels over a period of time. This would lead to severe health problems eventually.

Previous work has shown that when the heavy metals Ni, Cr & Cu are mixed in 1:1:1 ratio they show a synergistic toxic effect (Agarwal and Saxena, 2011). The sample analysis from the four chosen stations in this study indicate that approximately the same ratio, thereby indicating severe toxicity to the aquatic life and consequently to human beings as a result of bio-magnification.

The water samples collected from the four stations were analyzed with respect to the quality of the water indicator and oxygen consuming substances (pH, BOD and COD). The pH value is an indicator of the quality of water. The slight increase in pH of the river water (pH 7.72) at Yamuna river in comparison with the other three stations indicate the presence of carbonates of calcium and magnesium (Begum *et al.*, 2009). Our study shows that the Yamuna river has the unhealthiest levels of microorganisms which renders it unfit for human use. Municipal sewage disposal into the river can be the primary source of a high BOD, TSS and TDS. This in turn leads to high amount of coliform bacteria, which pose a severe hazard to human health. The discharge of various chemical effluents by the industries located around the river would lead to the high COD levels, further rendering it unfit for human use. Through our study, we thus show that the quality of our river water is steadily deteriorating and needs immediate attention.

5. Future Prospects

Through our study we have shown that three of the important rivers of India have a heavy metal load of varying concentrations. We would now like to extend this study to other major rivers of India, viz. Brahmaputra, Cauvery, Narmada, Tapti, Mahanadi, Krishna and Indus. This would give us a comprehensive view of the current situation of pollution levels in the river systems of the country. We would also like to extend this study to the aquatic life in the rivers in order to estimate the extent of biomagnifications occurring. The flora and fauna occurring on the banks of the rivers can also be analyzed for their content of heavy metals. Such a study would thus enable us obtain an insight into the extent of metal pollution of the rivers in India and the biomagnifications and bioaccumulation arising due to it.

6. Acknowledgement

The financial assistance provided by University of Delhi in University Innovation Research Project-SHC 203 to undertake this research work is gratefully acknowledged. We also acknowledge the students (Harshvardhan Singh, Astha Singla, Yashasvi Gupta, Manisha, Soumya Sharma, Karan Arora, Sweta, Anshu Sapra, Akashdeep Singh and Puru Sachan), working in the project for carrying out the experimental work.

References

- [1] Ray P. D., Yosim A. and Fry R. C. (2014). Incorporating epigenetic data into the risk assessment process for the toxic metals arsenic, cadmium, chromium, lead, and mercury: strategies and challenges. *Front. Genet.* 5:201 (Published online Jul 16, 2014).
- [2] Anand V. S. and Pandey J. (2014) Heavy metals in the midstream of the Ganges River: spatio-temporal trends in a seasonally dry tropical region (India), *Water International*, 1-13.
- [3] Malik D, Singh S, Thakur J, Kishore R, Kapur A and Nijhawan S, (2014) Microbial Fuel Cell: Harnessing Bioenergy from Yamuna Water. *International Journal of Science and Research*, 3 (6): 1076-1081.
- [4] Dey S. and Paul A. K. (2013). Hexavalent chromium reduction by aerobic heterotrophic bacteria indigenous to chromite mine overburden. *Brazilian J. Microbiol.* 44(1): 307-315.
- [5] Kaur S. (2012) Assessment of Heavy Metals in Summer & Winter Seasons in River Yamuna Segment Flowing through Delhi, India. *J. Environ. Eco.* 3(1): 149-165.
- [6] Agarwal A. and Saxena M. (2011). Assessment of toxic effect of industrial effluent on aquatic life in the River Gagan at Moradabad-India. *Pelagia Research Library Der Chemica Sinica.* 2 (2): 172-176.
- [7] Begum A., Ramaiah M., Harikrishna, Khan I. and Veena K. (2009). Heavy metal pollution and chemical profile of Cauvery river water. *e-J. Chem.* 6 (1): 47-52.
- [8] Bernard A. (2008) Cadmium and its adverse effects on human health. *Indian J. Med Res.* 128 (4):557-64.
- [9] Kar D., Sur P., Mandal S.K., Saha T. and Kole R.K (2008). Assessment of heavy metal pollution in surface water. *Int. J. Environ. Sci. and Technol.* 5: 119-124.
- [10] Das K. K., Das S. N. and Dhundasi S. A. (2008). Nickel, its adverse health effects and oxidative stress. *Indian J. Med. Res.* 128: 412-425.
- [11] Barbier O., Jacquillet G., Tauc M., Cougnon M. and Poujeol P. (2005). Effect of heavy metals on, and handling by, the kidney. *Nephron Physiol.* 99: 105-110.
- [12] Lenore S., Clesceria A., Greenberg E., and Andrew D.E., (2005). Standard methods for examination of water and wastewater. American Public Health Association.
- [13] Lenntech Water Treatment and Air Purification (2004) Water treatment. Lenntech, Rotterdamseweg, Netherlands (<http://www.excelwater.com/thp/filters/Water-Purification.htm>).
- [14] Bagchi D., Stohs S. J., Bernard W. O., Bagchi M. and Preus H. G. (2002). Cytotoxicity and oxidation mechanism of different forms of chromium. *Toxicol.* 180: 5-22.
- [15] Ali and Jain C. K. (2001). Pollution potential of Toxic metals in the Yamuna river at Delhi, *Indian J. Environ. Hydrol.* 9: 1-9.
- [16] Dayan A. D. and Paine A. J. (2001). Mechanisms of chromium toxicity, carcinogenicity and allergenicity: Review of the literature from 1985 to 2000. *Hum. Exp. Toxicol.* 20: 439.
- [17] Boxall A.B.A., Comber S.D., Conrad A.U., Howcroft J., Zaman N., (2000) Inputs, monitoring and fate modelling of antifouling biocides in UK estuaries. *Marine Pollution Bulletin* 40: 898-905.
- [18] Cunningham R. P. (1997) DNA repair: Caretakers of the genome? *Curr. Biol.* 7:576-579.
- [19] Kasprzak K. S. (1995) Possible role of oxidative damage in metal induced carcinogenesis. *Cancer Invest.*, 13:411-430.

- [20] USEPA (1999). National primary drinking water regulation. United States Environmental Protection Agency.
- [21] Schwartz J. (1994). Air pollution and daily mortality: A review and meta - analysis. *Environ. Res.* 64: 26-35.
- [22] Moore J. W. and Ramamoorthy S. (1984). Heavy metals in natural waters. Applied Monitoring and Impact Assessment. New York: Springer. 1-268.
- [23] Pillai, K. C. (1983). Heavy metals in aquatic environment, Water pollution and management (Varshey C.K.). New Delhi: Wiley, 74-93.
- [24] Dean J. G., Bosqui F. L. and Lannouette K. H. (1972). Removing Heavy Metals from Waste water. *Environ. Sci. Technol.* 6(6): 518-522.

Author Profile



Dr. Shashi Nijhawan (M.Sc & Ph.D), serving as **Principal, Shivaji College, University of Delhi**, Raja Garden, New Delhi since 2009. Earlier worked as Associate Professor in Department of Biochemistry, Deshbandhu College, University of Delhi. Her research areas of interest are Enzymology, Cell Biology and Molecular Biology. Published number of research papers in journals of National and International repute. Organized Workshops and International Conferences and presented no. of research papers in the National and International Conferences. She is the **Project Investigator** of an **University Innovation Project (SHC 203)** titled "**Biochemical assessment of toxic heavy metals in bacterial strain growing in the polluted water: A step towards Bioremediation**".



Dr. Darshan Malik (M.Sc & Ph.D), working as an Associate Professor in Department of Biochemistry, Shivaji College, University of Delhi. She is a recipient of **Erasmus Mundus (E.U) Scholarship for Academic Exchange Program** (Apr. 2009), worked for six months in the Department of Biology, Norwegian University of Science and Technology, Trondheim, **Norway**. Her research areas of interest are Cell Biology, Molecular Biology, Human Physiology, Environment and Disaster management. Published several research papers in journals of National and International repute. She has edited / published two books. Has organized Workshops and International Conferences. Presently supervising Ph.D. students registered with Guru Gobind Singh Indraprastha University, New Delhi, India. She is involved in research work on Bioremediation and guiding Undergraduate students of **University Innovation Project (SHC 203)**. She is also involved in the Environment & Disaster management plan of Delhi.



Dr. Amarjeet Kaur (M.Sc & PhD), serving as Professor in University School of Environment Management (USEM), **Guru Gobind Singh Indraprastha University, New Delhi**, India. She has more than 22 years of experience in research and training. Her areas of specialization are Ecology & sustainable development, Environmental Biotechnology and Geo-information System. She has worked as Principal /Co-Principal Investigator of various International/National projects related to Environment Management. She has visited USA, UK, Bangkok and Japan in connection with study exchange programmes and conferences. She was also involved in Environmental Management Plan of Delhi using Remote Sensing Technology and GIS. Presently she is leading a team of experts in reviewing/upgrading the District Disaster Management plan of Delhi as Director, Centre for Disaster Management of GGS Indraprastha University. She has twenty years experience of providing consultancy to various Industries.



Dr. Sunita Singh (M.Sc & Ph.D), working as an Assistant Professor in the Department of Biochemistry of Shivaji College, University of Delhi. Her research areas of interest are Nano-biocatalysts and their applications, Chemical modification and chemical cross linking of proteins and Industrial applications of enzymes. She has twenty research publications in National and International journals. Participated & presented papers /posters in National & International Conferences. She is the recipient of **Woman Scientists Scheme** (SR/WOS-A/LS-181/2007) of Department of Science & Technology (DST) for Research in Basic/Applied Sciences. She has availed **Young Scientist Award of DST** (under SERC FAST Track Scheme for Young Scientists 2001-02) and **Senior Research Associate (Pool Officer) of Council of Scientific and Industrial Research (CSIR)** (Pool no. 7446 A).



Jayita Thakur (M.Sc), is working as an Assistant Professor in the Department of Biochemistry of Shivaji College, University of Delhi. She is presently involved in University of Delhi Innovation Project along with the co-authors of this paper. She is pursuing her PhD from the **National Institute of Immunology**, New Delhi and is on the verge of completion. She has attended numerous National and International conferences, workshops and symposia. The main areas of interest include molecular biology, cell biology and genetics. She is also actively involved in research work, with interest and specialized skills in tools and techniques related to Molecular Biology, Virology and Cell Biology.



Dr. Raj Kishore Singh (M.Sc & PhD), working at University School of Environment Management (USEM), **Guru Gobind Singh Indraprastha University, New Delhi**, India. Research areas of interest are Industrial Pollution, Solid Waste Management, Contaminated Site Management, and Wastewater Treatment Technology. Has 10 research publications to his credit in peer reviewed journals. Twelve years of experience as co-coordinator and project manager of several consultancy projects. Presently in the Editorial Board of IJRASET and IJRAASE.