

Applications of Generalized Two Dimensional Fractional Fourier transform

V. D. Sharma

¹ Department of Mathematics, Arts, Commerce & Science College, Amravati, 444606 (M.S.) India

Abstract: As the one dimensional (1-D) Fourier transform can be extended into 1-D fractional Fourier transform (FrFT), we can also generalize the two-dimensional (2-D) Fourier transform. Recently several properties of FrFT have been developed by generalizing the properties of the ordinary Fourier transform (FrFT). In this paper Applications of Generalized two –dimensional fractional Fourier transform is presented.

Keywords: Fourier transform, Fractional Fourier transform, generalized function, signal processing optics.

1. Introduction

Fractional Fourier transform (FrFT) is a generalization of the ordinary FT. FrFT was first introduced as a way to solve certain classes of ordinary and partial differential equations arising in quantum mechanics [1]. FrFT has established itself as a powerful tool for the analysis of time varying signals, especially in optics [2]. FrFT has found applications in areas of signal processing such as repeated filtering, fractional convolution and correlation, beam forming, optional filter, convolution, filtering and wavelet transforms, time frequency representation. In every area in which Fourier transform and frequency domain concepts are used, then exists the potential for generalization and improved by using the FrFT[3].

Recently FrFT independently discussed by lots of researchers. Ozaktas etal [4] had given applications of FrFT in optics and signal processing. Alieva T [5] has developed fractional transforms in optical Information processing. Bultheel A. et al in [6] had given recent developments in the theory of fractional Fourier & linear canonical transforms. Djurovic [7] etal had established FrFT as a signal processing tool. Salazar F. [8] had given a new introduction to the FrFT and its applications.

In the present work generalization of two dimensional Fractional Fourier transform in presented. Application of two-dimensional Fractional Fourier transform in the form of examples are given. Two dimensional FrFT of some functions are obtained.

2. Preliminaries

2.1 Fourier Transform

The Fourier transform of function $x(t)$ in defined as

$$X(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x(t) e^{j\omega t} dt$$

And inverse is given by

$$x(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} X(\omega) e^{-j\omega t} d\omega$$

2.2 One dimensional fractional fourier transform

One dimensional fractional Fourier transform with parameter α of $f(x)$ defined as

$$FrFT\{f(x)\} = F_{\alpha}(u) = \int_{-\infty}^{\infty} f(x) k_{\alpha}(x, u) dx \quad (1.1)$$

where the kernel

$$k_{\alpha}(x, u) = \sqrt{\frac{1-i\cot\alpha}{2\pi}} e^{\frac{i}{2\sin\alpha}[(x^2+u^2)\cos\alpha-2(xu)]} \quad (1.2)$$

3. Distributional Two Dimensional Fractional Fourier Transform

3.1 Conventional Two- dimensional Fractional Fourier transforms

The two dimensional fractional Fourier transform with parameter α of $f(x, y)$ denoted by $FRFT\{f(x, y)\}$ performs a linear operation given by the integral transform

$$FRFT\{f(x, y)\} = F_{\alpha}\{f(x, y)\}(u, v) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) K_{\alpha}(x, y, u, v) dx dy, \quad (3.1)$$

Where the kernel,

$$K_{\alpha}(x, y, u, v) = C_{1\alpha} e^{iC_{2\alpha}[(x^2+u^2+y^2+v^2)\cos\alpha-2(xu+yv)]}, \\ C_{1\alpha} = \sqrt{\frac{1-i\cot\alpha}{2\pi}}, C_{2\alpha} = \frac{1}{2\sin\alpha} \quad (3.2)$$

3.2 The testing function space E

An infinitely differentiable complex values smooth function ϕ on R^n belongs to $E(R^n)$ if for each compact set $I < S_{a,b}$ where,

$$S_{a,b} = \{x, y: x, y \in R^n, |x| \leq a, |y| \leq b, a > 0, b > 0\}, I \in R^n,$$

$$\gamma_{E,p,q}(\phi) = \sup_{x,y \in I} |D_{x,y}^{p,q} \phi(x, y)|$$

$< \infty$, where $p, q = 1, 2, 3, \dots$

Thus $E(R^n)$ will denote the space of all $\phi \in E(R^n)$ with support contained in $S_{a,b}$.

Note that the space E is complete and therefore a Frechet space. Moreover, we say that f is a fractional Fourier transformable if it is a member of E^* , the dual space of E .

3.3 Distributional Two-dimensional fractional Fourier transform

The two distributional two-dimensional fractional Fourier transform of $f(x, y) \in E(R^n)$ can be defined by,

$$FRFT\{F(X, Y)\} = F_\alpha(u, v) = \langle f(x, y), K_\alpha(x, y, u, v) \rangle \dots \dots (3.1)$$

When $K_\alpha(x, y, u, v) = C_{1\alpha} e^{iC_{2\alpha}[(x^2+u^2+y^2+v^2)\cos\alpha - 2(xu+yv)]}$

$$, C_{1\alpha} = \sqrt{\frac{1-icot\alpha}{2\pi}}, C_{2\alpha} = \frac{1}{2\sin\alpha} \dots \dots (3.2)$$

The right hand side of (3.1) has a meaning as the application of $f \in E^*$ to $K_\alpha(x, y, u, v) \in E$.

It extended to the complex space as an entire function given by

$$FRFT\{F(X, Y)\} = F_\alpha(g, h) = \langle f(x, y), K_\alpha(x, y, g, h) \rangle \dots \dots (3.3)$$

Where

$$K_\alpha(x, y, g, h) = C_{1\alpha} e^{iC_{2\alpha}[(x^2+g^2+y^2+h^2)\cos\alpha - 2(xg+yh)]}$$

The right hand side of (3.3) is meaningful because for each $g, h \in c^n, K_\alpha(x, y, u, v) \in E$, as a function of x, y .

4. Examples on Generalized 2DFrFt

4.1- Prove that [2DFrFT(1)](u, v)

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{cot\alpha} e^{\frac{i\pi}{2} e^{\frac{i}{2} \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right] (u^2+v^2)}}$$

Proof:

$$2DFrFT\{f(x, y)\}(u, v) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} C_{1\alpha} e^{\frac{i}{2\sin\alpha}[(x^2+u^2+y^2+v^2)\cos\alpha - 2(xu+yv)]} f(x, y) dx dy$$

$$[2DFrFT(1)](u, v) = C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha}$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{\frac{i}{2}(x^2+y^2)cot\alpha - i(xu+yv)cosec\alpha} 1 dx dy$$

$$[2DFrFT \delta(x - a, y - b)](u, v) = C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha} e^{\frac{i}{2}(a^2+b^2)cot\alpha - i(au-bv)cosec\alpha} = C_{1\alpha} e^{\frac{i}{2\sin\alpha}(a^2+u^2+b^2+v^2)\cos\alpha - 2(au-bv)}$$

4.3- Prove that: [2DFrFT e^{i(ax²+by²)}](u, v) =

$$\sqrt{\frac{2\pi(1-cot\theta)}{(cot\alpha+2a)(cot\alpha+2b)}} e^{\frac{i}{2}(u^2+v^2)cot\alpha} e^{icosec^2\alpha \left(\frac{u^2}{2cot\alpha+4a} + \frac{v^2}{2cot\alpha+4b} \right)}$$

Proof:

$$[2DFrFT \{f(x, y)\}](u, v) = C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{\frac{i}{2}(x^2+y^2)cot\alpha - i(xu+yv)cosec\alpha} f(x, y) dx dy$$

$$[2DFrFT e^{i(ax^2+by^2)}](u, v) =$$

$$C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha} \int_{-\infty}^{\infty} e^{ix^2 \left[\frac{cot\alpha}{2} + a \right] - iuxcosec\alpha} dx$$

$$\int_{-\infty}^{\infty} e^{iy^2 \left[\frac{cot\alpha}{2} + b \right] - ivycosec\alpha} dy$$

$$\int_{-\infty}^{\infty} e^{iap^2+ibp} dp = \frac{e^{\frac{i\pi}{4}\sqrt{\pi}}}{\sqrt{a}} e^{\frac{ib^2}{4a}} \dots (4.1)$$

where, $a = \frac{cot\alpha}{2}, b = -ucosec\alpha$

$$= C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha} \left[\frac{e^{\frac{i\pi}{4}\sqrt{\pi}}}{\sqrt{\frac{cot\alpha}{2}}} e^{\frac{i(-ucosec\alpha)^2}{4\frac{cot\alpha}{2}}} \right] \left[\frac{e^{\frac{i\pi}{4}\sqrt{\pi}}}{\sqrt{\frac{cot\alpha}{2}}} e^{\frac{i(-vcosec\alpha)^2}{4\frac{cot\alpha}{2}}} \right]$$

$$= C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha} \left[\frac{e^{\frac{i\pi}{4}\pi}}{\frac{cot\alpha}{2}} \right] e^{\frac{iu^2cosec^2\alpha + iv^2cosec^2\alpha}{2cot\alpha}}$$

$$= \frac{\sqrt{1-icot\alpha}}{cot\alpha} \sqrt{2\pi} e^{\frac{i\pi}{2}} e^{\frac{i}{2}u^2 \left[\frac{cosec\alpha}{\sin\alpha} + \frac{1}{\sin\alpha\cos\alpha} \right]} e^{\frac{i}{2}v^2 \left[\frac{cosec\alpha}{\sin\alpha} + \frac{1}{\sin\alpha\cos\alpha} \right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2}u^2 \left[\frac{1+\cos^2\alpha}{\sin\alpha\cos\alpha} \right]} e^{\frac{i}{2}v^2 \left[\frac{1+\cos^2\alpha}{\sin\alpha\cos\alpha} \right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2}u^2 \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right]} e^{\frac{i}{2}v^2 \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2} \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right] (u^2+v^2)}$$

[2DFrFT(1)](u, v)

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2} \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right] (u^2+v^2)}$$

4.2- Prove that: [2DFrFT δ(x - a, y - b)](u, v) =

$$C_{1\alpha} e^{\frac{i}{2\sin\alpha}[(a^2+u^2+b^2+v^2)\cos\alpha - 2(au+bv)]}$$

$$[2DFrFT \{f(x, y)\}](u, v)$$

$$= C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)cot\alpha}$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{\frac{i}{2}(x^2+y^2)cot\alpha - i(xu+yv)cosec\alpha} f(x, y) dx dy$$

Taking $a = \left[\frac{cot\alpha}{2} + a \right], b = -ucosec\alpha$ etc

Therefore using equation (4.1)

$$[2DFrFT e^{i(ax^2+by^2)}](u, v)$$

$$= \sqrt{\frac{1-icot\theta}{2\pi}} e^{\frac{i}{2}(u^2+v^2)cot\alpha}$$

$$\frac{e^{\frac{i\pi}{4}\sqrt{\pi}}}{\sqrt{\frac{cot\alpha}{2}+a}} e^{\frac{i(-ucosec\alpha)^2}{4\left(\frac{cot\alpha}{2}+a\right)}} \frac{e^{\frac{i\pi}{4}\sqrt{\pi}}}{\sqrt{\frac{cot\alpha}{2}+b}} e^{\frac{i(-vcosec\alpha)^2}{4\left(\frac{cot\alpha}{2}+b\right)}}$$

$$= \sqrt{\frac{1-icot\theta}{2\pi}} \frac{2\pi e^{\frac{i\pi}{2}}}{\sqrt{(cot\alpha+2a)\sqrt{(cot\alpha+2b)}}}$$

$$e^{\frac{i}{2}(u^2+v^2)cot\alpha} e^{icosec^2\alpha \left(\frac{u^2}{2cot\alpha+4a} + \frac{v^2}{2cot\alpha+4b} \right)}$$

$$= \sqrt{\frac{2\pi(1-cot\theta)}{(cot\alpha+2a)(cot\alpha+2b)}} e^{\frac{i}{2}(u^2+v^2)cot\alpha}$$

$$e^{i \operatorname{cosec}^2 \alpha \left(\frac{u^2}{2 \cot \alpha + 4a} + \frac{v^2}{2 \cot \alpha + 4b} \right)}$$

4.4- Prove that: $[2DFrFT e^{i(ax+by)}](u, v)$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2\cot\alpha}}$$

$$e^{\{(a^2+b^2)-2\operatorname{cosec}\alpha(au+bv)\} + \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right] (u^2+v^2)}$$

Proof-

$$[2DFrFT \{f(x, y)\}](u, v) = C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)\cot\alpha}$$

Therefore using equation (4.1)

$$[2DFrFT e^{i(ax+by)}](u, v) = \sqrt{\frac{1-icot\phi}{2\pi}} e^{\frac{i}{2}(u^2+v^2)\cot\alpha} e^{\frac{i\pi}{4}\sqrt{\pi}} e^{\frac{i(a-u\operatorname{cosec}\alpha)^2}{4\left(\frac{\cot\alpha}{2}\right)}} e^{\frac{i\pi}{4}\sqrt{\pi}} e^{\frac{i(b-u\operatorname{cosec}\alpha)^2}{4\left(\frac{\cot\alpha}{2}\right)}}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2}\left[u^2\cot\alpha + \frac{(a-u\operatorname{cosec}\alpha)^2}{\cot\alpha}\right]}$$

$$e^{\frac{i}{2}\left[v^2\cot\alpha + \frac{(b-v\operatorname{cosec}\alpha)^2}{\cot\alpha}\right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2}\left[u^2\cot\alpha + \frac{a^2-2au\operatorname{cosec}\alpha}{\cot\alpha} + \frac{u^2\operatorname{cosec}^2\alpha}{\cot\alpha}\right]}$$

$$e^{\frac{i}{2}\left[v^2\cot\alpha + \frac{b^2-2bv\operatorname{cosec}\alpha}{\cot\alpha} + \frac{v^2\operatorname{cosec}^2\alpha}{\cot\alpha}\right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha} e^{\frac{i\pi}{2}} e^{\frac{i}{2}\left[\frac{a^2-2au\operatorname{cosec}\alpha}{\cot\alpha} + \frac{u^2(3+\cos 2\alpha)}{\sin 2\alpha}\right]}$$

$$e^{\frac{i}{2}\left[\frac{b^2-2bv\operatorname{cosec}\alpha}{\cot\alpha} + \frac{v^2(3+\cos 2\alpha)}{\sin 2\alpha}\right]}$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha}$$

$$e^{\frac{i\pi}{2}} e^{\frac{i}{2\cot\alpha}\left[\frac{(a^2+b^2)-2\operatorname{cosec}\alpha(au+bv)}{\cot\alpha} + \frac{(3+\cos 2\alpha)(u^2+v^2)}{\sin 2\alpha}\right]}$$

$$[2DFrFT e^{i(ax+by)}](u, v)$$

$$= \frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha}$$

$$e^{\frac{i\pi}{2}} e^{\frac{i}{2\cot\alpha}\left[\frac{(a^2+b^2)-2\operatorname{cosec}\alpha(au+bv)}{\cot\alpha} + \frac{(3+\cos 2\alpha)(u^2+v^2)}{\sin 2\alpha}\right]}$$

Generalized 2DFrFT

S.N.	$f(x, y)$	$\{2DFrFT f(x, y)\}(u, v)$
1	1	$\frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha} e^{i\pi/2}$ $e^{\frac{i}{2}(3\operatorname{cosec}^2\alpha + \cot 2\alpha)(u^2+v^2)}$
2	$\delta(x-a, y-b)$	$C_{1\alpha} e^{\frac{i}{2\sin\alpha}\left[(a^2+u^2+b^2+v^2)\operatorname{cosec}\alpha - 2(au+bv)\right]}$
3	$e^{i(ax^2+by^2)}$	$\sqrt{\frac{2\pi(1-\cot\phi)}{(cot\alpha + 2a)(cot\alpha + 2b)}} e^{\frac{i}{2}(u^2+v^2)\cot\alpha}$ $e^{i \operatorname{cosec}^2 \alpha \left(\frac{u^2}{2 \cot \alpha + 4a} + \frac{v^2}{2 \cot \alpha + 4b} \right)}$
4	$e^{i(ax+by)}$	$\frac{\sqrt{2\pi(1-icot\alpha)}}{\cot\alpha}$ $e^{\frac{i\pi}{2}} e^{\frac{i}{2\cot\alpha} e^{\{(a^2+b^2)-2\operatorname{cosec}\alpha(au+bv)\}}}$ $+ \left[\frac{3+\cos 2\alpha}{\sin 2\alpha} \right] (u^2+v^2)$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{\frac{i}{2}(x^2+y^2)\cot\alpha - i(xu+yv)\operatorname{cosec}\alpha} f(x, y) dx dy$$

$$[2DFrFT e^{i(ax+by)}](u, v)$$

$$= C_{1\alpha} e^{\frac{i}{2}(u^2+v^2)\cot\alpha} \int_{-\infty}^{\infty} e^{ix^2\cot\alpha - iux\operatorname{cosec}\alpha + iax} dx$$

$$\int_{-\infty}^{\infty} e^{iy^2\cot\alpha - ivy\operatorname{cosec}\alpha + iby} dy$$

Taking $a = \frac{\cot\alpha}{2}$, $b = a - u\operatorname{cosec}\alpha$ etc

5. Conclusion

In the present work generalization of two dimensional Fractional Fourier transform is presented. Some applications of 2DFrFt is obtained.

References

- [1] B.N.Bhosale, Integral transformation of Generalized functions. Discovery publishing hour, 2005.
- [2] H.M. Ozaktes, A. Zalevsky, M.A. Katay, the FRFT with Application in optics and signal processing, John Wiley, Chichester New York, USA, 2001.
- [3] Saleem I. Raza S. M. et.al., Fractional Fourier integral transform and fractional fourier sine and cosine transform, sci. Int. 24 (3) 233-238,2012
- [4] Zemanian A.H., Generalized Integral transform, Interscience Publication, New York, 1968.
- [5] Alieva T, Bastiaans M. etal, Fractional Transforms in optical Information processing. Euro. J. Appl. Sig. Proc. 10, (2005) 1498-1519.
- [6] Bultheel A. etal, Recent developments in the theory of the fractional Fourier and linear canonical transforms, Preprint,(2004) 1-33.
- [7] Sejdic E. and Djurovic I. etal, FRFT as a signal processing tool: An overview of recent developments, singal processing 91, (2011)1351-1369.
- [8] Felipe Salazar, An Introduction to the Fractional Fourier transform and its applications, Dec. 7,(2011), 1-8.
- [9] V.D.Sharma, Generalization of Two-dimensional Fractional Fourier transforms, Asian Journal of current Engineering and Maths, 46-48, 2012.
- [10] V.D.Sharma, operation transform Formulae on Generalized FrFT, processing of Int. conf. in Recent trends in Inf. Tech. and Comp. Science published in Int. J. of computer applications 15-18, 2012.
- [11] V.D. Sharma, Generalized Two-Dimensional Fractional Fourier Transform and Inversion Formula, Journal of Fractional Calculus and Applications, Vol. 4(2) July 2013, pp. 191- 195.

- [12] V. D. Sharma, Operational calculus on generalized two-Dimensional Fractional Fourier transform, International Journal of Engineering and Innovative Technology (IJEIT) Volume 3, Issue 2, August 2013 ,253-256.
- [13] V. D. Sharma, Modulation and Parsvels Theorem For Generalized Fractional Fourier Transform, Int. J. Engineering research and application , vol 3, issue 5, sept-octo 2013, PP 1463-1467.

Author Profile

Dr. V. D. Sharma is currently working as an Assistant professor in the department of Mathematics, Arts, Commerce and Science College, Kiran Nagar, Amravati-444606 (M.S.) India. She has got 18 years of teaching and research experience. She has obtained her Ph.D. degree in 2007 from SGB Amravati University Amravati. Her field of interest is Integral Transforms. Six research students are working under her supervision. She has published more than 60 research articles.

P. B. Deshmukh is an Assistant professor in the department of Mathematics, IBSS College of Engineering, Amravati 444602 (M.S.) India. She has obtained her master degree in 2006 from Sant Gadage Baba Amravati University, Amravati. She has got 7 years of teaching experience. She has 6 research articles in journals to her credit.

