

Effect of Parsley Extract on Protein Content and Histology of Submandibular Glands of Naturally Aged Male Mice

S. N. Khandare¹, N. K. Khandare²

¹Department of Zoology, Vidnyan Mahavidyalaya, Sangola, Dist- Solapur, (MS) -413 307, India

²Department of Botany, Krantisinh Nana Patil College, Walwa, Dist- Sangli, (MS) - 416 313, India

Abstract: *The aging is generally characterized by the declining in ability to respond to stress, increasing homeostatic imbalance and increased risks of diseases. Aging promotes free radical formation in the cell. The free radicals formed due to aging or various reasons are scavenged by antioxidants. In present studies antioxidant rich plant, viz- parsley (*Petroselinum crispum*. mill), was used to assess the free radicals generated during aging. Three groups of mice (Control, Naturally aged, Parsley receiving) were used during the experiments. Study was restricted to only submandibular glands. Histological sections of glands were stained with haematoxyline and eosine. Protein content was estimated in the experiment. Study shows damage in submandibular glands in naturally aged mice as compare to young control. Parsley corrected the histological structure of submandibular glands as well as protein content of submandibular glands in naturally aged animal.*

Keywords: Aging, submandibular glands, parsley, protein.

1. Introduction

The salivary glands are readily accessible and well characterized. Therefore, they are useful tools for the study of the normal aging process and the impact of stress on organ reserve and secretory functions (Baum *et al*, 1992). A common generalization associated with aging is that salivary gland function is altered (Storer, 1978) and diminished output results in dental carries, altered mucosal integrity and impaired taste and agglutination. Salivary glands dysfunction has been traditionally attributed to old age (Thomson *et al*, 1999). Submandibular glands are termed as acinar cells, together formed called acini, which secrete mainly glycoproteins. Acini are surrounded by myoepithelial cells and supported by parenchyma. There are various ducts as intercalated ducts, striated duct and granular ducts, which are also called as granular convoluted tubules. These ducts have been considered to be the sites of formation of many enzymes like Kallikrein (Hojima *et al*. 1977) Proteases (Sreebny and Meger 1964), nerve growth factor (Schwah *et al*, 1976), epidermal growth factor (Cohen, 1962, Young and Van Lennep 1978) and various mesodermal growth factors (Weimer and Haraguchi 1975). Salivary proteins are mucins these are effective lubricant, they control permeability of mucosal surface, limit penetration of potential irritants and toxins to mucous cells, protect cell membranes against proteases generated by bacteria, and regulate colonization of oral cavity by bacteria and viruses (Mandel, 1992). Parsley is considered to be one of the highest sources of flavonol glycosides (Kreuzaler *et al*, 1973). It has been shown to possess remarkable histological correction in sublingual glands (Khandare *et al*, 2013); anti inflammatory, antioxidant and anti carcinogenic property (Patel *et al*, 2007).

2. Material and Methods

2.1 Animals

Adult male albino mice (*Mus musculus*) of 23 weeks were used for the present investigation. The breeding pairs were obtained from Hindustan Antibiotics, Pune. They were reared and kept in the animal house with the maintenance of constant temperature (about 25 to 30°C) and light and dark cycles. They were supplied with Amrut Mice feed (Pranav Agro Industries, Sangli) and water ad libitum. Animals were randomly assigned to the following three groups.

2.2 Control Group

Adult male mice (Age 23 week's old and weight 42 ± 0.6 gm) was treated as control.

2.3 Naturally aged group:

Naturally an aged male mouse (Age 76 week's old and weight 38.66 ± 1.032 gm) was used.

2.4 Parsley receiving old male mice group:

Old male mice (Age 76 week's old weight 38.66 ± 1.032 gm) were injected parsley 40 mg / kg body weight / day subcutaneously for 20 days.

2.5 Preparation of plant extract

Fresh Parsley (*Petroselinum crispum*) mill was obtained from the market. Green fresh leaves were separated and washed properly with water and rinsed with distilled water. Washed leaves were blotted properly with blotting paper and kept for drying in the shade. The dried leaves were crushed, powdered and sieved. The powder was soaked in double

distilled alcohol (ethanol) for 72 hours for extraction. The mixture of ethanol and powder of parsley leaves was filtered. The alcohol was evaporated by using high speed (Buchi Type) vacuum evaporator to obtain a thick paste like extract. This extract was collected by spatula and stored in a glass bottle. It was kept refrigerated (at 4°C) for further use.

2.6. Histology

The animals were sacrificed by cervical dislocation. The salivary glands were excised out. They were weighed and kept in the freezer for freezing and thawing and used for protein estimation. For the histology of sublingual glands, tissue were fixed in 2% CAF (Calcium acetate formalin) + 2% calcium acetate in 10% formalin fixative for 24 hours at 4°C. The tissue was washed in running tap water for 24 hours dehydrated through alcohol grades, cleared in xylene and embedded in paraffin. The sections were at a thickness of 6 microns and stained with eosin and haematoxyline.

2.7. Estimation of Proteins: (Lowry et al, 1931)

2.7.1. Preparation

1. Reagent A – 2% Na₂CO₃ in 0.1 N NaOH.
2. Reagent B – 0.5% CuSO₄ in 1% freshly prepared Na⁺, K⁺ tartarate.
3. Reagent C – 50 ml A+1 ml B (made freshly at the time of use).
4. Reagent D – Folin c (90% Ciacolteu Phenol) reagent.

2.7.2. Procedure

The additions for the estimation of proteins were made as follows.

	Blank	Sample
Distilled Water	4 ml	3.5 ml
Sample		0.5 ml
Reagent C	5.5 ml	5.5 ml

The tubes were shaken well and kept for 10 min, 0.5 ml reagent D was added mixed well and kept for 30 min. The optical density was measured 660 nm. The final colour production is the result for biuret reaction of protein with copper ions in an alkaline medium at reduction of phosphomolybdc phosphotungstic reagent by the tyrosin and tryphophan present in treated protein.

2.7.3. Calculation:

The protein concentration 1 mg tissue was determined using standard graph of bovine serum albumin.

2.8. Statistical analysis: Statistical analysis was performed by Student ‘t’-test., P<0.001

3. Result

Fig no. “a.” describes the cross section of submandibular glands of adult mice (Age 23 weeks) was having well formed acini (AC) and convoluted granular tubules (GCT) and ducts were clear and darkly stained.




Figure (a): Control group, cross section of Submandibular gland]

In naturally aged mice, Fig no. “b.” (Age 76 weeks) Submandibular glands, number and size of acini were reduced. Organization of acini and granular convoluted tubules with respect to one another was lost. Ducts were disorganized cell layer of the duct was thin, reduced staining reactivity of acini (AC) and GCT was lost and also their integrity.




Figure (b): Naturally aged group, Cross section of submandibular gland]

Fig. no. “c” depicts the structure of submandibular gland of parsley receiving mice nuclear staining reactivity of both the granular cells (GC) and acini (AC) was increased as well as architecture acini became normal, but not of the granular convoluted tubules.




Figure (c): Parsley receiving group, Cross section of submandibular gland]


Table no. 1; graph.1, shows that, Protein content estimated in submandibular glands in control group was 430.18 ± 0.934 µg/mg weight tissue it was significantly (1:2 p<0.001) decreased in naturally aged mice. Naturally aged mice

receiving parsley group showed increase in protein content. The increase was significant (2:3 $p < 0.001$) compared to naturally aged mice

Table 1: Effect of *Petroselinum crispum* extract on protein content ($\mu\text{g}/\text{mg}$) of submandibular gland of naturally aged male mice

Sr. No.	Animal Group	Age in weeks	Protein content $\mu\text{g}/\text{mg}$	
			Mean \pm SD	Statistical significance
1	(6) Control	23	430.18 \pm 0.934	1 : 2 t = 55.973
2	(6) Naturally aged	76	197.89 \pm 2.2250	($p < 0.001$) 2 : 3
3	(6) Parsley receiving recovery	76	376.83 \pm 1.9714	T = 39.509 ($p < 0.001$)

Values are mean \pm SD $P < 0.001$, highly significant, Values in the parenthesis denote number of animals.



Graph 1: Protein content of submandibular gland of naturally aged male mice

4. Discussion

The purpose of this study was to examine the effect of dietary antioxidants (parsley) on stressed salivary glands. Dietary antioxidants are polyphenolic compounds which are ubiquitously present in foods of plants origin. Quercetin is a strong antioxidant studied against the aging and free radical toxicity in brain (Durate *et al*, 2001; Molina *et al*, 2003), heart (Mohanty *et al*, 2004) and other organs (Gurento *et al*, 2000). According to Giugliano *et al*, 2000; sufficient supply with antioxidants from diet might help to prevent the occurrence of pathological changes associated with oxidative stress. There are several antioxidants suggested in the literature to prevent or acceptable aging effect. Several attempts have been made to knock out antioxidants vitamin E (Zhang *et al*, 2002), vitamin C, melatonin (Reiter *et al*, 1996), lipoic acid (Packer *et al* 1997; Kalia *et al*, 2013) in mice and other animals. *Petroselinum crispum* possess strong antioxidant properties (Hirano *et al* 2001). It contains rich amount of vitamins C and A (Pattision *et al*, 2004). It is one of the richest sources of flavonoids, especially quercetin

(Fejes *et al*, 1998). *Petroselinum crispum* improved the histological structure of sublingual glands in naturally aged male mice (Khandare *et al*, 2013) In present investigation the *Petroselinum crispum* shows promising results in proteins and corrected histological structure of submandibular glands.

5. Conclusion

Present experiments investigated that, Parsley (*Petroselinum crispum*), improved the histological structure and showed recovery in protein contents of Submandibular glands in naturally aged male mice.

6. Abbreviations

GCT - Convolved granular tubules

AC - Acinar cells

Dc - Duct

References

- [1] B.J. Baum, J. A. Ship, and A.J. Wu, "Salivary gland function and aging a model for studying the interaction of aging and systemic disease." Crit. Rev. oral Biol. Med, (4) pp.53-64 1992.
- [2] D. Giugliano, "Dietary antioxidants for cardiovascular prevention." Nutr. Metab Cardio Vasc. Dis, 10C1,pp.38-44,2000.
- [3] D. J. Pattision, A. J. Silman, Good Son, , M. Lunt, D. Bunn,R. Luben, A. Welch, Bingham, K.T . Khaw ,N. Day, and S..D. Jymmon, "Vitamin C and the Risk of developing inflammatory polyarthritis: Perspective nested case control study." An. Rheum. Dis, 63(7), pp.843-847, 2004.
- [4] D. Patel,, S. Shukla, "Apigenin and cancer chemoprevention progress potential and promise (Review)" Int. J. Oncol 30(1),pp.233-245,2007.
- [5] E. W. Van Lennep, J. A. Young, "Transport in salivary and salt glands." Part II: Salt glands. In: G Giebisch. D C Tosteson, H H Ussing (eds), *Membrane Transport in Biology Volume 4B,Transport Organs*. Springer Verlag, Berlin, pp. 675-692, 1979.
- [6] F. Kreuzaler and K. Hahbrock, "Flavonoid glycosides from illuminated cell suspension culture of *Petroselinum hortense*." Phytochemistry, (12), pp.1149-1152, 1973.
- [7] I. D. Mandel, C.E. Barr, L. Turgeon, "Longitudinal study of parotid saliva in HIV-1 infection." J oral pathol Med. (21),pp. 209-213,1992.
- [8] J. Durate, M. Galisteo , M.A. Ocete, F. Petez Vizeaino, A. Zarzuelo and Tamargo, "Effect of chronic quercetin treatment on hepatic oxidative state of spontaneously hypertensive rats." J. Mol. Cell. Biochem.,221(1-2),pp.155-160, 2001.
- [9] K. Kalia, H. M. Chiragni, and P. P. Sood. "Effect of Antioxidants (alpha-lipoic acid and bamboo shoot extract either alone or in combination) in Lead induced oxidative stressed animals" J. Cell Tissue Res., 13(1),pp. 3431-3438 ,2013.

- [10] L. Guarente and C. Kenyon, "Genetic pathways that regulate ageing in model organisms." *Nature*, (408), pp. 255-262, 2000.
- [11] L. M. Sreebny, J. Meyer, "Hormones inanition and salivary glands. In salivary glands and their secretion" (L. M. Greebng and J. Meyer (eds) prgamon oxford, pp. 83-103, 1964.
- [12] L. Packer, H. J. Tristchler, and K. Wesset, "Neuroprotection by the metabolic antioxidant lipoic acid." *Free rad. Biol. Med.* 22(1-2), pp.339-378, 1997.
- [13] M. E. Schwab, K. Stockel, H. Thoenen, "Immunocytochemical localization of nerve f. growth factor (NGF) in the submandibular gland of adult mice by light and electron microscopy." *Cell tiss. Res.*, (169), pp. 289-299, 1976.
- [14] M. F. Molina, I. Sanchez Rues, I. Iglesias and Benedi, "Quercetin a flavonoid antioxidant, prevents and protects against ethanol-induced oxidative stress in mouse liver." *J. Biol. Pharm. Bult.* 26(10), pp. 1398-1402, 2003.
- [15] O. H. Lowry, N. J. Rosenbrough, A. L. Farr, and R. J. Randall, "Protein measurement with the folin-phenol reagent." *J. Bil. Chem.*, (193), pp.265-275, 1951.
- [16] R. Hirano, W. Sasa moto and A. Mastsumoto , "Antioxidant ability of various flavonoids against DPPH radicals and LD1 oxidation" *J. Nutr. Sci. Vitaminol (Tokyo)*, 47(6), pp. 357-362 , 2001.
- [17] R. J. Reiter, M. Z. Pablos, T. T. Agapita, and J. M. Guerrero, "Melatonin in the context of the free radical Theory of aging". *Ann. New York, Acad. Sci.* (786), pp.362-378 ,1996.
- [18] R. Storer, The oral tissues. In.: *Text Book of Geriatric Medicine and Gerontology*, (Brocklehurst, J. C. ed) London, Churchill Livingstone, pp. 330-340, 1978.
- [19] S. Cohen, "Isolation of a mouse submaxillary gland accelerating incisor eruption and eyelid opening in the animal." *J. Biol. Chem.*, pp.237-1555,1962.
- [20] S. Fejes, A. Krey, A. Blazovics, A. Lugasi, E. Lemberkovics, G. Peteri, and E. Szoke, "Investigation of the in vitro antioxidant effect of *Petroselinum crispum*." *Acta pharmtung.*, 68(3), pp.150-156,1998.
- [21] S. M. Zhang, M.A. Herman, H. Chen, D. Spiegelman, W. C. Willett, and A. Ascherio, "Intake of Vitamin E and C Carotenoids, Vitamin Supplements and P D risk." *Neurology*, 59(8), pp.1161-1169, 2002.
- [22] S. N. Khandare, M. M. Pillai and N. K. Khandare, "Effect of parsley extract on sublingual gland of naturally aged male mice." *J. Cell and Tissue Research* Vol. 13(2), pp.3761-3764, 2013.
- [23] V. L. Weimer and K.H. Haraguchi, " A potent new mesodermal growth factor from mouse submaxillary gland: A quantitative, comparative study with previously described submaxillary gland growth factor." *Physol chem. Phy.*, (7), pp.7-21,1975.
- [24] W. M. Thomson, J. M. Chalmers, A.J. Spencer, and M. Ketabi, "The Occurence of xerostomia and salivary gland hypofunction in a population based sample of older south Australians spec." *Care Dentist*, (19), pp. 20-23 1999.
- [25] Z. Mohanty, A. D. Singh, A. Dinda, K. K. Talwat, S. Joshi, and S. K. Gupta, "Mechanisms of cardioprotective effect of flavonoids in experimentally induced myocardial infarction." *Pharmacol. Toxicol.* 94(4), pp.184-190,2004.

Author Profile



S. N. Khandare¹ is a postgraduate in Zoology and awarded Ph.D in cell biology (specialization in ageing), from Shivaji University Kolhapur, (Maharashtra). He is assistant Professor of Zoology and has 9 years teaching experience and actively engage in ageing research at Vidnyan Mahavidyalaya, Sangola (affiliated to Solapur University), Dist- Solapur, (Maharashtra), India.



N. K. Khandare² is a postgraduate in Botany and awarded Ph.D in plant pathology, from Shivaji University Kolhapur, (Maharashtra). He is assistant Professor of Botany and has 18 years teaching experience and actively engage in research in disease management and mycorrhizal fungi at Krantisinh Nana Patil College, Walwa (affiliated to Shivaji University, Kolhapur), Dist-Sangli, (Maharashtra), India.